



- |                                  |  |  |
|----------------------------------|--|--|
| Pleistocene and Recent           |  | Alluvium   |
|                                  |  | Alluvial fan deposits                              |
|                                  |  | Landslide debris                                   |
| Late Tertiary(?) and Pleistocene |  | Basaltic Cinder Cone                               |
|                                  |  | Basalt / Undetermined angular unconformity         |
| Miocene                          |  | Sevier River formation unconformity                |
|                                  |  | Dry Hollow formation                               |
| Upper Oligocene                  |  | Bullion Canyon volcanics                           |
|                                  |  | Hornblende Rhyolite                                |
| Paleozoic                        |  | Augite andesite                                    |
|                                  |  | Conglomerate                                       |
|                                  |  | Undifferentiated Upper Paleozoic Sedimentary rocks |
|                                  |  | Coconino ss  |

- Contact  
Dashed where approximately located.
- Fault  
Dashed where approximately located; dotted where concealed or inferred. U-upthrown side, D-downthrown side.
- Strike and dip of beds.
- Unit area outline

**GEOHERMAL DIVISION**  
**LOS ANGELES, CALIFORNIA**  
**COVE FORT-SULPHURDALE UNIT AREA**  
 GEOHERMAL RESERVOIR ASSESSMENT CASE STUDY  
**GEOLOGIC MAP**  
 Plate III  
 MILLARD AND BEAVER COUNTIES, UTAH  
 SCALE: 1" = 4000'  
 JULY, 1976