

# GLOVALLO

USGS Cascades Workshop

22 May 85

Wednesday Morn

Pat Muffler -

Reserve = 1% of accessible resource base  
① Tuseamy, 1978.

- see his paper for all geoth energy
- Spoke of Coc 790 undiscovred reserves  
10,000 x 10<sup>18</sup> joules estimated in  
identified + undisc. convection systems 790°C
- 790 indicates a large undiscovred  
resource base.

Marianne Geffert

- High level silic magma chambers not indheated  
by geophynes
- Lack of surface hydrothermal manifestations
- Bob Smith / Herb Shaw model
  - shallow silic magma chambers of age ad idune
  - single pulse of magma, conductive cooling
  - effects of magmatic reheating isolated by  
hydrothermal convective cooling

- 62 young igneous systems identified
- plot log age vs log volume
- evolved 11 Cascades systems  
Lassen etc. -  
Medicine L  
Shasta

Total calculated heat energy of these 11 total  $3900 \times 10^{18} J$ , which is about equal to Clear Lake system and smaller than Long Valley.

Lassen, Crater Lake, Newberry, Glacier Peak, Meager etc. - identified hydrothermal systems  
In addition few small systems in boundary West Cascades / High Cascades.  
also not in Cascades - not associated of a volcanic system

For Cascades systems, took ratio of hydroth to igneous energy where possible

Glacier Peak 4%

Newberry 11%

Lassen 5%

Hyd Syst known

Baker

Glacier also new known

Rainier Crater L.

Wood Medicine L.


Newberry

Lassen

Igneous

Dal Stanley - Craig Weaver's Talk

- Craig has much updated data base better than what is in literature.



This area N of Rainier is low in geologic potential. Portland volcanics might be on split in plate.

Shallow dipping bench in wash, steeply dipping plate in Oregon.

Highly compressive in wash from St Helens north - lack of basalt.

- electrical data indicates same thing strongly

Cross sections  $\approx W$  - JGR article

Green unit =  $< 5$  km - m conductor --

looks like compression unit to him -

- north Cascades  $> 30$  km then very sensitive

- has a conductor  $\approx$  accreted sediment

normal

$\Rightarrow$  highly compressive nature of  $x$  set in wash

- for comparison, in Oregon,

simple cross sections

deep conductor 0-5 km below 20-25 km

w/ interval 100-500 km unit most of

crust w/ surface perturbation

- regard profile thru Newberry

shows trough under high Cascades, Newberry

shows a buildup

dep of damaging plate steps beneath

Coast Ranges -

- if crustal tectonics controls reaching to surface of intrusions

- electrical sections show  $\approx 5$  km

underthrust unit in wash and also Alps

Compartments.



- underthrust body is conductor
- St Helens seismic zone -- is a strike-slip energy release system

Bob Fuis, USGS

Seismicity under Cascades

↳ kinematics to BOP -

- modelled stresses + gravity simultaneously -
- Geol. context

↳ kinematics - accreted oceanic rocks

↳ nodoch - Sierra Nevada continues beneath them

Cascades are between, in transition zone

Seem to be pulled SOR

Norm MacLeod

- showed side-lighted photo of 1:250K

plastic terrain map -

- Cascades show as buried topography -

- base unit is ash flow 36 my extends Shasta to Eugene -


- Slide K-Ar ages - 687 dates

lots dates 0-2 my -- reflects interest --

compilation by George Walker, Jim Smith -

- high Cascades younger than any

North Cascades separated by NE-trending line from W-O border




- In W Cascades, trend of volc. is  $\perp$  to direction of convergence = NE
- South - continuous ~~the~~ belt: known normal faults, fissure vents, etc indicate extension. Low seismicity by contrast indicates E-W extension.
- about 3000 known vents in Cascades - many lakes are craters. Shoto volc are only a small population.
- distrib of vents has a well defined western limit -- thin line -- not an obvious fault control.
- Silicic rocks - mostly assoc w/ strato volc - few isolated occurrences between.
- almost all among silicic volc rocks in Cascades are park or welder types.
- In other parts of world, andites form hydro syst.
- Rocks between strato volcanoes are not andites
  - mean of 62 andites 53.5%  $\text{SiO}_2$  - - basaltic andites. Hammond got 57.9% @ Indian Heaven for  $n=62$ . So broad volc of Cascades are ~~basalts~~ - basaltic andites (52% is split up basalt, > 57% is andites).

- Blackwell contends that the background gradient is 600°C/km and 100 mW/m<sup>2</sup>.
- MacLeod -- Columbia Plateau "Basalt" is 58% SiO<sub>2</sub>
- Jim Smith - USGS - has been compiling <sup>geologic</sup> data -

Dave Blackwell

- always believed that "active andesites indicate subduction".
- low heat flow on coast assoc. w/ description of heat by slabs -- ~~then~~ changes to high heat flow in High Cascades over 1/2 width 10 km. Anomaly is broad, w/ volcanoes just pinpoints on overall map. -



- Believes that this is real picture, of hydrologic pattern taken out --

- JGR, 1982 paper

- Good - models of magma chambers compared to HF would allow a small chamber that may not be still molten

- pattern of transition to high HF persists in BC. major tectonic sets E of transition, on top of broad anomaly -


- gravity changes in west of axis of Cascades model has been made, etc that show ~~the~~ low density @ depth -- at variance w/ proposed model.

→ what has been detail of rock density studies?

- "HF generated gravity" - did joint modeling of HF's gravity -- gravity low west of Bight could be an edge effect of low- $\rho$ , hot body -- wants low- $\rho$  body under of Cascades.

-- if subduct out body, just get NE-body contour - contours fit up NW as regional gravity -- he subducts at a Cascades on. -

skewed E-W prof



models : oregon top ~~at~~ melt 10km  
washington - top melt 13-15km

- believes in major staging chamber @ 10 km.
- Sierran - scale heat source, w/ volcanoes just above brecciation occurs -

- gravity model - temp induced gravity changes produce only 25% of desired changes -

- why, if there is a shallow chamber, are there few Holocene flows? why <sup>few</sup> shows of protraction?

- uses Hildreth - doesn't feel that there is any evidence for a siberic magma chamber at all - basaltic-andesite is too well spread to indicate a siberic material.

- Blachw -

prob is what is partitioning between basalt that makes it to surface vs ~~to~~ remains in crust -

- believes that more than compression - extension going on in w/o to account for differences - feels there is a basic difference in heat input in these areas.

Eyer - good teleseis data from Canada border to Calif. -

There is a high velocity anomaly at depth - have faster arrivals -


but magma chamber delay could be subberged - he sees no evidence for velocity low @ 10 km -

Blachw has used 3 km for thicknesses of gravity block - unresolvable from p-wave delay -

-- steep HF G gravity is west of volcano - why?

Blockwell model -

up to date version  
is SP pap 15 Heat F



deep circulation must be in deep, high level metabasites or granites / ultramafics that can be part of -- below west case sealed, low grade rocks -

Bob Christensen

Shasta -

each cone has been formed in a short period of time - years to a few thousand years -

- 25 km<sup>3</sup> avalanche 5000 yrs ago destroyed much of cone -- so there was a new history


- 65% SiO<sub>2</sub>, 2-pyrox andesites

- both domes within Shasta and flank domes to 75% SiO<sub>2</sub>

- dacitic pyroclastic flows on all parts of volcano

- calc-alkaline lavas on flanks

olivine tholeiites around it, & not from it.




both long-lived, relative to axes.



Lassen

andesitic flows, pyroclastics  
thick silice flows  
silicic flanky eruptions

- origin of various rock types is by  
crystal fractionation



## Medicine Lake - Lake Dore

Shield volcano above Modoc plateau  
1200 m x 600 km<sup>3</sup> lavas over wide  
area.

Shasta - Med. Lake 1 Cascade Blvd  
45 - 77% silica - bimodal.

most mafic - no obvious evolution  
thru time. In historic time basalt,  
andite, dacite have all erupted.

7 volume silicic eruptions

Glass mtn

Little Glass mtn

Hoffman Dater - 5 vents


Medicine Dater

- no qp evidence for large silicic chamber -  
2 km across resolution of qp feat.
- granitic inclusions occur in rhyolite
- med lake lava come within a few km  
of Klamath Falls - she speculates  
that med lake may supply hot water.
- climatic eruption 600 yrs BP -  
2 phases - rhyolite erupt + rhyolite  
phase also pyroclastic - thick,  
ventric deposits @ 9500 C from  
pyroclastic growth.

Top magma before eruption ~~level~~  
was  $\approx 3$  km

Crater Lake - good area to study  
differentiation and alteration

Newberry -  
calls for underplating of a rhyolite magma  
to heat up mag slightly - observed few  
phenocrysts of feld - - not calcic;  
crystallized -



Rhyolite cholor in W side of  
lake underplating prod ~~to~~ basalts around

Ed Taylor →

3 - sites group

siberia volcanism present

most recent rhyolites 1900 BP


- no reg. chronologic appearance of differentiated  
volcs like indep evolution of new magmas which  
are related due to tectonic events - but over  
a long period of time -

Adams -

only shasta is bigger in vol.  
can less than 200 yrs old, on  
wood base now  $\times 10^3$  years old -

- several basaltic fields full in whole  
fringe -

- fundam. a basaltic magmatic system.  
- system is much bigger than can



- all major siberia eruptions are  $> 100,000$  years old

Jack Southern -

Meager  
~~Caley~~ Caley  
Barabaldi

along  
three N-S eruptive  
center

Basaltic adobe → 2 pyrox adobe → h<sub>2</sub> dactyl


Nathan Green - differentiation red ug  
JUGR public + GSA


Area here, eruptive vents have  
moved north.

- two groups mud uggs - Meager h<sub>2</sub> &  
Pebble Creek

Meager h<sub>2</sub> Noel - displaced  
Pebble Nattoz - meteoric base

art - layered - earth water ? polarization  
suggests dikes as source -





Coley -


underground magmatic cell in basement rocks

- wall springs -
- sunburst anomaly -
- 100° C/cu in of anomaly

- heat source hidden - - gives few springs -

Al Truitt - observes that in peaks of new  
systems, organic alteration - wonders why?  
are there structures that lead gasses to surface?

Fed Syer -




① depth, body anomaly  
becomes azimuth-sensitive -  
- use linear immersion technique to get  
velocity in subsurface - introduces "fuzziness"

Studies: Yellowstone, NHP, etc - dozen or so areas -

Lang Vally, Mars, NHP, COSO -- all seismic centers have shown area larger than 5 km of low velocity @ depth. Velocity contrasts 10-12%

- Hood, Lassen, Shasta - have found no anomalies in Cascades centers - also profile 300 km long across Crater L.





Shasta shows same thing - its a -1 sec residual, meaning high vel. at depth. Believed to assoc w/ subduction.

- inversion shows high vel body 20-200 km depth - 7-10% relative high. when applied crustal corrections, high v even sharper, dips to E?  
-- intep still in process.

There is a 1 sec change across Cascades for 300 sta network -

- cause is a large-scale body - Blackwell's body would give only 1/4 of desired anomaly -

- midlake lake & Newberry are better constrained -

- shows a slab between 10E-20 km beneath Newberry w/ 10% velocity high

- picture similar in mid-lake -

- intep - high veloc due to repeated mafic dikeing

- to improve resolution, asked if DOE funding, worked @ Newberry 10 km x 10 km, 100 m instr grid - explosive sources - 9-10 shots

- high velocity zone @ Newberry is believed due to ring dikes complex -- but need much more intep to solidify ideas

### Card Figure

-  $\rho = 2.43$  g/cm<sup>3</sup> for regional data set, but  $\rho = 2.20$  needed for volcanoes -

- map of  $\lambda > 10$  km

- Gravity lows down axis of High Crater - steep gradient in transition zone -

- subtracted to get residual map.

- Medicine Lake - +24 ugals area -

modeled. results 2.8-2.9 density for causative body. top body 1 km over center to 4 km over edges

- same at Newberry -

- Crater Lake also shows + w/ 2.8 beneath.

- also Lassen - + body. very shallow body.

- Goat rocks - large area of grassy area,  
mag anomaly & gravity almost same,  
so did joint modeling

- She says that worldwide, calderas  
w/ dia  $> 15$  km, gravity lows  
 $< 15$  km gravity highs  
but sign of anomalies depends on country  
rock, etc.

23 May 85

Bob Mariner


- Few Springs - - Newberry = drowned fumaroles - -  
difficult to work with

- Springs on boundary of W & Cascades  
Baker, Glacier Peak, Condit, St Helens -

- Speke about doing a budget study  
of stream above & below spring to get  
Chemistry?


- Macready -  $N_2$  discharge  
Brody -  $CO_2$  discharge

Discharge rate vs temp yields smooth curve




Gas gradient -  $N_2/Ar$  ratios are high  
 use 60-100 -- found up to 300. Means  
 destroy/keep Ar or generate  $N_2$ . Latter  
 means organic matter involved. Also find  
 $CH_4, C_2H_6$

Sorey - models of Hydro Syst




Newberry

- water table 200m outside volcano
- he suggests water table @ 300m to put 260°C bottom hole @ 930m under enough pressure to be liquid



Crater Lake


Hydro npt expressed by elev SO<sub>2</sub>, Cl in lake and high HF in lake by Dave working  
 3.3 HFH in lake bottom -  
water budget




For total h f 2-7X  
 Casdiental 251-5 cfs

Lassen Peak

see also his Stanford paper



each outflow zone (2-3 in the) could supply  $\pm 25$  MW



He concentrations indicate long travel times for this syst

1800C hand on rest of US - but long travel time

Estimated area over which water  
absorbs mg heat flow @ depth  
to produce heat out at springs -

Sprng	conductive m	Avg kwt	Conduct mwt/m <sup>2</sup>
Austin	83	185	410
Proctor	9	85	106
Bellhop/Fol	21	123	122

- See the Flower took USGS to task  
over Williams' HF measurements ~~what~~ in bottom of  
Crater Lake which showed ~~no~~ temp anomalies  
of only few tenths of degree - people are  
now saying that these springs may be disrupted  
by development.

Norm Coe's firm - see paper -

- believes we need to get as close as possible  
to summit area.
- EM sounding work closer to summit.

Adel Zohdy

MT, DC RES, DIGHEM, TDEM, FDEM

In Cascades, low  $\rho$  will be high-temp brines, or altered rocks.

DC RES - 1. prob is very high contact resistance.

- Medicine Lake - 1-2 soundings found near basement
- 2. layout of line in area limited access. Did corrections for being off line.

Med Lake;

1. DIGHEM - got 3 anomalies; med Lake staff  
E of lava flow on E, north spot; Telephone plat.  
900 Hz; anomalies down to 10-30  $\mu$ m -

Newberry

MT, Schlumberger, TDEM



- got low  $\rho$  zone under caldera @ about  
depth of temp increase. To west get  
deep  $\rho$  low west of edge of caldera

Doc Stanley -

MT results @ Newberry

- power layer of 10-20  $\mu$ m, w/ higher to west,  
not closed to E
- maps depth to "alt'd layer" - conductor.  
shallow as 200-300  $\mu$ m -
- depth to chert basement 1500  $\mu$ m





Dave Fitterman  
Transect  $\leftarrow$   $\rightarrow$  @ Newberry

- plot apparent  $\rho$  vs time - layered earth interpretation

[ 60  $\Omega$ m to 450 m ]  
[ 12  $\Omega$ m basement ]

interface corresponds as first  
conductive zone in USGS well -

in caldera, this section is then  
outside caldera, upper layer 400-800m  
of 200-300  $\Omega$ m, w/ 20-70  $\Omega$ m below

Depth to conductor up  
inside caldera 400m  
outside " to 800m

cause conductor not clear, esp outside -

Frank Fischbein

Medicine Lake

TDEM Soundings

## Cascades models Panel

Chris, Al T., Wes, Don W., Pat Welch, Dave B., Mike S. -

Chris.

1. Cascades - subduction controlled volc. arc system  
- w/ regional tectonics quasi-ridge of subd features  
i.e. B&R tectonics on E side, etc.  
- maybe have to think of Bell overlap S of  
Rainier split, look toward north.
2. Geologists seem favor models of small blob like bodies,  
gp's favor large bodies, Chris feels that big  
plutons are probably extent.

Al T.

1. Has lost faith in theory of cold water blankets  
- shield are wrong shape to form gt systems,  
but are narrow column  
- success in Cascades will be w/ cinderas - they conform  
w/ Taupo, other cinderas areas
2. Geochron has not been fully utilized yet.  
- <sup>3</sup>He/He ratios studied, show  $\pm 5 \times$  atmosphere rates,  
shows broad magmatic signature to Cascades  
- geochron not as informative as one would hope -  
are Cascades waters diff from B&R? Amount of SO<sub>2</sub>  
of interest, formed from H<sub>2</sub>S?

## Dove Blackwell

1. Gt systems are where you find them
  - some volcanos may not have gt systems
  - dict tech - can make models for drilling both low & high p, etc -
  - deep circulation systems can occur -- background gradient is 24 B&R
  - counter A.T.'s argument - big volcanos are rather inaccessible compared to Calderas

## Mike Sorey

1. Can form models of what we see from present data -
  - its possible in each setting to have high T, but then flows appear to be very low -- maybe too low for development.
  - caldera lobes prob driven a lot of surface manifestations
  - regional systems may exist -- if gp can't see, it will be very expensive to locate by drilling -
  - look for structure on surface

## Wes Hildreth -

1. Don't know depths of magma storage
  - ages of magmatism - long duration, but cones constructed in short time, often overlapping volcanos
  - lots have large volumes, but each event may be small -- silicic chambers indicated to be small

for connection

[heat  
permeability

- heat sources, no quest

- permeability big probs

- can't quite understand using sketches -- miles of intr contacts

as no evidence of hydrothermal syst -- veering, alt, unmarked

- so if find heat source, search for hydrothermal syst only begun

- from pc grad, many and other systems seem to intrude  
own volc pile, so that dior. grads into volc above, and  
some other structures while thing is unmarked.

- how to develop permeability?

the geyser structures are important, but  
pc's are often shattered areas, <sup>veined</sup> ~~veined~~ in course  
of cooling

can conceivably of such a system more or less confined  
below areas of low permeability and/or cold water removal of  
outflow products -- low permeability exist below alt of

- there is not type of system we can analyze @ usgs, due  
few sketches carry heat, but may be type present under  
Lassen and may be spearhead.

gas geodesic?

rocking in contact

- He believes most rhyolite chambers are  $< 1 \text{ km}$  across
- Ductile could come from chambers too small to be detected g.p.
- much of magma is in pods or dikes -- maybe too much emphasis on big magmas
- Petrograph textures, mixing, inclusions  $\Rightarrow$  complex plumbing -
- can <sup>magm</sup>dike swarms supply gravity anomalies, account for lack of seis anomalies
- can dike swarms form gt systems
- How important are crustal structures vis a vis mantle focusing of intrusion?
- Does crustal extension (either magma/beltwater storage or cause energy descent)

## Don White

1. what's possible of cascades being a perp - An guarding sept.

- if so, how to decide which area is best?
- look @ volcanoes of variety of volc product
- multiple intrusion of small bodies
- magmatic eruption history not good
- want recent fumarolic activity -


2. what is longevity of ht systems:

- Steamboat  $\approx 3 \text{ my}$ , perhaps not continuous -

→ How about electrical anisotropy re. regional structure -- to measure

## Notes

1. Believes we don't have good field feel for actual geoth models - in Cascades -
2. Ned took @ used Bretanish model, which has pump in a bottle, front of ~~activity~~ unit - what about Puse units?
3. Bretanish Lake Suedo
  - BHT prob 160°C
  - aquifer temp 136°C by bottom-reading thermometer
  - Blackwell probe @ bottom
  - George wants to deepen to 4cm
  - Suedo took no samples of aquifer or any other



George pitches a drill hole to the east as Bretanish -- wants to follow aquifer back into cascades

Bob Fournier

1. Saltan Sea well

- DOE has given scientists 6 weeks to come up w/ scheme to get more core -- maybe core all way w/ stem hole
- Sept is mud date may be pushed back
- argument that need large hole to get good fluid sample -- pit to dispose of water is too small to clean out hole anyway -

2. Fluid samples are avail < 300°C for stem holes -- USGS has one; also commercial

- Phil Bethke has suggested sealing down a g/f-z crystal fract d -- it will soak up fluids of bed itself -- do fluid inclusion type work
- for non-artesian diff wells, look off w/ N<sub>2</sub> lift -



→ ask George about regard metamorphic studies.

Alwahel

Brecciated -

- no sig faults
- perm limit to either perm in flows & subvolc material
- w increased depth/metamorphism at TD rocks were brittle and would have maintained fractures if they had been
- So where is permeability?  
as you increase in depth, rock becomes brittle - structural features will provide permeability -  
permeability varies w depth
- to south where Mesozoic is made of white luffs, spec is that permeability would be deeper -

Heat Source

- "Subvolcanic" feature that controls permeability is an intrusion, basaltic andesite, that is presumed to dip east via geophysics - etc

George Priest Meeting

24 May 85

1. Candidate drill sites selected on basis of data available

- Mt Hood
  - lower heat flow
  - study of stratcone is less likely to be useful in developing a model for typical high cascades
- Sautiam Pass area <sup>max</sup>
  - better because data avail, typical of cascades development
  - Breitenbush deep drill hole avail able

2. Talked about DSECC meeting

- Dave B. pissed -
  - unsolicited proposal not in communication of scientific community
  - IRIS succeeded in getting community involved, DSECC failed
  - \$2M of NSF funds to look @ Applause area
  - SAC will pass on merits of proposals
  - subgroup was inquisition part -- panel had own programs proposed -- well written proposals were deemed to be higher rating
  - they have made a presentation to Jim Hayes that was received well -

- no chance to impact FY86 budget
- 1) - some chance to impact FY87 -- money would come in in fall 1987 at soonest  
must be some perseverance
- 2) - continued pressure -- worrying on the Fed budget
  - need broad base of congressional support for sig funds in program at all
  - need concurrence of DoS, ECISAC and/or OBEES community to be rated highly
- 3) OSECC will renew/updates every year, so time for details, but first fact in door counts, too

George talked of caps of Zeolite facies rock sitting on top of higher grade metamorphics in western Cascades that may be fractured and permeable

Pullington - central problem is fluid/permeability at depth - we know there is heat around - deep results Sweden + Newberry show not much fluid/permeability -

George - ~~can~~ Dean would be more in favor of drilling on structure. Then -

There is a COCORP Line thru Siskiyou pass -

Paul Hammond } working to organize a Penrose  
Norm Goldstein } or other workshop to put out a pub.  
to stop on table

Rawley gave/showered George copy of proposal and explained in detail how it was put together, offered help, etc.

John G's outline

Cascades

B, ER

- Theory
- Methodology
- Techniques
- Logistics
- Interpretation
- Integration
- Proofing

autographs — no one really working on it  
 - Keith Berger — took mel Beeson's prog  
 marks Park — good @ area ident / fluid idea  
 Structure — George Walker —

Structure is not well covered in Co program

→ write something on prop — a relationship as applies to Cascades

Two nearby areas of pc observation

- 1) Little N Santiam Dist — townships, potassic alt
- 2) Detroit pluton also attend —

How proposed program would shed light on mineralization.

Fields of Oreg states good contact  
 unless in Oreg cascades we see only tops of  
 plutons of same phases as BC.

how does pc model fit  
 w/ both litho & v. calc  
 are  
 exp on the  
 same  
 kind

V14 no.7  
GSA abstracts 1982 p 463 Pb isotopes - Church  
GS Canada abstracts 1983 v 8 p A12 Pb isotopes - evidence  
of mixing Church

OF 84-0587 selected geoph bibliography of Snake River Plain  
& E Oregon

OF 83-0909 SP Alewherry

OF 83-0177 - gravity principle faults Deanard Peak

OF 83-0466 - " " " Mt Jefferson

OF 84-0512 A/M way E-Central Oreg

OF 84-0205 AM SW Wash NW Oreg

OF 84-0391 A/M W-Cent Oreg

OF 83-0474 mag & geoth res Mt Adams wilderness, Wash

OF 83-0386 principle faults + density contrasts backhole gravity

RRL-3, -4, -5 etc. Harland site

OF 83-0176 principle faults 106 grav sta Goat Rocks wilderness

OF 84-0511

OF 84-0510 } A/M maps

OF 83-0650

JGR 178 p 4447 - Mt/Stanley

MF-1653-6. grav/mag Goat Rocks map

HEAT FLOW OF THE CASCADE RANGE; D. D. Blackwell and J. L. Steele,  
Department of Geological Sciences, Southern Methodist University, Dallas, TX  
75275

The heat flow pattern in the Cascade Range has been studied in quite great detail compared to other island arc terrains and even compared to any other areas of regionally focused heat flow studies in the United States. These heat flow studies have been summarized in very brief form by Blackwell and Steele (1983). A heat flow map of the Cascade Range and adjoining areas taken from that report is shown as Figure 1.

Between 45° and 43° north latitude, there is a 200 km long N-S zone of the Cascade Range which has been most intensely studied and the geophysical characteristics of that area are summarized in idealized form on Figure 2. The characteristics of the heat flow pattern are low heat flow values in the western part of the western Cascade Range and in the Coastal Provinces (averaging 40 mWm<sup>-2</sup>) and exceptionally high values throughout most of the eastern part of the western Cascade Range and the adjoining part of the High Cascade Range (averaging 100 mWm<sup>-2</sup>). The transition from low to high values occurs over a distance of about 20 km.

Between 45° and 47° N latitude heat flow values along the axis of the Cascade Range are lower and average between 70 to 80 mWm<sup>-2</sup>. The data are very sparse north of 47° but some values in the Cascade Range between Snoqualmie Pass and Mt. Rainier suggest values of 40-60 mWm<sup>-2</sup>. Heat flow in the Puget Sound and Vancouver Island regions is considerably less than 40 mWm<sup>-2</sup>. Closely spaced heat flow values in a fjord in British Columbia across the province boundary suggest a heat flow pattern similar to that found in southern Washington and a similar transition zone width of about 20 km. Values in the high heat flow terrain are greater than 60 mWm<sup>-2</sup>. Data south of 43° N

are sparse; the low heat flow coastward is well documented, but the regional heat flow along the Cascade Range axis is not well determined.

The significance of the heat flow data in central Oregon has been discussed in detail by Blackwell et al. (1982). The data require a large heat source with a width of 40 to 60 km at a depth of 7 to 10 km below the surface. Interpretation of the data are permissive of the occurrence of a locally partially molten zone or "magma chamber" throughout this whole region. Confirmation of the significance of the heat flow data is provided by interpretation of gravity data (see Blakeley et al., 1985), depth to Curie point aeromagnetic analyses (Connard et al., 1983), and regional magnetotelluric studies (Stanely, 1982). These results are summarized in Figure 2. Interpretations of the gravity data are permissive of the existence of a body at a depth of between 7 and 10 km with a negative density contrast of  $0.15 \text{ gm/cm}^3$  with respect to the country rock. A density contrast of this magnitude would require at least partial melting within this zone. Further to the north the data are not so easily interpreted and the heat source zone, although it still must be at a depth of approximately 7 to 10 km may have a mean temperature of about  $500^\circ\text{C}$ , or significantly below the solidus of any likely crustal material at those depths (see Figure 3).

Drilling of intermediate depth holes (1-2.6 km deep) confirms the interpretation of regional gradients based on the heat flow studies in the shallower holes (100 to 600 m deep). In particular holes over 2 km deep near Mt. Hood and west of Mt. Jefferson near Breitenbush Hot Springs confirm that regional gradients are in the range of  $60$  to  $70^\circ \text{ C/km}$  to depths of at least 2+ km. The results from the Breitenbush hole further indicate that at least some of the hot springs are related to lateral outflow at relatively shallow depths



from thermal aquifers originating beneath the axis of the High Cascade Range. Therefore only drilling studies which give a true picture of the thermal structure along the axis of the Cascade Range will allow a precise evaluation of the geothermal characteristics and the ultimate crustal significance of the observed high heat flow.

Blakely, R.J., Jackens, R.C., Simpson, R.W., and Couch, R.W., 1985. Tectonic setting of The Southern Cascade Range as interpreted from its magnetic and gravity fields. Geol. Soc. Am. Bull., 96, 43-48.

Connard, G., Couch, R., and Gemperle, M., 1983. Analysis of aeromagnetic measurements from The Cascade Range in Central Oregon. Geophysics, 48, 376-390.

Leaver, D.S., Mooney, W.D. and Kohler, W.M., 1984. A seismic refraction study of the Oregon Cascades. J. Geophys. Res., 89, 3121-3134.

Stanley, W.D., 1981. A regional magnetotelluric survey of the Cascade Mountains region. U.S. Geol. Surv. Open-file Rept., 24pp.

Blackwell, D.D., and Steele, J.L., 1983. A summary of heat flow studies in the Cascade Range. Geothermal Resources Council Trans., 7, 233-236.

Blackwell, D.D., Bowen, R.G., Hull, D.A., Riccio, J., and Steele, J.L., 1982. Heat flow, arc volcanism and subduction in Northern Oregon, Jour. Geophys. Res., 87, 8735-8754.

Blackwell




Figure 1. Heat flow map of the Cascade Range and adjoining areas (1 ucal/cm<sup>2</sup> sec = 41.84 mWm<sup>-2</sup>).





Figure 2. Includes data from Leaver et al. (1984), Connard et al. (1983) and Stanley (1982).

CENTRAL WASHINGTON CASCADE RANGE



CENTRAL OREGON CASCADE RANGE

Figure 3. Intrusive models of high heat flow in the Cascade Range

WRIGHT

Undiscovered Geothermal Resources of the United States

by

L. J. Patrick Muffler  
U. S. Geological Survey  
Menlo Park, CA 94025

Resource assessment can be defined as the broad-based estimation of future supplies of minerals and fuels. This assessment includes not only the quantities that could be produced under present economic conditions, but also the quantities not yet discovered or that might be produced with improved technology or under different economic conditions. Resource assessment requires (1) estimation of the amount of a given material in a specified part of the Earth's crust and (2) estimation of the fraction of that material that might be recovered and used under assumed economic, legal, and technological conditions. Geothermal resource assessment is thus primarily the estimation of the thermal energy in the ground, referenced to mean annual temperature, coupled with an estimation of the amount of this energy that might be extracted economically and legally at some reasonable future time.

Estimates of geothermal resources of the United States vary by as much as six orders of magnitude, from a high of 10 billion megawatt-centuries (MWe-c) to several low estimates that cluster around 10,000 MWe-c (Muffler, 1973). Part of the reason for the discrepancy in published geothermal resource estimates lies in a confusion among the terms resource base, reserve, and resource. These terms come from the oil and minerals industries and have been defined as follows:

Resource base: "\*\*\* the sum total of a mineral raw material present in the earth's crust within a given geographic area \*\*\* whether its existence is known or unknown and regardless of cost considerations and of technological feasibility of extraction." (Netschert, 1958; Schurr and Netschert, 1960, p. 297)

Reserve: "quantities of minerals \*\*\* that can be reasonably assumed to exist and which are producible with existing technology and under present economic conditions." (Flawn, 1966, p. 10).

Resource: "\*\*\* that part of the resource base (including reserves) which seems likely to become available given certain technologic and economic conditions." (Netschert, 1958; Schurr and Netschert, 1960, p. 297)

For geothermal energy, geothermal resource base was defined by Muffler and Cataldi (1979) as all the heat in the Earth's crust beneath a specific area, measured from local mean annual temperature. Accessible resource base was defined as the thermal energy at depths shallow enough to be tapped by drilling in the foreseeable future. The geothermal resource was defined as that fraction of the accessible resource base that might be extracted economically and legally at some reasonable future time. The geothermal resource can be divided into economic and subeconomic categories depending on present-day economics. This logic can be displayed as the vertical axis of a McKelvey diagram; this diagram defines the terms resource and reserve.

In the United States, geothermal resource assessment is the responsibility of the U.S. Geological Survey. In meeting this responsibility, the USGS has during the past decade produced three geothermal resource assessments (White and Williams, 1975; Muffler, 1979a; Reed, 1983a). Collectively, these national resource assessments evaluated geothermal energy in five categories:

1. Regional conduction-dominated regimes
2. Igneous-related geothermal systems
3. Geopressured-geothermal energy
4. Hydrothermal convection systems at temperatures  $>90^{\circ}\text{C}$ .
5. Geothermal regimes at temperatures  $90^{\circ}\text{C}$ .

Geothermal resources were estimated only for categories 3, 4, and 5. Energy calculations made for categories 1 and 2 are not resource estimates, but are useful in giving a conceptual framework for the occurrence of geopressured and hydrothermal convection systems. In particular, for igneous-related geothermal systems, the thermal energy still remaining in silicic intrusions and adjacent country rock was calculated by conductive cooling models using estimates of the size and age of intrusions (Smith and Shaw, 1979). This calculation assumes that cooling of the igneous body by hydrothermal convection was offset by the effects of magmatic preheating and additions of magma after the assumed time of emplacement (Smith and Shaw, 1975).

The figures estimated and calculated in the 1978 USGS geothermal resource assessment of the United States suggest that the energy in igneous-related systems to a depth of 10 km is 100 times greater than the energy in all identified and undiscovered hydrothermal convection systems to a depth of 3 km, and nearly 1000 times greater than the energy in all hydrothermal convection systems identified to date. From this comparison it can be inferred that very large amounts of geothermal energy yet remain to be found. The major geothermal questions facing us for the United States in general and for the Cascade Range in particular are:

- What is the true value of this igneous-related energy?
- What is the distribution of this energy among magma, hot dry rock, and hydrothermal convection systems
- Where is all this energy located?

## References

- Flawn, P. T., 1966, Mineral resources: Chicago, Rand McNally and Co., 406 p.
- Muffler, L. J. P., ed., 1979a, Assessment of geothermal resources of the United States--1978: U. S. Geol. Survey Circ. 790, 163 p.
- Muffler, L. J. P., and Cataldi, R., 1978, Methods for regional assessment of geothermal resources: Geothermics, v. 7, no. 2-4, p. 53-89.
- Netschert, B. C., 1958, The future supply of oil and gas: Baltimore, Johns Hopkins University Press, 134 p.
- Reed, M. J., ed., 1983a, Assessment of low-temperature geothermal resources of the United States--1982: U.S. Geological Survey Circular 892, 73 p.
- Schurr, S. H., and Netschert, B. C., 1960, Energy in the American economy, 1850-1975: Baltimore, Johns Hopkins Press, 774 p.
- Smith, R. L., and Shaw, H. R., 1975, Igneous-related geothermal systems, in White, D. E., and Williams, D. L., eds., 1975, Assessment of geothermal resources of the United States--1975: U. S. Geol. Survey Circular 726, p. 58-83.
- Smith, R. L., and Shaw, H. R., 1979, Igneous-related geothermal systems, in Muffler, L. J. P., ed., Assessment of geothermal resources of the United States--1978: U. S. Geol. Survey Circ. 790, p. 12-17.
- White, D. E., and Williams, D. L., eds., 1975, Assessment of geothermal resources of the United States--1975: U. S. Geol. Survey Circular 726, 105 p.

PREVIOUS ESTIMATES BY THE U.S. GEOLOGICAL SURVEY  
OF GEOTHERMAL RESOURCES OF THE CASCADE RANGE

by Marianne Guffanti  
U.S. Geological Survey, Reston, VA 22092

Introduction

As part of a second assessment by the U.S. Geological Survey of geothermal resources of the United States (Muffler, 1979), estimates were made of thermal energy in igneous-related geothermal systems and in hydrothermal convection systems with reservoir temperatures  $\geq 90^{\circ}\text{C}$  in the Cascade Range. The assessment is based on data available July 1, 1978, and is a revision of the first national geothermal resource assessment (White and Williams, 1975). This paper summarizes results of the 1978 assessment for the Cascade Range, in order to provide a starting point for a revised regional assessment of the Cascades. Medicine Lake volcano, California, and Newberry volcano, Oregon, are considered as part of the Cascades in this discussion, although in the 1978 assessment they were not included in that province.

Igneous-related systems

Thermal energy stored in magma, solidified pluton, hot country rock and associated hydrothermal convection systems to a depth of 10 km was estimated for young igneous centers in the western U.S. (Smith and Shaw, 1979). This thermal energy was calculated using a conductive cooling model based on estimates of volumes and ages of presumed shallow silicic magma chambers. The primary assumption of the model is that for a given system a single pulse of magma was emplaced instantaneously and cooled conductively from the time represented by the youngest age of the system. Another assumption is that effects of magmatic preheating and resupply to the system are balanced by hydrothermal cooling. As a best guess, it was estimated that about half of igneous-related geothermal

energy exists as magma and nearly that much as solidified intrusion and hot surrounding rock, with only a few percent expressed by hydrothermal convection systems (Muffler, 1981).

The thermal energy calculated for 11 igneous-related systems in the Cascades is approximately  $3900 \times 10^{18}$  joules (table 1), an amount approximately equal to that calculated for the Clear Lake volcanic center near the The Geysers geothermal field in California. The 11 Cascades systems represent about 8% of the igneous-related energy in the United States, excluding the thermal energy in the very large Yellowstone-Island Park system. Six igneous-related systems in the Cascades were not evaluated for thermal energy -- Mt. Hood, Mt. McLoughlin, Mt. Jefferson, Mt. Baker, Mt. Rainier, and Mt. Adams -- because adequate age or size data were not available or the magma chamber was thought to be too small and/or at a depth greater than 10 km. Of the 11 evaluated systems, eight appear to be at least partially molten and thus may serve as geothermal heat sources. These are Lassen Peak, Medicine Lake, and Shasta in California; Crater Lake, Newberry, and South Sister in Oregon; Glacier Peak and Mt. St. Helens in Washington.

One aspect of a revised assessment of the Cascade geothermal regime might involve a refinement of igneous-related thermal energy estimates. Based on detailed geologic mapping and geochronological and geophysical studies at several volcanic centers in the Cascades, new age and size data may permit revision of existing estimates, as well as calculation of thermal energy at some of the previously unevaluated systems. However, the assumptions of the conductive cooling model may be poorly approximated in the Cascade Range. Although the Cascades has high regional heat flow and extensive Quaternary volcanism, substantial magma chambers in the upper crust have not been detected by various geophysical techniques (Iyer, 1984). What, then, are the geometries



and depths of crustal magma reservoirs underlying Cascades volcanic centers? At what rates is magma supplied to these igneous centers of predominantly intermediate composition and how is it stored in the upper crust? How efficient is hydrothermal cooling of Cascades magma bodies? What is the division of thermal energy among magma, solidified intrusion, and hydrothermal systems in the Cascades? What is the nature of magmatic supply and storage to areas between the major stratovolcanoes?

### Hydrothermal systems

Fourteen hydrothermal systems in the Cascades (including Newberry volcano) were estimated in 1978 to contain  $84 \times 10^{18}$  joules (table 2), about 5% of the hydrothermal energy of all identified systems in the United States. The recoverable thermal energy (resource) in 3 high-temperature systems -- Morgan-Growler H.S. near Lassen Peak, California; Gamma Hot Springs near Glacier Peak, Washington; and Newberry volcano, Oregon -- is equivalent to 883 MW of electricity for 30 years (as a comparison, the current electrical generating capacity at The Geysers is about 1400 MW.) Because the Lassen system and Ohanapecosh Hot Springs are in National Parks and thus not available for exploitation, the resource was not calculated there. Beneficial or usable heat of  $0.5 \times 10^{18}$  joules was estimated for hydrothermal systems with reservoir temperatures between  $90^{\circ}$  and  $150^{\circ}\text{C}$ .

Determination of the portion of igneous-related energy that is expressed hydrothermally for various Cascades volcanoes is a first step in characterizing the relationship between magmatic heat sources and hydrothermal systems. Unfortunately, assessment data for this purpose are incomplete in the Cascades. Only three systems have both igneous and hydrothermal estimates to permit calculation of the hydrothermal-to-igneous ratio: Lassen, 5%, Glacier Peak, 4%, and

Newberry, 11%. These values are fairly high compared to some other systems such as Long Valley, Valles, and Yellowstone, where the hydrothermal portion is 1-3%. The overall hydrothermal-to-igneous ratio in the Cascade Range is 2%, excluding hydrothermal systems not obviously associated with igneous systems.

Undiscovered thermal energy in hydrothermal systems in the Cascades is estimated to be very large amount, approximately 20 times the amount stored in identified systems. The undiscovered estimate is simply a multiple of the identified component, chosen subjectively on the basis of the favorable geologic setting. The transport of heat into the upper crust by magma movement, as evidenced by the abundance of young volcanic rocks, and the occurrence of hydrothermal systems along the Range, coupled with the possibility of groundwater masking, suggest that a large resource may exist. To confirm the large undiscovered estimate would require some combination of the following:

- 1) Previous thermal energy estimates be revised upward by applying new temperature and volume data. For example, drilling results at Newberry volcano in Oregon (Sammel, 1981) indicate a higher subsurface temperature there than previously thought, 265°C compared to 230°C. Also, the hot-spring systems along the Western/High Cascades boundary may represent larger, hotter systems at depth. Nominal volumes of 3.3 cu km that were assigned to these systems should be re-evaluated in light of new data;
- 2) Previously unidentified hydrothermal systems be evaluated. For example, no hydrothermal resources have yet been quantified at Medicine Lake or Crater Lake. The discovery of new Lassen-type systems is unlikely, but more "Newberries" might be uncovered beneath the rain curtain at other volcanic edifices. And we need to determine what, if any, concealed hydrothermal systems exist at depth between the major stratovolcanoes.

Certainly, the educated guess made in 1978 should be re-evaluated in light of a refined magmatic-hydrothermal model for the Cascade Range. What distinguishes the volcanic centers that have hydrothermal systems from those that do not? To what extent does masking of hydrothermal systems by cool groundwater occur? Where are convenient structures for localization of hydrothermal activity? Can we apply information gained at a few type hydrothermal reservoirs to systems elsewhere in the Cascades? If hydrothermal reservoirs are masked and not expressed at the surface, how do we obtain the necessary data on temperatures and sizes for energy estimates, short of extensive drilling? If the Cascade Range does not have large hydrothermal resources, why not?

#### References Cited

- Iyer, H. M., 1984, Geophysical evidence for the locations, shapes, and sizes, and internal structures of magma chambers beneath regions of Quaternary volcanism: *Phil. Trans. Royal Society of London*, v. 310, p. 473-510.
- Brook, C. A., Mariner, R. H., Mabey, D. R., Swanson, J. R., Guffanti, M., and Muffler, L. J. P., 1979, Hydrothermal convection systems with reservoir temperatures  $> 90^{\circ}\text{C}$ , in Muffler, L. J. P., ed., *Assessment of geothermal resources of the United States -- 1978*: U.S. Geol. Survey Circular 790, p. 18-85.
- Muffler, L. J. P., 1979, ed., *Assessment of geothermal resources of the United States -- 1978*: U.S. Geol. Survey Circular 790, 163p.
- Muffler, L. J. P., 1981, Geothermal resource assessment, in Rybach, L. and Muffler, L. J. P., eds., *Geothermal Systems and Case Histories*: John Wiley and Sons, p. 181-198.
- Sammel, E. A., 1981, Results of test drilling at Newberry volcano, Oregon: *Geothermal Resources Council Bulletin*, v. 10, n. 11, p. 3-8.
- Smith, R. L., and Shaw, H. R., 1979, Igneous-related geothermal systems, in Muffler, L. J. P., ed., *Assessment of geothermal resources of the United States -- 1978*: U.S. Geol. Survey Circular 790, p. 12-17.
- Smith, R. L., Shaw, H. R., Luedke, R. G., and Russell, S. L., 1978, Comprehensive tables giving physical data and thermal estimates for young igneous systems of the United States: U.S. Geol. Survey Open-File Report 78-925, 27p.

Table 1. Thermal energy calculated for young igneous systems of the Cascade Range. Data are from Smith and others (1978).

	Age, Years	Chamber Area, km <sup>2</sup>	Chamber Volume, km <sup>3</sup>	Heat Now 10 <sup>18</sup> J
CA Lassen Peak	61	80	400	960
CA Medicine Lake	<1000	64-74	300	724
CA Mt. Shasta	<200; 5-9500 ?	50	300	724
OR Crater Lake	6600	50	>320	>770
OR Newberry	1300, >6600	32	100	240
OR South Sister	<2000	30-40	100	240
OR Melvin-3 Crks	400,000 ?	10	40	76
OR Cappy Burn	2,500,000 ?	8	40	26
OR Bearwallow	<1,000,000 ?	8	30	41
WA Glacier Peak	12,000; <70,000	5	12.5	35
WA Mt. St. Helens	Active	5	>12.5	>35
TOTAL				3900 x 10 <sup>18</sup> J

Table 2. Thermal energy estimates of hydrothermal systems in the Cascade Range. Data are from Brook and others (1979). Values of temperature, volume, and stored energy are mean values; associated standard deviations are omitted here for simplicity of presentation but are given in Brook and others (1979).

	Temp. °C	Volume km <sup>3</sup>	Stored Energy 10 <sup>18</sup> J	Recoverable Energy 10 <sup>18</sup> J	Electricity or Beneficial Heat
<u>&gt;150°C</u>					
CA Lassen	237	71	42	Nat. Park	--
CA Morgan-Growler	217	8.3	4.5	1.1	116
OR Newberry	230	47	27	6.9	740
WA Gamma H.S.	165	3.3	1.4	0.3	27
					883 MW-30 years
<u>90°-150°C</u>					
CA Big Bend H.S.	116	3.3	0.91	0.23	0.055
OR Mt. Hood area	122	3.3	0.96	0.24	0.058
OR Carey H.S.	104	3.3	0.80	0.20	0.048
OR Breitenbush H.S.	125	3.3	0.99	0.25	0.059
OR Belknap H.S.	113	3.3	0.88	0.22	0.053
OR Foley H.S.	99	3.3	0.76	0.19	0.046
OR McCredie H.S.	91	3.3	0.68	0.17	0.041
OR Umpqua H.S.	112	3.3	0.87	0.22	0.052
WA Baker H.S.	134	3.3	1.07	0.27	0.064
WA Ohanapecosh H.S.	127	3.3	1.00	Nat. Park	--
TOTAL					84 x 10 <sup>18</sup> J
					0.5 x 10 <sup>18</sup> J

SEGMENTATION OF THE JUAN DE FUCA PLATE:  
A FIRST-ORDER GEOTHERMAL ASSESSMENT OF THE CASCADE RANGE?

Craig S. Weaver  
U. S. Geological Survey

If the assessment of the geothermal potential of the Cascade Range were a straight-forward application of existing techniques, one could argue rather persuasively that the gross plate tectonic setting, although interesting, was likely of marginal use in the assessment process. That is, the real world of geothermal assessments deals with the nature of magmatic/hydrothermal systems in the shallow crust (0-10 km) in localized areas (horizontal scales of at most a few 10's of km). The location of the seismic Benioff zone, the dip of the Juan de Fuca plate, and the like, although ultimately responsible for the volcanism observed in the Cascade Range, would not be the primary input (and in reality probably not considered at all) to the geothermal assessment.

On the other hand, during the past seven years, it has become increasingly apparent that the Cascade Range does not represent a simple assessment problem that is amenable to application of existing methodology. Part of the problem with attempting to understand the geothermal potential of the range is the tendency to treat the entire range as a single province- in essence we have failed to recognize the significant scale changes that exist throughout the range. As an example, it is widely known that the Cascade Range is the volcanic arc for the subduction zone created by the convergence between the Juan de Fuca and the North American plates. On the other hand, the tectonic setting of the range changes markedly from north to south, and these changes are in large part the result of changes in the geometry of the Juan de Fuca plate and the interaction with the North American plate. In this paper I suggest that the changes in the tectonic framework, when coupled with the distribution of late Cenozoic and Quaternary volcanism in the Cascade Range, provide a simple first order

geothermal assessment of the Cascade Range from the American border south to about Crater Lake: 1) north of Mount Rainier- zero potential, 2) Mount Hood to Mount Rainier- slight potential (but probably not much above zero), and 3) Mount Hood to Crater Lake- moderate potential (perhaps locally some areas with high potential). Although I have not yet modeled the southern portion of the range (Crater Lake to Lassen Peak), this entire region probably should have moderate potential (again with some local areas of high potential).

The changes in the tectonic framework that allows this first order geothermal assessment is postulated to be the result of the segmentation of the Juan de Fuca plate as it subducts beneath the North American plate. (In this report, segmentation refers to the change in dip of the Juan de Fuca plate landward of the coastal thrust zone). In subduction zones, the tectonic framework is determined in part by both geometry (including convergence rate, direction, dip angle) and the type of interaction between the plates (aseismic- without large magnitude thrust earthquakes or seismic). The lack of Benioff zone earthquakes along most of the subduction zone has made it difficult to determine those aspects of plate geometry related to the dip of the Juan de Fuca plate, and there remains considerable uncertainty as to the nature of the plate interaction.

Although northwestern Washington is the simplest portion of the Cascade Range from the point of view of geothermal assessments (zero), it does provide the only area in the range where there is a tie between the geometry of the Juan de Fuca plate as deduced from the occurrence of Benioff zone earthquakes, and the distribution of crustal seismicity and volcanism in the overlying North American plate. The lack of Cenozoic volcanism probably is the result of three factors: 1) the shallow dip of the Juan de Fuca plate and the thick crust beneath the Olympic Mountains, 2) the thickness of the continental lithosphere in the North Cascades, and 3) the continuous nature of the slab as it subducts to the

northeast. The thick crust beneath the Olympic Mountains allows a significantly increased area for the direct interaction between the Juan de Fuca and the North American plates, and this results in horizontal compression dominating the fore arc. The stresses from the plate interaction produce high rates of seismicity in the Puget Sound basin, and work to minimize volcanism in the arc. The thickness of the lithosphere beneath the North Cascades and the continuous nature of the slab may work together to inhibit volcanism by acting as either a heat sink or preventing hot asthenosphere from coming in contact with the subducting slab. As a consequence there is little late Cenozoic volcanism within the Cascade Range, and the geothermal potential is zero.

Between Mount Rainier and Mount Hood, the geometry of the Juan de Fuca plate is not as well constrained as to the north, but the few Benioff zone earthquakes suggest that the dip of the subducting plate (landward of the thrust zone) is greater than to the north. The sudden end of the Benioff zone earthquakes has been used to suggest that the Juan de Fuca plate may not be continuous south of Mount Rainier, but has broken off as a result of repeat thrust events on the interface with the North American plate. This breakoff of the Juan de Fuca plate may decrease internal gravitational stresses to the point that seismicity ceases within the slab. However, because the slab (oceanward of the proposed break) is still strongly coupled with the overlying North American plate, horizontal compression again dominates the arc region. Earthquake focal mechanisms are largely strike-slip on vertical fault planes, in good agreement with the horizontal stress direction expected from gross plate convergence.

The poor geothermal prospects in most of this area reflect the lack of volcanism observed at the surface and the compressional nature of the crustal stresses. As an example, Mount St. Helens is located at a left-stepping offset along a regional strike-slip fault zone characterized by right-lateral motion. It is

likely that the resulting compression at Mount St. Helens has limited the development of secondary vents away from the immediate vicinity of the cone and is responsible for the very small scale of the upper crustal magmatic system ( $\sim 10 \text{ km}^3$  in the upper 7-10 km). My designation of a slight potential for this region reflects the existence of strike-slip faulting (which may have a favorable opening geometry in some local areas) and the increase in late Cenozoic volcanism southward toward Mount Hood.

Finally, south of Mount Hood, seismic data offer essentially no constraints on the geothermal potential of the Cascades. As with the Mount Rainier to Mount Hood segment of the Juan de Fuca plate, the lack of Benioff zone earthquakes beneath Oregon may suggest that the plate is broken near the coastal thrust zone. Further, the crust beneath the Oregon Coast Range is significantly thinner than that beneath the Olympic Mountains in Washington, as a consequence the contact between the Juan de Fuca and the North American plates is reduced. Superficially, this geometry might lead to extensional processes dominating the arc region, and the volume of late Cenozoic volcanics found in the Oregon Cascades is consistent with this proposed extension. Simply because the crust is under extension rather than compression, the geothermal potential should be higher than to the north in Washington. In addition, local areas with abnormally high rates of crustal extension may exist, and these areas would be expected to have the highest geothermal potential.



## Volcanism in the Cascade Range

N. S. MacLeod and D. A. Swanson, Cascades Volcano Observatory, U. S. Geological Survey, Vancouver, WA.

Volcanism in the Cascade Range started about 36 m.y. ago and produced such widespread units as the Ohanapecosh Formation in Washington and the Little Butte Volcanics in Oregon. Since then, volcanic activity has been more or less continuous on a time scale of a few million years, although certain segments of the range have apparently had relatively short periods of increased or decreased activity. For example, in Washington few volcanic rocks were erupted 17-15 m.y. ago at the time of maximum outpouring of the Columbia River Basalt Group from fissure vents farther east. In Oregon, on the other hand, that period produced voluminous flows and tuffs of the Sardinia Formation. Despite contrary claims, evidence is weak or lacking for regional volcanic episodicity in the Cascades, a concept whose testing is beset with sampling problems.

There is no compelling evidence that volcanism formed a narrow linear chain before about 3-4 m.y. ago. For example, in central Oregon vents of Oligocene age extend from the western Cascades more than 100 km to central Oregon. Even since 3 m.y. ago, volcanism has not been confined to a narrow belt; Mount Adams is 50 km east of Mount St. Helens, Newberry Volcano 60 km east of the Cascade crest, and Medicine Lake 55 km east of Mount Shasta. Minor vents are even more widely scattered. Nonetheless, most eruptions during the last few million years in Oregon have occurred along a very linear chain forming the High Cascades. This chain is also exemplified by north-trending linear fissures, normal faults, and alignments of cinder cones and widely spaced stratovolcanoes.

Most volcanism in the Cascades during the last few million years has been south of Mount Rainier. Only a few volcanoes occur farther north, and all of these fall along a northwest-trending linear belt between Glacier Peak, Washington, and Meager Mountain, British Columbia, that is strongly discordant to the chain that trends south from Mount Rainier. The Glacier Peak-Meager Mountain belt projects southeastward through several known Tertiary structures to the western Snake River Plain and thus may be partly controlled by an old crustal weakness. The belt is nearly perpendicular to the direction of plate convergence, and moderately high seismicity outboard of the belt has been interpreted to define a Benioff zone with low dip. In contrast, the area west of the Oregon and northern California Cascades has low seismicity except along spreading ridges and related transform faults, and no Benioff zone. In that area the Cascade crest is oblique to the convergence direction, normal faults and fissure vents indicate east-west extension, and volcanism is much more voluminous than farther north.

More than 2,000 volcanoes of late Pliocene or Quaternary age have been identified in the Cascade Range; many others lie buried by younger flows. About 50 percent of the volcanoes are in Oregon, 35 percent in California, and 15 percent in Washington and British Columbia. The vast majority of these volcanoes are monogenetic cinder cones or fissure vents that erupted basalt or basaltic andesite flows. In some parts of the range the cinder cones and fissure vents are randomly dispersed, but in other parts they occur in distinct clusters. One such cluster of late Holocene basalt and basaltic-andesite vents occurs between the Three Sisters and Three Fingered Jack in Oregon, and another somewhat older cluster occurs in the Indian Heaven area between Mount St. Helens and Mount Adams, Washington.

Small stratovolcanoes and relatively steep-sided lava shields form perhaps 5 to 10 percent of the total vents. Many appear to be of basaltic-

andesite composition, with lesser basalt and andesite, and they rise as much as 500 m above the surrounding lava fields. Most of these volcanoes are probably short-lived, perhaps being formed in a few years to as much as a few thousand years.

Large stratovolcanoes form only about 1 percent of the total volcano population, yet their imposing size leads to popular identification of the range with stratovolcanoes. The largest volcanoes, Mount Rainier in Washington and Mount Shasta in California, lie near the ends of the chain of greatest volcanism in the Cascades. The deeply eroded remains of several large Pliocene and Pleistocene stratovolcanoes have been identified along the chain; few stratovolcanoes persist long as imposing edifices because of erosion, especially by glaciers.

Clusters of stratovolcanoes occur at Three Sisters, Crater Lake, and near Lassen Peak. These clusters represent long-lived volcanic systems that have produced relatively large volumes of intermediate and silicic magma; the cluster at Crater Lake was largely destroyed by the cataclysmic eruption that formed Crater Lake caldera.

Large shield-shaped volcanoes or volcanic fields lie in a north-trending belt along the east side of the Cascade Range. These include Newberry volcano in Oregon and Medicine Lake volcano in northern California; Lassen Peak may lie along this trend. Similar volcanoes or volcanic fields several million years older occur along this same alignment at Yamsay Mountain and Bald Mountain, Oregon, and in the Simcoe, Washington, area. All of these eastern vent areas have erupted silicic as well as mafic lava, and several have summit calderas.

The petrochemistry of many of the stratovolcanoes has been studied, but most of the lava fields between stratovolcanoes have received no such attention and few generalizations can be made about them. Basaltic andesite and

basalt appear to dominate in areas between stratovolcanoes, and even some of the stratovolcanoes consist mainly of basaltic andesite rather than andesite; calc-alkaline rocks dominate, but tholeiitic types also occur. Andesite is locally abundant, especially on some of the stratovolcanoes, and in a few areas dacite, rhyodacite, and rhyolite are common. With the possible exception of youthful Mount St. Helens, most areas in which silicic rocks are abundant appear to be products of long-lived magmatic systems from which eruptions have occurred over periods of hundreds of thousands of years. These include Lassen Peak, Shasta, and Medicine Lake in California, Crater Lake, Newberry Volcano, and the Three Sisters-Broken Top highland in Oregon, possibly Glacier Peak in Washington, and Mount Garibaldi and Meager Mountain in British Columbia. Local centers of silicic volcanism that have generated voluminous pyroclastic flows occur at Crater Lake, Medicine Lake, and Newberry Volcano, all of which have calderas; a caldera may lie buried in the Three Sisters-Broken Top highland, judging by the distribution of ash-flow tuffs around that area. This paucity of Quaternary calderas suggests that large high-level silicic magma bodies are not common.

Geothermal resources have been identified at Meager Mountain, B. C., high temperatures have been encountered in shallow holes on Newberry volcano, and several of the volcanoes have fumaroles. Geothermal resources are presumed to be most likely to occur at long-lived volcanic systems that have produced silicic rocks. Some andesite and basaltic-andesite stratovolcanoes may have geothermal resources, although drilling at Mount Hood was not encouraging. A background of relatively high heat flow in the Cascades as a consequence of long-lived volcanism throughout most of the chain may indicate an extensive deep resource.

## Quaternary Volcanism in the Southernmost Cascade Range, California

Michael A. Clynne  
U. S. Geological Survey  
345 Middlefield Road MS 910  
Menlo Park, CA 94025

Volcanism in the southernmost Cascade Range can be characterized on two scales. On a regional scale volcanism is basaltic to andesitic. Individual volcanoes have small volumes and relatively short lifetimes. Many eruptive centers coalesce to form the crest of the Cascade Range. Individual volcanoes range from monogenetic basalt and basaltic andesite cinder cones to larger lava cone and shield volcanoes of basaltic to andesitic composition. Rocks with greater than approximately 60 per cent  $\text{SiO}_2$  are sparse. Superimposed on the regional mafic volcanism are long-lived, larger volume volcanic centers that have produced eruption products spanning a wide range of composition from mafic andesite to rhyodacite or rhyolite.

Each volcanic center consists of a large andesitic composite cone and flanking silicic domes and flows. Three of these volcanic centers younger than about 3 Ma have been recognized in the Lassen area. Each has had a similar history, consisting of three stages: (1) an initial cone-building period of mafic andesite and andesite lava flows and pyroclastics; (2) a later cone-building period characterized by thick andesite and silicic andesite lava flows, and (3) silicic volcanism flanking the main cone. The silicic magma chamber of the third stage provides a heat source for development of a hydrothermal system within the core of the main cone. Alteration of permeable rocks of the cone facilitates increased glacial and fluvial erosion of the central area of the volcano. The result is preservation of a resistant rim of the thick later cone-building lava flows and flanking silicic rocks surrounding a depression in the altered and eroded core of the composite cone.

The two older volcanic centers have reached this stage, and their hydrothermal systems are extinct. The younger Lassen volcanic center (LVC) however, hosts active silicic volcanism and a well developed hydrothermal system, including the thermal features in Lassen Volcanic National Park (LVNP) and Mill Canyon.

The LVC developed in three stages. Stages I and II produced the Brokeoff Volcano, an  $80 \text{ km}^3$  andesitic stratocone. Stage I deposits consist of olivine-augite and olivine-hypersthene-augite andesite lava flows and pyroclastic rocks erupted from a central vent during the period from about 0.7 to 0.55 Ma. Stage I culminated in eruption of a small volume of hornblende-pyroxene dacite. An interval of erosion lasting 0.1 Ma followed stage I. During stage II, which lasted 0.1 Ma, thick flows of porphyritic augite-hypersthene silicic andesite, generally lacking interbedded pyroclastic material, were erupted. Volcanism then shifted to the north and east and became more silicic. Eruption of a minimum of  $50 \text{ km}^3$  of rhyolitic magma at 0.35 Ma as air-fall tephra and ash flows initiated stage III. This eruption probably produced a collapse caldera now filled by a dacite domefield. The dacite domefield consists of two groups of rocks totaling on the order of  $30\text{-}50 \text{ km}^3$ . During the period from 0.25 to 0.2 Ma pyroxene-hornblende dacite magma produced a group of six domes and flows. Hornblende-biotite rhyodacite has been erupted as domes, lava flows and pyroclastic flows in at least ten episodes over the past 0.1 Ma. At least six times during the past 0.1 m.y. hybrid andesite totaling  $10 \text{ km}^3$  and consisting of thoroughly mixed mafic and silicic magma has been erupted from the margins of the dacite domefield.

Porphyritic andesite and dacite with high  $\text{Al}_2\text{O}_3$ , low  $\text{TiO}_2$ , and medium  $\text{K}_2\text{O}$  contents and  $\text{FeO/MgO}$  ratios of 1.5-2.0 are the most abundant rock types in the LVC. Early mafic andesite, late rhyodacite, and hybrid andesite are subordinate in abundance. Rocks of the LVC resemble other calc-alkaline.

volcanic rocks emplaced on a continental margin overlying silicic crust. Harker variation diagrams of the major element chemistry of rocks from the LVC show smooth trends from 55-73 per cent  $\text{SiO}_2$ . The overall evolution is from mafic to silicic. However the evolution is not strictly sequential (see fig). Sparse isotope data suggest that crustal interaction is not significant in the development of the silicic rocks. However, partial melting of young mafic crust may play a role in the origin of the silicic magmas at LVC. Smooth geochemical trends, increasing silica content with time, and isotope data all suggest the origin of LVC magmas by crystal fractionation of a mafic parent. The long time span of activity at LVC and the presence of a hydrothermal system imply the presence of an evolving magma chamber. However, the current conceptual model of the magmatic system of the LVC as outlined below is poorly constrained and somewhat speculative.

The Lassen volcanic center differs from other Cascade volcanoes by having a larger volume of silicic rocks. The stress regime in the southern Cascades is moderate east-west extension, which favors retention of mafic magma in the crust and development of zoned systems with large volumes of derivative silicic magma. Systematic geochemical variation of the rocks of the Lassen volcanic center and the mineralogic and geochemical homogeneity of the youngest rocks suggest derivation from a single source. The magma system of the Lassen volcanic center can be envisaged as an evolving pluton-sized body of magma in the middle crust (10-20 km depth). The pattern of young vents suggests that an 5-8 km diameter magma body underlies the northwestern corner of LVNP. The young volcanic vents and hydrothermal system may indicate the position of a cupola on the magma chamber, which may be as shallow as 6 km beneath Lassen Peak. The hybrid andesites are produced by mixing of mafic magma intruded into silicic magma and may indicate a sill-like projection of


the magma chamber beneath the central plateau of LVNP. Alternatively the hybrid andesites may be erupted by tapping the main chamber at depth.

The upper portion of the chamber contains crystal-rich rhyodacite, and has varied little in composition over at least the last 50,000 years. The system is probably zoned to more mafic compositions at depth. Regional basalt provides heat and material input to maintain the system in the partially molten state. There is abundant evidence for the interaction of mafic and silicic magma. A small volatile-rich cap has periodically developed at the top of the magma chamber but is removed by venting before increasing to substantial volume. Assuming that the vertical extent of the magma chamber is 10 km results in volume estimates of 250-500 km<sup>3</sup>.

Despite the volcanologic and petrologic evidence of a magma system beneath the LVC, geophysical evidence for the presence of a magma chamber is lacking. A 25 km oval, -50 mgal gravity anomaly is centered on the dacite domefield and central plateau (hybrid andesites) of LVC. The gravity low is probably an expression of light volcanic rocks in the near surface and the presence of Quaternary plutonic rocks below the volcanic field.

Teleseismic and seismic refraction studies have failed to show the presence of a magma chamber beneath LVC. However the resolution of these studies is such that magma chambers smaller than 5 km in diameter could not be seen. A compilation of regional seismicity reveals an area lacking crustal earthquakes that corresponds to the gravity low. Future geophysical studies will have to be of higher resolution to detect the magmatic system of LVC.





Evolution of the Lassen Volcanic Center

## The Mount Shasta Magmatic system

Robert L. Christiansen

U.S. Geological Survey

Menlo Park, California

The still-active compound stratocone of Mount Shasta, 14,162 feet high and more than half a million years old, has grown mainly during several distinct episodes of cone building from single central vents, each of which was active only briefly--possibly only a few hundred years (Christiansen and Miller, 1976; Christiansen and others, 1977; Christiansen, 1982). Four such cone-building episodes younger than about 250,000 years have been recognized, two of them Holocene; major cone building was separated by longer, predominantly erosional periods during which smaller volumes of lava were added to the cones. In addition, an older but similar edifice at the site of Mount Shasta was largely destroyed by a huge volcanic sector avalanche about 300,000 years ago (Crandell and others, 1984). The oldest exposed rocks of Mount Shasta, on its west flank and having a K/Ar age of about 590,000 years (G. B. Dalrymple, written commun. 1979), probably are a remnant of that earlier edifice.

The bulk of the stratocone consists of silicic andesites to mafic dacites, probably averaging just over 60% SiO<sub>2</sub>. More or less uniform silicic 2-pyroxene andesites constitute most of each cone of the compound volcano, probably aggregating about 80-85% of the whole. More silicic rocks, including hornblende-bearing andesites,

dacites (mainly 63-66% SiO<sub>2</sub>), and rare rhyolites, occur principally as domes or relatively small lava flows, erupted both within the earlier central craters and on the flanks of the edifice. Significant volumes of dacitic pumice occur largely as fallout to the northeast. Numerous pyroclastic flows were emplaced during early Holocene time, especially on the west flank of the volcano (Miller, 1978); older pyroclastic flows occur on all sectors. In the past few thousand years the system has erupted every few hundred years (Miller, 1980).

Mount Shasta also has erupted relatively small volumes of more femic magma, generally as calc-alkalic basalts to mafic andesites, from cinder-cone or small-shield vents on its flanks. The highest such vent was noted by Anderson (1974) on the south side at about 9,000 feet elevation. No magmas similar to regional high-alumina olivine tholeiites (Hart and others, 1984) are known to have vented from Mount Shasta. Preliminary petrologic studies suggest that much of the range of compositions constituting the bulk of the volcano may have evolved by fractional crystallization and mixing of variously evolved magmas from a calc-alkalic basaltic parent like some of those erupted from Mount Shasta's flank vents (Baker and others, 1984).

Both the andesite-dacite stratocone of Mount Shasta and the more or less bimodal shield volcano of Medicine Lake, probably at least a million years old (Donnelley-Nolan, this volume), are voluminous long-lived systems near the margins of a large negative gravity anomaly. This gravity low extends across the High Cascade axis,

Mount Shasta lying distinctly on the west and Medicine Lake on the east side of the nearly north-south axis at the ends of a linear system of young vents that trends east-northeast, suggesting a locally anomalous stress-field control. Between the Klamath and McCloud Rivers, the Pliocene to Quaternary volcanoes of the main Cascade axis are predominantly rather simple mafic-andesite shields. Within this sector of the chain, substantial volumes of silicic andesites to dacites have been produced only near the intersection of this main axis with the Shasta-Medicine Lake trend.

Taken together, the foregoing relationships suggest a tentative view of the Mount Shasta magmatic system. The areally extensive gravity anomaly, local stress-field control of adjacent volcanic systems, and the systematic relation of two major long-lived and highly evolved systems to the main Cascade axis suggest that the Shasta-Medicine Lake zone represents an anomaly in this sector of the Cascade subduction zone, perhaps related to excess production of felsic magma in the mantle. Growth of the Mount Shasta system in this excess-production region during a few rapid increments may represent distinct upper-crustal emplacements of magma that had evolved at deeper levels to silicic pyroxene andesite from parental compositions like the common mafic calc-alkalic lavas of the region. During the intervening much longer times between these major emplacement events continued mantle magmatism probably drives relatively slow evolution of the Shasta magmas in upper-crustal chambers by open-system differentiation processes, including fractional crystallization accompanied by wallrock assimilation and mixing of variously evolved magmatic batches.

It is during these extended periods of upper-crustal magmatic evolution, during which dacitic and even rhyolitic magmas are produced, that any extensive hydrothermal convection might develop in the vicinity of the magmatic system. Based on both the slow eruptive rate and the lack of major cone building during the past several thousand years and on the abundance of dacites among its Holocene volcanic products, Mount Shasta may now be in such a magmatic state. No evidence is known for the existence of a cryptic hydrothermal system beneath Mount Shasta, but the high downward flux of cold water through the volcanic edifice from its heavy winter snowpack and extensive summer melting could mask hydrothermal convection at shallow crustal depths.

#### References cited

- Anderson, A. T., Jr., 1974, Evidence for a picritic, volatile-rich magma beneath Mt. Shasta, California: *J. Petrol.*, v. 15, p. 243-267.
- Baker, M. B., Grove, T. L., and Price, R. C., 1984, Petrogenesis of Mt. Shasta, N. California andesites by fractional crystallization and mixing: *Geol. Soc. Am. Abstr. Progr.*, v. 16, p. 435.
- Christiansen, R. L., 1982, Volcanic hazard potential in the California Cascades, in Martin, R. J., and Davis, J. F., eds., Status of volcanic prediction and emergency response capabilities in volcanic hazard zones of California: *Calif. Div. Mines Geol. Spec. Pub.* 63, p. 43-65.
- Christiansen, R. L., and Miller, C. D., 1976, Volcanic evolution of Mt. Shasta, California: *Geol. Soc. Am. Abstr. Progr.*, v. 8, p. 360-361.
- Christiansen, R. L., and others, 1977, Resource appraisal of the Mt. Shasta Wilderness Study Area, Siskiyou County, California.: *U.S. Geol. Surv. Open-file Rep.* 77-250, 53 p.

Crandell, D. R., Miller, C. D., Glicken, H. X., Christiansen, R. L., and Newhall, C. G., 1984, Catastrophic debris avalanche from ancestral Mount Shasta volcano, California: *Geology*, v. 12, p. 143-146.

Hart, W. K., Aronson, J. L., and Mertzman, S. A., 1984, Areal distribution and age of low-K, high-alumina olivine tholeiite magmatism in the northwestern Great Basin: *Geol. Soc. Am. Bull.*, v. 95, p. 186-195.

Miller, C. D., 1978, Holocene pyroclastic-flow deposits from Shastina and Black Butte, west of Mt. Shasta, California: *U.S. Geol. Surv. J. Res.*, v. 6, p. 611-623.

Miller, C. D., 1980, Potential hazards from future eruptions in the vicinity of Mt. Shasta volcano, northern California: *U.S. Geol. Surv. Bull.* 1503, 43 p.

## GEOHERMAL POTENTIAL OF MEDICINE LAKE VOLCANO

Julie M. Donnelly-Nolan, U. S. Geological Survey

Mail Stop 910, 345 Middlefield Road, Menlo Park CA 94025

Medicine Lake Volcano (MLV) lies east of the main Cascades trend about 50 km ENE of Mt. Shasta. The volume of MLV is estimated to be 600 km<sup>3</sup>, larger even than Mt. Shasta which is the largest of the Cascade stratovolcanoes. Several lines of reasoning suggest that MLV has geothermal potential, including the suggestion by Eichelberger (1981) that a large silicic magma chamber underlies the volcano. Several late Holocene silicic units are present across the top of the volcano. Two of these, the rhyolite of Glass Mountain on the east side of the central caldera and the rhyolite of Little Glass Mountain and Crater Glass flows 16 km to the west, have nearly identical major- and trace-element chemistries. The close similarity of composition suggests that a single large silicic magma body is the source for these two units. Heiken (1978) explained the Glass Mountain-Little Glass Mountain chemical similarity by proposing tapping of a central small magma body by ring dikes. Other silicic lavas have erupted throughout the history of the volcano although the total volume of such eruptions appears to be relatively small. The earliest such lava is a rhyodacite ash flow about 1.25 million years old. Growth of the volcano apparently followed eruption of large volumes of high-alumina basalt that form the principal rock type of the Modoc Plateau. No obvious pattern can be discerned in the chemical evolution of the volcano except that higher silica eruptions occur on the higher parts of the volcano

while primitive high-alumina basalts have continued to erupt around the lower flanks. MLV consists of these parental tholeiitic basalts plus calc-alkaline derivatives. The average composition of MLV is probably andesitic, and andesitic lavas are abundant especially on the upper slopes of the volcano. Anderson (1941) suggested that the caldera probably formed by eruption of voluminous andesite flows. Andesitic lavas at MLV are typically crystal-poor and fountain-fed, both characteristics suggesting that they were very hot when they erupted. Most of the volcano's lavas are crystal-poor suggesting relatively high temperatures or short residence times before eruption. Rather than a single large magma body, this may suggest small, short-lived bodies of magma that erupt relatively soon after they form.

Geophysical evidence provides no support for a large silicic magma body although a large mafic intrusion can be inferred from seismic refraction and gravity studies. Electrical and teleseismic studies are not definitive about the presence of magma at depth; small bodies less than 2 km across could be present. Intrusion of primitive basalt appears to be intruded into shallow crustal levels at the volcano, followed by fractional crystallization and assimilation, resulting in small, relatively silicic magma bodies. Small bodies of rhyolitic magma were probably produced by fractional crystallization of andesitic magma (Grove and Donnelly-Nolan, 1983). Given similar bodies of andesitic magma, derivative rhyolites of nearly identical composition might be produced by the same set of processes. Thus, nearly identical rhyolitic products such as those seen at Glass Mountain and Little Glass Mountain might be produced. Both lavas contain a variety of inclusions ranging from granitic (probable country rock) to andesitic (liquid when incorporated) to cumulate hornblende gabbros. The variety and abundance of these inclusions argues



against the presence of a single large silicic magma chamber. The granites would have been quickly digested and the cumulate hornblende gabbros should represent material off the bottom of the chamber. Also, none of these young silicic lavas are truly high-silica rhyolites. None contains more than 74% SiO<sub>2</sub>, compared to the older rhyolites at MLV that contain 75-77% SiO<sub>2</sub> and are inclusion-free. The younger rhyolites may have erupted after disruption by intruded mafic magma before the silicic magma body had completed its evolution.

Structurally, MLV lies in an E-W extensional environment. It is located at the intersection of a major lineament trending ENE across the Cascades and the westernmost N-S trending normal faults of basin and range type. One of these faults forms Gillem Bluff at the northern edge of the volcano. Farther south, this fault is buried by younger lavas of MLV, but alignments of vents as well as the relatively abrupt eastern topographic margin of MLV strongly suggest that the fault bends SSE and continues S at least 35 km through the vents for the Glass Mountain flow and domes. A strong NE structural trend is manifested high on the NW side of the volcano by open ground cracks, the Little Glass Mountain-Crater Glass flows and domes, and late Holocene basaltic eruptions. A similar direction is evident at a set of Holocene andesitic pit craters on the SE side of the caldera. This evidence together with the E-W extension and dominant N-S alignment of vents over 50 km on MLV, the ring faults of the volcano's caldera, and other alignments of vents and open ground cracks, provide evidence of fracture permeability at depth.

Is there enough fluid for power production? Although springs are rare on the volcano, very large volumes of cold water emerge at the southern edge of MLV lavas to form the Fall River, suggesting that adequate sources of water are available. Whether water is stored and heated directly under MLV is

unknown. Hot springs are absent, but this does not preclude the existence of hot water at depth.

Whether or not a large silicic magma body exists under MLV, the volcano still has the potential to be a major geothermal resource. Sufficient heat could be supplied by continued intrusion of mafic magma into shallow crustal levels. Numerous Holocene vents and intermittent rhyolitic volcanism for at least 1.25 m.y. suggest that MLV is a prime target for geothermal energy exploration. Several companies are actively engaged in exploration. One deep hole was drilled during the summer of 1984 and more are planned. Results of the drilling by private companies are proprietary but the amount of interest shown suggests that the companies believe there is geothermal potential at MLV.

Outflow of hot water from MLV might also be possible, particularly in the dominant N-S structural direction. Only 30-50 km N of MLV, geothermal fluids in and near Klamath Falls, Oregon, are used for low-temperature applications such as space heating. Klamath Falls lies equidistant between MLV and Crater Lake Volcano, at the lowest point in the Klamath graben that lies between these two major young volcanoes. The Klamath Falls geothermal system, unrelated to any recent volcanism at the surface, may be at least partly supplied by hot water from MLV.

#### References

Anderson, C. A., 1941, Volcanoes of the Medicine Lake Highland, California:

University of California Publications, Bulletin of the Department of Geological Sciences, v. 25, no. 7, p. 347-422.

Eichelberger, J. C., 1981, Mechanism of magma mixing at Glass Mountain,

Medicine Lake Highland Volcano, California: U. S. Geological Survey  
Circular 838, p. 183-189.

Grove, T. L., and Donnelly-Nolan, J. M., 1983, Role of amphibole in the  
differentiation history of Medicine Lake Highland lavas, EOS, v. 64, no.  
45, p. 900.

Heiken, Grant, 1978, Plinian-type eruptions in the Medicine Lake Highland,  
California, and the nature of the underlying magma: Journal of  
Volcanology and Geothermal Research, v. 4, p. 375-402.

## Mount Mazama and Crater Lake Caldera

Charles R. Bacon  
U.S. Geological Survey  
345 Middlefield Road  
Menlo Park, California 94025

Geologic mapping and K-Ar dating (Bacon and Lanphere, 1983) provide the data to define the eruptive history of Mount Mazama (Bacon, 1983). Events that led up to the catastrophic, caldera-forming eruption have been reconstructed through detailed field studies of late products (Bacon, 1983, 1985), including paleomagnetic determinations (Champion, 1983). Chemical analyses of rocks from all periods in the growth of Mount Mazama and microprobe analyses of phenocrysts in late products (C. R. Bacon and T. H. Druitt, unpub. data, 1985) help to suggest models for the magmatic system beneath the volcano.

Mount Mazama was composed of overlapping short-lived shields and stratocones and thick lava flows, constructed west and north of a voluminous field of rhyodacite domes and flows. The rhyodacite evidently was extruded in a few pulses, probably of short duration, at various times between ~750 and ~500 ka (thousand years B.P.). It seems likely that basalt lies beneath the rhyodacite, the thickness of which is unknown.

The oldest exposed stratovolcanoes of the Mount Mazama cluster are Phantom Cone and Mount Scott, ~420-395 ka. These are overlapped by andesitic and dacitic lavas erupted ~355-230 ka. Mafic andesitic shields were built near Cloudcap ~290-220 ka and near Llao Rock ~185-110 ka. Andesitic to dacitic lavas were erupted near Llao Rock ~130-110 ka, followed by rapid growth of a complex andesitic stratocone near Hillman Peak ~70 ka. Shortly after the Hillman Peak cone was constructed, dacite pyroclastic flows descended the southwest flank of Mount Mazama and dacite airfall pumice was erupted from vents near Llao Rock and Cloudcap. Provisional Fe-Ti oxide equilibration

temperatures for the three sets of dacitic pyroclastic rocks are ~930, 990, and 960°C, respectively. Dacite lava flows, one of which is dated at ~70 ka, were associated with the pyroclastic activity. Although there is no evidence preserved for caldera formation, several km<sup>3</sup> of dacite magma must have been erupted. Following these eruptions, the existing evidence suggests that Mount Mazama reverted to producing only andesite for ~20 ka.

About 50 ka the dacitic Watchman flow was erupted from a vent now exposed in the west caldera wall, adjacent to the south side of Hillman Peak. Like many earlier dacite and silicic andesite flows of Mount Mazama, the Watchman flow contains inclusions of andesite that were largely molten when incorporated into the host lava. Their presence indicates that the underlying magmatic system was compositionally zoned, containing magma at least as mafic as andesite that was incompletely mixed into the more silicic magma before eruption. Similar inclusions are present in dacitic blocks found in monolithologic avalanche breccias or lithic-pyroclastic-flow deposits on the south and southwest flanks of Mount Mazama. These deposits were formed by collapse of lava domes near the summit of the volcano ~40 ka. Stratigraphic evidence and K-Ar dating indicate that andesite lava flows on the north and west caldera rims were erupted between ~50 and 40 ka. Known younger precaldere lavas were almost exclusively silicic in composition.

A small hornblende rhyodacite dome on the north caldera rim was emplaced ~30 ka. Similar lava domes of the Sharp Peak group northeast of the caldera may be of similar age. Williams Crater, formerly "Forgotten Crater", west of Hillman Peak may be also approximately the same age. Williams Crater and a vent about 1 km to the west erupted basalt and caused the emplacement of an extensive lava flow and several domes of commingled andesite and dacite. Thus, Williams Crater probably marks the west margin of the silicic magma reservoir at this time. The Redcloud Cliff flow, a thick hornblende

rhyodacite flow on the east caldera wall, evidently is latest Pleistocene in age, and may be as old as the ~30 ka dome on the north rim. There is no evidence preserved for any volcanic activity after eruption of the above units and before Holocene volcanism began.

About 7,000 ka the shallow chamber began to leak significant volumes of rhyodacitic magma in the form of three lava flows: Liao Rock, Grouse Hill, and Cleetwood. Other rhyodacitic flows may have been present within the area that collapsed to form the caldera. Paleomagnetic evidence suggests that the Liao Rock and Grouse Hill flows were erupted in short succession. About 140 yr. later, the Cleetwood flow was erupted. The bulk composition (on a volatile-free basis) and Fe-Ti oxide phenocryst equilibration temperature (~940°C) of this lava are virtually identical with those of the pumice of the climactic eruption, except that the lava had a much lower water content upon eruption. Surprisingly, the climactic eruption began before the Cleetwood flow had cooled. Although most, if not all, of the precursory lava eruptions began with an explosive phase, flow of magma to the surface apparently was sufficiently sluggish to allow the magma to de-gas during ascent so that conduits eventually were plugged with viscous rhyodacite. Only the Liao flow was preceded by a voluminous pyroclastic phase, but this spread tephra over parts of several western states. Loss of several km<sup>3</sup> of magma from the top of the chamber over a comparatively short period to form the lava flows and related pyroclastic rocks may have depressurized the system, without accompanying loss of volatile pressure, so that vapor saturation occurred at greater depth than before and the climactic eruption could be sustained.

The climactic eruption took place in two steps: (1) a single-vent phase and (2) a ring-vent phase. The first phase began with eruption of the widespread airfall deposit from a high Plinian column, which later collapsed to produce pyroclastic flows as the vent became enlarged. The pyroclastic

flows deposited the Wineglass Welded Tuff in valleys on the north and east of the caldera, its distribution being limited by the summit of Mount Mazama which lay to the south of the vent. Phase (1) ended when the caldera began to collapse, initiating phase (2) which produced much more mobile pyroclastic flows from higher eruption columns. These flows deposited nonwelded to partly welded tuff in the valleys all around Mount Mazama and left a lag deposit of lithic breccia near the rim and on high ground, including even the summit of Mount Scott. These deposits show spectacular compositional zonation of homogeneous rhyodacite ( $70-71\% \text{SiO}_2$ ) below andesite that decreases fairly regularly in silica content with stratigraphic height ( $\sim 62-51\% \text{SiO}_2$ ). Equilibration temperatures for Fe-Ti oxide phenocrysts in the silicic part of the eruption group tightly near  $\sim 950^\circ\text{C}$  and, like other "climactic" rocks, are more oxidized than the  $\sim 70$  ka dacites. Over  $50 \text{ km}^3$  of magma were erupted in the climactic eruption, most being rhyodacite. Although the exact depth is unknown, the existence of the caldera suggests that the top of the magma chamber must have been only a few km below the surface.

A provisional model of the magmatic system beneath early Holocene Mount Mazama would have relatively differentiated (up to  $\sim 72\% \text{SiO}_2$ ) and cooler ( $\geq 870^\circ\text{C}$ ) porphyritic magma overlying well-mixed, convecting rhyodacite magma, which was in turn underlain by zoned, crystal-rich andesite. Presumably, basalt underlay the entire reservoir, because basaltic eruptions from monogenetic cones and small shields took place beyond and on the lower flanks of Mount Mazama throughout its history. Such a long-lived crustal magmatic system must have had hydrothermal phenomena associated with it. The uppermost deposits on the caldera rim were formed by phreatic explosions that took place when this hydrothermal system was disrupted by caldera collapse. Postcaldera volcanism, consisting of andesitic activity outboard of rhyodacitic eruptions, suggests that the remains of the old magmatic

system--or a new one--are still viable. High heat flow in the floor of Crater Lake, elevated chloride concentration in the lake water, and the well mixed nature of the lake suggest that hydrothermal convection continues beneath the lake floor (Williams and Von Herzen, 1983). Conceivably; lateral flow of thermal waters might produce geothermal reservoirs beyond the boundaries of Crater Lake National Park where they might be utilized for extraction of geothermal energy.

Beyond the potential for geothermal resources, Crater Lake caldera provides a natural laboratory in which to study not only the evolution of a crustal magma body, but also hydrothermal alteration within an andesitic stratovolcano. Alteration of the older rocks of the caldera walls is ubiquitous, and the lithic breccias of the climactic eruption contain partly-fused plutonic rocks and highly altered fragments from deep within Mount Mazama. These may offer clues to the precaldern hydrothermal history of this long-lived volcanic system.

#### References

- Bacon, C. R., 1983, Eruptive history of Mount Mazama and Crater Lake caldera, Cascade Range, U. S. A.: J. Volcanol. Geotherm. Res. 18:57-115.
- Bacon, C. R., 1985, Implications of silicic vent patterns for the presence of large crustal magma chambers: J. Geophys. Res. 90:in press.
- Bacon, C. R., and Lanphere, M. A., 1983, K-Ar chronology of eruptions at Mount Mazama before the formation of Crater Lake Caldera: EOS 64 no. 45:879.
- Champion, D. E., 1983, Paleomagnetic "dating" of Holocene volcanic activity at Mount Mazama, Crater Lake, Oregon: Geol. Soc. Am. Abstr. Prog. 15 no. 5:331.
- Williams, D. L., and Von Herzen, R. P., 1983, On the terrestrial heat flow and physical limnology of Crater Lake, Oregon: J. Geophys. Res. 88:1094-1104.



The Magmatic System of Newberry Volcano, Oregon

N. S. MacLeod, Cascades Volcano Observatory, Vancouver, WA

D. R. Sherrod, U.S. Geological Survey, Menlo Park, CA

Newberry Volcano, 60 km east of the crest of the Cascade Range in central Oregon, has been active for more than 500,000 yr and erupted most recently about 1350 yr ago. The surfaces of its flanks are formed mostly of basaltic-andesite and basalt flows, which issued from hundreds of cinder cones and fissure vents. Widespread ash-flow tuffs exposed on some flanks probably underlie the basaltic flows; dacite to rhyolite domes and flows are common on the flanks. The summit caldera is filled by rhyolitic ash-flow and air-fall tuff, by rhyodacite and rhyolite domes and flows, and by a few mafic flows and vent deposits. Drill cores show that fragmental debris fills the caldera to about 500 m below the surface; it overlies rhyodacite to basalt flows that extend to at least 932 m. The caldera resulted from several periods of collapse during voluminous eruptions that formed the ash-flow tuffs on the flanks.

The most recent eruption cycle was 10,000 to 1350  $^{14}\text{C}$  yr ago. Widespread basaltic-andesite flows on the flanks, estimated to be about 10,000 to 8000 yr old, are approximately coeval with several rhyolitic flows and domes in the caldera and on the upper southeast flank. Rhyolitic eruptions in the caldera produced domes, flows, and tephra deposits between 6845  $^{14}\text{C}$  yr (approximate age of Mazama ash, a regional stratigraphic marker) and 6100  $^{14}\text{C}$  yr ago. Numerous basaltic-andesite

flows issued from vents on the northwest and southeast flank about 6100  $^{14}\text{C}$  yr ago; eruptions since then have been rhyolitic. Eruptions 4000-3000 yr ago produced obsidian flows and pumice deposits, and 1350 yr ago air-fall tuff, ash-flow tuff, and an obsidian flow were erupted. Although 10,000-8000 yr and 6100 yr basaltic flows are common on the flanks, they are nearly absent in the caldera; those that are present were erupted from fissures on the caldera walls. Mixed basalt-rhyodacite lavas that erupted from near the top of the east wall suggest that basaltic magma may have intercepted silicic magma beneath the caldera.

Analysed rhyolitic rocks younger than about 10,000 yr are chemically similar. They are readily distinguished from abundant older silicic rocks at Newberry by their relatively evolved major- and trace-element composition, especially their higher Rb/Sr ratio ( 2, compared to 1 in older rhyolitic rocks). Also, even though the range in composition is quite small, successively younger rocks show a progressively more evolved composition. The young rhyolites are all very low in phenocryst content and successively younger rocks have fewer phenocrysts; the youngest obsidian flow contains less than 0.01 percent phenocrysts. This suggests that the rhyolites were erupted from a magma chamber whose upper part was near the liquidus and that the tapped part of the magma did not crystallize during a much longer period than that since the most recent eruption. We attribute the decrease in phenocryst content to increasing water content in the magma or to influxes of heat owing to underplating of a silicic magma chamber by hot basaltic magma 10,000-8000 and 6100 yr ago. The basaltic flows erupted during the last 10,000 yr are presumed to be from a source much deeper than the inferred silicic magma body.

The silicic body beneath Newberry (whether still magma or a solidified pluton) appears to be a single body, because the chemical compositions and petrography of the young rhyolites are so similar. The restriction of the vents for the young rhyolites to the eastern part of the caldera and the absence of mafic vents there (a "shadow effect") suggest that the magma body was centered under the eastern half of the caldera. A 932-m-deep hole drilled by the USGS in the caldera encountered temperatures as high as 265°C and high heat flow, such as would exist above a magma body or a hot but solidified intrusion. Hydrothermal convection beneath the drill hole, however, could extend an unknown distance downward to a deep hot body. If magma is still present, the body apparently is small, deep, or complex in shape inasmuch as it has not been definitely detected by geophysical methods.

## THREE SISTERS TO MOUNT JEFFERSON: MAGMATIC SYSTEMS AS HEAT SOURCES

U.S.G.S. Workshop on Geothermal Resources of the Cascade Range

E.M. Taylor, Oregon State University, May 22, 1985

A summary of central High Cascade volcanism, magmatism, and tectonism between Three Sisters and Mount Jefferson, as related to sources of geothermal heat, should include their pre-Pleistocene antecedents because modern heat flow is probably influenced by structural features produced prior to development of the modern High Cascades and by the long-continued evolution of a few isolated silicic magma systems in a mafic volcanic province. The most complete records of early High Cascade volcanism are found in the Deschutes Formation on the east flank of the range and in correlative deposits exposed in the Western Cascades. An episode of crustal tension and production of associated mafic lavas was initiated over a broad part of the eastern Cascades, 8-10 m.a. A combined tholeiitic and calc-alkaline volcanic arc was constructed in the position of the modern High Cascade axis. Shields and composite cones of basalt and basaltic andesite predominated. However, many of the volcanoes produced silicic lavas; in the distal deposits there is a record of hundreds of discrete eruptions of air-fall pumice and pyroclastic flows ranging in composition from mafic andesites to rhyolite. Tensional stresses in the crust apparently increased to a maximal condition by 4-5 m.y., at which time the early High Cascade volcanic arc subsided along discontinuous north-south normal faults, leading by intermittent stages to a broad, graben-like depression of the volcanic

axis. Explosive volcanism was enhanced at this time and was characterized by the appearance of andesitic ash-flow tuffs and abundant, inhomogeneous, mixed magmas of diverse compositions. Much of the volcanoclastic output was probably trapped within segments of the subsiding graben.

After 3 m.a. the intensity of volcanism decreased and the magmas brought to the surface were almost entirely basalt and basaltic andesite. Consequently, the modern central High Cascade Range is a broad platform of mafic volcanoes ranging from large composite cones and shields such as North Sister, Mount Washington, and Three Fingered Jack, to small cones and cone groups with associated lava flows such as Belknap Crater, Sand Mountain, and South Cinder Peak. Late High Cascade silicic volcanism has been restricted to the vicinity of Three Sisters and Mount Jefferson. The Three Sisters volcanic group has been intermittently active for several  $10^5$  years and has produced basalt, basaltic andesite, andesite, dacite, and rhyodacite in the form of lavas, domes, and volcanoclastic deposits representing three clearly distinct magmatic series. All of these compositional types have been erupted during Holocene time; the most recent example is the 1900-year-old rhyodacite of Rock Mesa at the southwest base of South Sister. The development of Mount Jefferson is less well known; it is a composite cone of basaltic andesite surmounted by younger andesite lavas and dacite domes. The most recent eruptions produced dacitic air-fall pumice and pyroclastic-flow deposits that appear to have issued from vents on the northeast.

flank, while late Pleistocene glaciers were retreating.

The central High Cascade province has been a locus of intrusion by mafic dikes, sills, and plugs during the last 9 million years; this has probably contributed to a regional background of geothermal heat. Perturbations in this background are to be anticipated where hydrothermal circulation is influenced by buried graben faults and thick accumulations of volcanoclastic material. Of particular significance in this context is the prolonged localization of silicic volcanism in the vicinity of Three Sisters and Broken Top. Ash-flow tuffs were produced from source volcanoes near the sites of modern Three Sisters 4-5 m.a. Later, an extensive highland of silicic domes was constructed and several late High Cascade ash-flow sheets spread as far east as Bend. The modern Three Sisters group is the latest manifestation of a long-lived source that has been contributing silicic magmas to the shallow crust. It would be surprising indeed if young, hot, plutonic bodies were not present beneath this area.

MOUNT ADAMS: Eruptive history of an andesite-dacite stratovolcano at the focus of a fundamentally basaltic volcanic field.

Wes HILDRETH and Judy FIERSTEIN

U.S. Geological Survey, Menlo Park, CA 94025

Mount Adams (3742 m) is a late Quaternary stratovolcano 50 km due E of Mount St. Helens, 75 km SSE of Mount Rainier, and 90 km NNE of Mount Hood. At 350 km<sup>3</sup>, its volume is exceeded in the High Cascades only by Mount Shasta, but the Simcoe, Newberry, and Medicine Lake back-arc systems are all larger. Mount Adams stands upon the Cascade crest and is drained radially by many streams fed by glaciers radiating from its summit icecap. Only 2.5% of its 650-km<sup>2</sup> area is glacier-mantled today, but as much as 90% was ice-covered in latest Pleistocene time.

There have been no recorded eruptions of Mount Adams and probably just one in the past 3500 years, but at least 7 Holocene eruptions and the persistence of sulfurous summit fumaroles indicate that the volcano remains potentially active. The 7 Holocene eruptions on the stratocone took place at flank vents 2000-2500 m in elevation and produced a wide range of compositions, 49-61% SiO<sub>2</sub>.

Nearly all of the high cone above 2300 m in elevation was constructed during latest Pleistocene time, probably between 20 and 10 Ka, explaining the abundance of late-glacial till and the scarcity of older till. Products of this eruptive episode range from 54 to 62% SiO<sub>2</sub> on the main cone. Contemporaneous and younger peripheral vents yielded lavas and scoriae in the range 48-57% SiO<sub>2</sub>, and, along with Mount Adams, they define a recently active N-S eruptive alignment 40 km long and only 5 km wide. Basalts within this zone are extremely varied compositionally, containing, e.g., 0.16 to 1.6%

K<sub>2</sub>O at 48-48.5% SiO<sub>2</sub>.

Beneath the young edifice, deeply eroded elements of a long-lived compound stratovolcano are as old as 500 Ka, representing several eruptive episodes and an unbroken spectrum of rock compositions from 52 to 69% SiO<sub>2</sub>. Peripheral to the stratovolcano complex, coalescing shields of basalt (49-53% SiO<sub>2</sub>) underlie and interleave with lavas from the andesite-dacite focus of the overall magmatic system.

Persistent eruption of peripheral mafic shields and cinder cones throughout the lifetime of the stratovolcano suggests their fundamental role in transport of heat and magma into the crust. The term "parasitic" should be abandoned for such vents, as it promotes the view that mafic flank eruptions are leaks from a central chamber and implies dependence on the main cone. For large andesite-dacite systems, nearly the opposite is true: The stratocones are the more derivative features, built up of magmas that evolve by concurrent fractionation, crustal melting, assimilation, and recurrent internal mixing within the foci of the larger domains injected by the various mantle-derived basalts that also yield variable arrays of peripheral eruptions.

Dacitic lavas and a few block-and-ash flows (63-69% SiO<sub>2</sub>) erupted several times from the focal area, but probably none are younger than 100 Ka. A single rhyolite (72% SiO<sub>2</sub>), as yet undated, is also old. The antiquity of all known silicic units, in conjunction with the andesitic compositions of the late Pleistocene summit cone and the Holocene lavas that erupted high around that cone, make it appear unlikely that Mount Adams is now underlain by a magma reservoir large enough to support either a major hydrothermal convection system or fractionation of voluminous silicic differentiates.

The high precipitation on the stratocone, estimated to be ~3500 mm/a, makes Mount Adams an important site of ground water recharge. The high permeability of its rubbly lava flows suggests that much of this ground water



moves rapidly downward and outward from the stratocone, failing to remain in the warmer central region of the volcano long enough to develop a hydrothermal convection pattern but, instead, dispersing and dissipating whatever heat is supplied from depth to the fumarolically altered core. The weak and diffuse fumarolic emissions on the summit are the only manifestations of possible hydrothermal activity anywhere on the stratocone. The warmest spring that we have found on the stratovolcano measured 3°C, on a summer day. Nonetheless, in surrounding areas, springs as warm as 24°C and gradients of 50°C/km in heat-flow holes offer some hope that ground water sufficiently warm for local space heating or agricultural uses might be found by exploring the lowland periphery of the volcano.

Cascades Workshop, Menlo Park, May 22-23, 1985

Characteristics of Cascades magmatic systems determined  
from seismic studies

H. M. Iyer

Introduction

Seismic techniques (refraction, reflection, seismicity mapping, and velocity and attenuation measurements using seismic waves from local earthquakes, regional earthquakes, and teleseisms, have been used to delineate magma chambers in several silicic volcanic centers in western U.S.A. (Iyer, 1984). However so far, such techniques have only been marginally successful in detecting magma chambers in the Cascade volcanoes. Seismic refraction and seismicity data have provided valuable information on the volcano-tectonic setting of the Cascades (California: Berge and Stauber, 1985; Zucca et al., 1985; Oregon: Leaver et al., 1982; Washington: Rohay, 1979; Taber and Smith, 1985). The most exciting result from teleseismic residuals is the detection of the subducting Juan de Fuca plate beneath Washington and Oregon (Michaelson and Weaver, 1985; Michaelson, personal communication). Teleseismic-residual data at the andesitic stratovolcanoes of the Cascades do not reveal detectable magma chambers. Examples are Mt. Hood, Mt. Shasta, and Mt. Lassen. On the other hand the Newberry and Medicine Lake volcanoes show evidence for massive high-velocity intrusions into the crust.

In this talk, I discuss P-wave residual modelling of the subducting Juan de Fuca plate beneath the Cascades, and results from teleseismic-residual experiments in Cascade stratovolcanoes and shield volcanoes. I also present preliminary results from a high-resolution experiment at the Newberry volcano to detect small (diameter 1 km or more) magmatic intrusions.

### Subduction Zone

Unlike in other major subduction zones of the world, due to the lack of a Benioff zone of deep earthquakes, it has been difficult to delineate the subducting lithosphere beneath the continental margin in western north America. At least in Washington and northern California, intermediate depth earthquakes seem to define a zone that looks like the shallow segment of the subducting plate. In Oregon, on the other hand, subduction seems to be taking place completely aseismically.

Michaelson and Weaver (1985) used teleseismic P-wave arrivals from earthquakes recorded by a dense network of seismic stations in Washington and northern Oregon, and using inverse and forward modelling techniques, delineated an eastward dipping high-velocity anomaly in the upper mantle, inferred to be the subducting plate. It is about 50 km thick, has P-velocity higher than the surrounding region by 3-7%, and extends from 45 to 300 km depth. Its dip angle is of the order of 60-70 Degrees. Michaelson and Weaver also found evidence for segmentation of the plate into three different configurations between northern Washington and northern Oregon.

Evidence for the slab beneath Oregon comes from the inversion of P-residual data from a USGS seismic network operated for two years in the Oregon Cascades and from a linear, northwesterly, profile of stations in

southern part of the state. Both data sets reveal the presence of an east-dipping, tabular, high-velocity anomaly in the upper mantle. The inversion and forward modelling of residual data from the profile (Caryl Michaelson, personal communication), show an eastward dipping 50-km thick slab in the depth range of 40-200 km. The dip angle is about 50 degrees and the velocity contrast in the slab from the surrounding rock is about 10% at the top and about 4% at depth.

#### Stratovolcano

We have looked at the three-dimensional crustal structure of three stratovolcanoes in the Cascades, Mt. Hood in Oregon (Weaver et al., 1982), and Mt. Shasta and Mt. Lassen in California, using teleseismic residuals collected by dense seismic networks. In all these cases the results show only regional heterogeneities in crustal structure with no clear evidence for the presence of magma chambers of horizontal or vertical dimensions greater than about 5 km, the resolution limit of the teleseismic technique.

#### Shield Volcano

Seismic and other geophysical data show the presence of high-velocity intrusions in the crust in the Newberry volcano, Oregon, and in the Medicine Lake volcano, California. Three-dimensional modeling of teleseismic P-wave residuals collected using a dense seismic array over Newberry volcano delineate a column of high-velocity material extending from within 10 km of the surface below the volcano summit to mid-crustal depths near 25 km (Stauber et al., 1985). The P-wave velocity in the column is about 6-14% higher than

in the surrounding crustal rocks. Stauber et al., (1985) interpret this intrusive body to be an expression of a swarm of predominantly sub-solidus gabbroic dikes and sills which were intruded as the volcano was built. A similar high-velocity body has been tentatively identified beneath the Medicine Lake volcano by Evans (1982).

#### High-resolution experiment

Neither the teleseismic results nor results from other geophysical experiments, however, can explain the high temperatures encountered in two shallow drill-holes drilled in the Newberry caldera. To investigate if magma intrusions undetectable due to the limited resolution of the teleseismic method are responsible for these high temperatures, we carried out a high-resolution version of the teleseismic-residual experiment by recording 8 large shots using a dense array of 120 seismographs deployed over a 12 km diameter area centered on the caldera. Preliminary interpretation of the traveltime residuals reveals a ring of high velocity material coinciding with the inner ring fault system of the caldera in the upper 2 km and a zone of lower velocity extending deeper than 2 km in the center of the caldera (Stauber et al., 1985). IS THE LOW-VELOCITY ZONE THE MAGMA CHAMBER WE HAVE BEEN SEARCHING FOR IN THE CASCADES?

#### REFERENCES

- Berge, P. A., and Stauber, D. A., 1985, Seismic refraction study of Lassen Peak, California (in review).

- Evans, J. R., 1982, Compressional-wave velocity structure of the Medicine Lake Volcano and vicinity from teleseismic relative traveltime residuals, in "Technical Programs, Abstracts, and Biographies", Society of Exploration Geophysicists Fifty-Second Annual International Meeting and Exposition", p. 482-485.
- Iyer, H. M., 1984, Geophysical evidence for the locations, shapes and sizes, and internal structures of magma chambers beneath regions of Quaternary volcanism, in "The Relative Contributions of Mantle, Oceanic Crust and Continental Crust to Magma Genesis", (Ed: S. Moorbath, R. N. Thompson, and E. R. Oxburgh), Phil. Trans. Royal Soc. London, Series A., v. 310, 473-510.
- Leaver, D. S., Mooney, W. D., and Kohler, W. M., 1984, A seismic refraction study of the Oregon Cascades, J. Geophys. Res., v. 89, 3121-3134.
- Michaelson, C. A., and Weaver, C. S., 1985, Upper mantle structure from teleseismic P-wave arrivals, J. Geophys. Res. (in press).
- Rohay, A., 1979, A refraction study of the north Cascades and northeastern Washington, in "Annual Technical Report on Earthquake Monitoring of the Hanford Region, Eastern Washington", (Ed: S. Malone), Geophysics Program, University of Washington, Seattle.
- Stauber, D. A., Green, S. M., and Iyer, H. M., 1985, Three-dimensional P-velocity structure of the crust below Newberry volcano, Oregon, J. Geophys. Res. (in press).
- Stauber, D. A., Iyer, H. M., Mooney, W. D., and Dawson, P. B., 1985, Three-dimensional P-velocity structure of the summit caldera of Newberry Volcano, Oregon, Geothermal Resources Council Trans. (in press).

- Taber, J. J., and Smith, S. W., 1985, Seismicity and focal mechanisms associated with the subduction of the Juan de Fuca plate beneath the Olympic Peninsula, Washington, Seismol. Soc. Am. Bull., v. 75, 237-249.
- Weaver, C. S., Green, S. M., and Iyer, H. M., 1982, Seismicity of Mt. Hood and structure as determined from teleseismic P-wave delay studies, J. Geophys. Res., v. 87, 2782-2792.
- Zucca, J. J., Fuis, G. S., Milkereit, B., and Mooney, W. D., and Catchings, R. D., 1985, Crustal structure of northeastern California from seismic refraction data, J. Geophys. Res. (in press).

# GRAVITY AND MAGNETIC STUDIES IN THE CASCADE RANGE

Carol Finn

U.S. Geological Survey

P.O. Box 25046, Denver, Colorado 80225

## ABSTRACT

A compatible set of gravity data for the entire Cascade Range has been compiled. From this data set I prepared a series of interpretive color gravity maps including a free air anomaly map, Bouguer anomaly map at a principle ( $2.67 \text{ g/cm}^3$ ) and an alternate ( $2.43 \text{ g/cm}^3$ ) reduction density, and filtered and derivative versions of the Bouguer anomaly map. The set is accompanied by a color terrain map at the same scale.

The regional anomaly pattern and gradients outline the various geological provinces adjacent to the Cascade Range and delineate major structural elements in the range. The more local anomalies and gradients may delineate low density basin and caldera fill, faults, and shallow plutons.

## Introduction

The U.S. Geological Survey (USGS) has supported gravity studies as part of its Cascades Geothermal Program. Gravity data are particularly useful in areas where there are large lateral density contrasts such as in volcanic regions. These density contrasts cause variations in the earth's gravity field which can often be related to subsurface geologic features such as faults, intrusions, ore bodies, etc.



The purpose of this paper is to provide preliminary interpretations of some of the local gravity and magnetic features in the Cascades.

The Pacific Northwest, the location of the Cascade Range, has a complex tectonic history involving changes in plate tectonic setting from a passive Atlantic-type margin to an active subduction zone (Dickinson, 1976; Hamilton, 1978; and Hammond, 1979), changes in the location of the subduction zone and its associated magmatic arc, accretion of various terranes and micro-plate rotations. The Cascade Range is a volcanic arc extending from Lassen Peak, California to Mt. Garibaldi, British Columbia and is related to subduction of the Juan de Fuca and Gorda plates. The Cascades consist of a wide range of rock types and ages ranging in composition from basalt through rhyolite and in age from Miocene to recent (they do, however, overlie Mesozoic and Paleozoic rocks in Washington and Canada). The diversity of rock types results in numerous density contrasts that produce a wide variety of gravity anomaly patterns.

#### Local Gravity Anomalies

Local gravity anomalies in the Cascades are generally caused by sediment filled basins (gravity lows), an incorrect Bouguer reduction density (can cause either gravity highs or lows, but lows are more common when a Bouguer reduction density of  $2.67 \text{ g/cm}^3$  is used), intrusions (gravity highs) and caldera fill (gravity lows). Since the Cascades south of Mt. Rainier are mantled by a layer of low density volcanics, gravity anomalies in this region usually delineate rocks which differ from the density of these volcanic rocks. Although the proper Bouguer reduction density for much of the High Cascades is between  $2.67$  and  $2.43 \text{ g/cm}^3$ , most of the young, recently active volcanoes in the range have a bulk density of  $2.2 \text{ g/cm}^3$ . When this density is

used to reduce the gravity data for most individual volcanoes, gravity highs, not lows, are commonly observed (Finn and Williams, 1982; Williams and Finn, in press<sup>b</sup>). These positive anomalies are due to shallow, dense intrusions that probably range in composition from intermediate to mafic. Many other local gravity highs have been identified with old, eroded volcanic centers. Examples are the Goat Rock volcano in the Goat Rocks Wilderness, Washington and the Still Creek and Laurel Hill plutons just southwest of Mt. Hood (Williams and Finn, 1983).

#### Aeromagnetic Data

All of the stratocone volcanoes in the Cascades have had aeromagnetic surveys flown over them. Studies of the data from Mt. Shasta (Blakely and Christiansen, 1978), Mt. Hood (Flanigan and Williams, 1982), Medicine Lake (Finn, 1982), Newberry (Griscom and Roberts, 1984) Goat Rocks Volcano (Williams and Finn, in press<sup>b</sup>), and Mt. St. Helens have been made. In most cases, i.e. at Medicine Lake, Newberry, and Mt. Shasta, the magnetic anomaly has been interpreted to be solely due to the terrain. At Mt. Hood, an old reversely-magnetized volcano upon which the present mountain was built was delineated with the magnetic data. Preliminary results from a combined gravity and magnetic study of Mt. St. Helens indicate that a magnetic and gravity high, unrelated to the terrain is associated with Mt. St. Helens. They may be related to a pluton under the volcano. More interesting results, in terms of delineation of geothermal resources, come from a study of gravity and magnetic data from Goat Rocks Volcano (Williams and Finn, in press<sup>(b)</sup>). Two and one-half-dimensional modeling of the positive gravity and magnetic anomalies delineated a shallow pluton. In most places the gravity and magnetic anomalies defined the pluton. However, some differences exist.

There are places where a magnetic low occurs where there is the gravity high associated with the pluton. These lows correspond to mapped areas of alteration, indicating that the zones of alteration is not just surficial, but must be meters thick. Other areas of mapped alteration do not have magnetic lows associated them, showing that the alteration is surficial. I hope to apply a similar approach to other volcanoes.

### Geothermal Resources

The delineation of structural features like faults and local features such as intrusions is useful for geothermal exploration. Faults can be zones of increased permeability. Intrusions under active volcanoes are often asymmetric to the volcanic cone and could be reached at a shallower depth if drill holes are sited with the benefit of the gravity data.

## References

- Blakely, R. J., and Christiansen, R. L., 1978, The magnetization of Mount Shasta and implications for virtual geomagnetic poles determined from seamounts: *Journal of Geophysical Research*, v. 83, p. 5971-5978.
- Dickinson, W.R., 1976, Sedimentary basins developed during the evolution of Mesozoic and Cenozoic arc-trench systems in western North America: *Can. Jour. of Earth Sci.*, v. 13, pp. 1268-1287.
- Finn, Carol, 1983, A magnetic study of Medicine Lake Volcano, California: *EOS*, v. 63, p. 1150.
- Finn, Carol, and Williams, D.L., 1982, Gravity evidence for a shallow intrusion under Medicine Lake Volcano, California: *Geology*, v. 10, pp. 503-507.
- Griscom, A., and Roberts, C. W., 1984, Gravity and magnetic interpretation of Newberry Volcano, Oregon: U.S. Geological Survey Open-File Report 84-\_\_\_.
- Hamilton, Warren, 1978, Mesozoic tectonics of the western United States: in D. G. Howell and K.A. Mc Dougall, eds., *Mesozoic paleogeography of the western United States: Pacific section*, Society of Econ. Paleontologists and Mineralogists, *Pac. Coast Paleogeography Symp. 2*, pp. 33-70.
- Hammond, P.E., 1979, A tectonic model for evolution of the Cascade Range, in J.M. Armentrout, M.R. Cole and H. TerBest Jr., eds., *Cenozoic paleogeography of the western United States: Pac. sect.*, Soc. of Econ. Paleontologists and Mineralogists, *Pacific Coast Paleogeography Symp. 3*, pp. 219-237.
- Williams, D.L., and Finn, Carol, 1985, Analysis of gravity data in volcanic terrain and gravity anomalies and subvolcanic intrusions in the Cascade Range and at other selected volcanoes: *in press* (A)

Williams, D. L., and Finn, Carol, in press,<sup>(b)</sup> Evidence for a mafic pluton  
beneath the Goat Rocks Wilderness, Washington, from gravity and magnetic  
data: J. Geophys. Res.

## GEOCHEMICAL FEATURES OF CASCADES HYDROTHERMAL SYSTEMS

By Robert H. Mariner  
U.S. Geological Survey, Menlo Park, CA

Thermal waters of the Cascade Range include at least 4 distinct types: Na-Cl, Na-SO<sub>4</sub>, CO<sub>2</sub>-charged, and dilute Na-HCO<sub>3</sub>. The most interesting, from a geothermal point of view, are the Na-Cl waters which occur in the Oregon Cascades and at Lassen Volcanic National Park. The large chloride content of these waters, up to 3,350 mg/L, must be derived from rock deposited in a marine environment, and in most of the Cascades this requires circulation to considerable depth. Most thermal springs in the central Oregon Cascades discharge Na-Cl waters which have a Ca-Cl component. In "Ca-Cl component" waters, such as Belknap Springs, some dissolved calcium must be used to electrically balance the chloride in the chemical analysis because insufficient sodium is present.

### Belknap Springs


	<u>t°C</u>	<u>pH</u>	<u>SiO<sub>2</sub></u>	<u>Ca</u>	<u>Mg</u>	<u>Na</u>	<u>K</u>	<u>HCO<sub>3</sub></u>	<u>SO<sub>4</sub></u>	<u>Cl</u>
mg/L	86	7.64	91	210	.34	660	15	20	150	1,200
meq/L				10.5		28.7				33.8

Although Ca-Cl thermal waters do not occur in the Cascades, several Ca-Cl mineral springs occur in western Oregon and Washington. Pigeon Springs, a cold mineral spring in western Washington, is an example of a Ca-Cl water:


### Pigeon Springs

	<u>t°C</u>	<u>pH</u>	<u>SiO<sub>2</sub></u>	<u>Ca</u>	<u>Mg</u>	<u>Na</u>	<u>K</u>	<u>HCO<sub>3</sub></u>	<u>SO<sub>4</sub></u>	<u>Cl</u>
mg/L	11	8.23	9.2	7,450	5.4	6,400	7.2	250	250	22,500
meq/L				372		278				635

Ca-Cl waters develop from Na-Cl waters by extensive albitization of plagioclase (Hardie, 1983). In the albitization of plagioclase, sodium and silica are consumed, and calcium is released:



A secondary reaction, brought about by the increase in calcium concentration is precipitation of anhydrite:



Provided that little or no water-rock reaction has taken place after the thermal water left the aquifer at depth, and that mixing of thermal and cold groundwater are not important, we can estimate the aquifer-temperature at depth by increasing the water temperature until the activity product constant for the reaction of interest equals the equilibrium constant. At Belknap Springs, equilibrium with respect to the albite-anorthite pair would be attained at 152°C. A similar calculation with respect to anhydrite, also produces an equilibrium temperature of 152°C.

Testing the "Ca-Cl" component waters of the Oregon Cascades for equilibrium with the mineral pair albite-anorthite and anhydrite, produces temperatures similar to those estimated from the  $\text{SO}_4\text{-H}_2\text{O}$  isotopic geothermometer (table 1). Quartz geothermometry does not work in this environment, probably because quartz is not present in the deep aquifer. Dissolved silica concentrations are limited by the albitization reaction not the dissolution of quartz. With the exception of Breitenbush,  $t_{\text{Na-K-Ca}}$  indicates temperatures within 25°C of the measured spring temperatures.

Albitization of plagioclase should be a common process in volcanic terranes such as the Cascades. However, before trying these equations in other areas, the effects of dilution with a low TDS water should be discussed. As an example, Belknap Hot Spring water was "diluted" with 10%, 25% and 50% distilled water. The apparent temperatures of equilibrium of these waters with the mineral pair albite-anorthite and anhydrite were calculated.

Percent dilution of Belknap water:	<u>0%</u>	<u>10%</u>	<u>25%</u>	<u>50%</u>
$t_{\text{anhydrite}}$	152	158	166	185
$t_{\text{albite-anorthite}}$	152	148	140	125

Dilution increases the apparent anhydrite saturation temperature but decreases the apparent albite-anorthite equilibrium temperature.

At Lassen, Growler and Morgan hot springs have anhydrite saturation and albite-anorthite equilibrium temperatures which range from 197 to 226°C.  $\text{SO}_4\text{-H}_2\text{O}$  isotopic equilibrium temperatures indicate 225 or 210°C, depending on whether conductive or adiabatic cooling is assumed. Gas geothermometer calculations indicate possible equilibrium temperatures of 242°C (Muffler and others, 1982).

$\text{Na-SO}_4$  waters of the Modoc Plateau and Klamath Falls are also low in magnesium. In the Modoc Plateau,  $t_{\text{Anhy}}$  and  $t_{\text{Ab-An}}$  give temperatures of 150 to 200°C, generally slightly cooler than the 190 to 205°C indicated by  $t_{\text{SO}_4\text{-H}_2\text{O}}$ . At Klamath Falls,  $t_{\text{Anhy}}$  and  $t_{\text{Ab-An}}$  indicate 165 to 175°C, while  $t_{\text{SO}_4\text{-H}_2\text{O}}$  indicates 185 to 200°C. This may indicate that chemical equilibrium last occurred in an aquifer at about 170°C but that  $\text{SO}_4\text{-H}_2\text{O}$  isotopic compositions preserve evidence of a hotter aquifer at about 190°C.



Na-SO<sub>4</sub> waters (up to 88°C) occur at Mt. St. Helens (M. J. Thompson, unpub. data) and have  $t_{\text{quartz}}$ ,  $t_{\text{Na-K-Ca}}$ , and  $t_{\text{Ab-An}}$  values which range from 183<sub>Qtz</sub> to 169<sub>Ab-An</sub>. The anhydrite saturation temperatures are lower, about 130°C probably due to dissolution of a sulfate mineral after the water left the thermal aquifer but before it discharged at the spring site. Magnesium concentrations are about 15 mg/L indicating that low temperature water-rock reaction has taken place. Na-SO<sub>4</sub> waters were also encountered in the Pucci well on Mount Hood (Robison, unpub. data), where  $t_{\text{Anhy}}$  and  $t_{\text{Ab-An}}$  indicate 140 to 150°C.

The CO<sub>2</sub>-charged waters of the Cascades are generally of moderate to low discharge temperature. Most carry several hundred mg/L chloride and up to a hundred mg/L magnesium. Except for springs in the Meager Mountain area, none have high SO<sub>4</sub>-H<sub>2</sub>O isotopic equilibrium temperatures. The springs at Meager Mountain ( $t_{\text{SO}_4\text{-H}_2\text{O}}$  of 220 to 290°C) are also higher in dissolved silica than the other CO<sub>2</sub>-charged thermal waters of the Cascades. Virtually all CO<sub>2</sub>-charged waters of the Cascades have  $t_{\text{Anhy}}$  and  $t_{\text{Ab-An}}$  values in the 170 to 275°C range. However, the SO<sub>4</sub>-H<sub>2</sub>O isotopic equilibrium temperature range is much wider, 10-290°C. It appears that the albitization reaction and anhydrite saturation temperatures are meaningless in CO<sub>2</sub>-charged water. The cation geothermometer (Na-K-Ca ± magnesium correction) gives temperatures near the spring temperatures for most CO<sub>2</sub>-charged waters. Although high temperatures are required for CO<sub>2</sub> generation, there is no proof that it is being generated in the thermal aquifer. It could just as well be generated at greater depth and dissolved in the water as the gas rises towards the land surface. The depth at which the CO<sub>2</sub> is being generated could be considerably beyond economic drilling depth.

Discharge rates for individual spring systems in the Oregon Cascades were determined from the chloride content and discharge rates of the streams at U.S.G.S. gaging sites. Total discharges (L/S) for individual spring groups, in the Oregon Cascades, correlate with the maximum spring temperature at the respective sites (figure 1). This correlation indicates that conductive cooling is the primary factor which determines spring temperature. A major entry of chloride water not associated with a hot spring occurs at Oakridge on the North Fork of the Middle Fork of the Willamette River. Both a cool saline well (22°C) and a saline spring (18°C) occur in Oakridge (Brown and others, 1980).

Chloride data for streams on the east side of the Cascades in central Oregon indicate that two large discharge cold springs contribute most of the chloride. Discharge rate measurements for the day and site we sampled are not yet available, but using discharge measurements for previous years, the headwater springs of the Metolius River seem to carry about 25 L/S of 1,200 mg/L chloride equivalent waters (Breitenbush equivalent), while the headwater springs of the Spring River carry about 10 L/S of 1,200 mg/L chloride equivalent water.

Deuterium compositions of thermal waters of the Oregon Cascades match the composition of cold springs along the crest of the Cascade Range. If the Cascade crest is the source of recharge of the thermal water, this requires that the circulation time is very short or that, fortuitously, the isotopic distribution of precipitation is the same now as when recharge occurred. Development of "Ca-Cl component" waters is thought to be a very slow process, so circulation times of these "Ca-Cl component" thermal waters could be longer than for most thermal waters. Dissolved helium

concentrations provide evidence for circulation times of at least 25,000 years, and perhaps much longer.

Gases discharged from the "Ca-Cl component" waters are principally nitrogen (>92%). They are peculiar, for geothermal water at least, in that their  $N_2/Ar$  ratios correlate with estimated aquifer-temperatures ( $t_{SO_4-H_2O}$ ). He versus  $N_2$  plots show two distinct trend lines, one including Belknap, Bigelson, Breitenbush, Foley, and perhaps Austin hot springs, while the other includes McCredie and Wall Creek hot springs.

#### REFERENCES

- Brown, D. E., McLean, G. D., Woller, N. M. and Black, G. L., 1980, Preliminary geology and geothermal resource potential of the Willamette Pass Area, Oregon: Oregon Department of Geology and Mineral Industries, Open-File Report 0-80-3, 65 p.
- Hardie, L. A., 1983, Origin of  $CaCl_2$  brines by basalt-seawater interaction: Insights provided by some simple mass balance calculations: Contributions to Mineralogy and Petrology, v. 82, pp. 205-213.
- Muffler, L. J. P., Nehring, N. L., Truesdell, A. H., Janik, C. J., Clynne, M. A. and Thompson, J. M., 1982, The Lassen Geothermal Systems: Proceedings of the Pacific Geothermal Conference, Auckland, New Zealand, November 1982, pp. 349-356.
- Thompson, 1985, Chemistry of thermal and nonthermal springs in the vicinity of Lassen Volcanic National Park: Journal of Volcanology and Geothermal Research, in press.

Table 1.--Estimated Aquifer Temperatures (°C) - Central Oregon Cascades

Name	t <sup>°C</sup>	t <sub>SO<sub>4</sub>-H<sub>2</sub>O</sub>	t <sub>Anhy</sub>	t <sub>Ab-An</sub>	t <sub>Quartz</sub>	t <sub>Na-K-Ca</sub>
Austin Hot Springs	86	181	186	155	126	84
Breitenbush Hot Springs	84	176	174	190	166	148
Bigelow Hot Springs	59	157	155	147	120	83
Belknap Hot Springs	86	148	152	155	132	82
Foley Hot Springs	79	--	100	130	113	45
Rider Creek Hot Springs	46	136	135	123	99	49
Wall Creek Hot Springs	40	--	160	137	115	69
McCredie Hot Springs	74	118	130	146	120	78
Kitson Hot Springs	44	--	134	133	96	82

N



S




Figure 1.--Measured spring temperature as a function of calculated discharge rate.

TYPES OF HYDROTHERMAL CONVECTION SYSTEMS IN THE  
IN THE CASCADE RANGE OF CALIFORNIA AND OREGON

Michael L. Sorey  
U.S. Geological Survey, Menlo Park, California

Three general types of hydrothermal convection systems can be delineated within the Cascade Range of California and Oregon. For the purposes of this abstract, they are termed summit-crater systems, caldera systems, and lateral-flow systems. Hydrothermal systems at most major Quaternary volcanic centers within this part of the Cascade Range can be identified with one of these types on the bases of test drilling, geophysical surveys (including heat flow), and the distribution and chemical characteristics of waters from thermal springs and fumaroles.

Summit-crater systems occur at Mt. Shasta and Mt. Hood and include surficial discharge of sub-boiling point fumaroles and steam-heated hot springs. Calculations based on empirical gas geothermometers applied to fumarolic samples yield subsurface temperature estimates in excess of 200°C at both areas (C.J. Janik and R.H. Mariner, personal communication, 1985). Similar thermal manifestations have not been detected at other andesite composite cones within the High Cascades of Oregon, possibly because of an absence of hot rocks at shallow depths. Summit-crater systems may include zones where lateral flows of thermal water cool conductively and mix with nonthermal groundwater before discharging in warm springs on the volcano flanks, as for example at Swim Springs south of Mt. Hood.

Caldera systems occur at Newberry Volcano, Medicine Lake Volcano, and Crater Lake. At these locations, ring fractures associated with caldera subsidence may provide conduits for deep

fluid circulation. In addition, long-lived silicic volcanism and Holocene eruptions indicate the presence of relatively shallow magmatic heat sources at each area. Test drilling at Newberry has provided temperature measurements and fluid samples that suggest that permeable strata containing water at temperatures near 265°C exist within the volcanic section at depths of 1-2 km. At Crater Lake, heat flow and geochemical investigations indicate the presence of hydrothermal convection to depths of 1-2 km beneath the lake floor and thermal-spring discharge at the lake floor. No clear evidence is yet available that hydrothermal circulation extends outside the caldera structures at these three locations, although attempts to delineate such circulation by test drilling and geophysical surveys are currently planned or in progress, and areas of anomalous soil mercury have been detected along a northeast-trending fissure/fault system on the southeast flank of Newberry Volcano (Priest and others, 1983).

A lateral outflow system occurs at Lassen Park and in the KGRA to the south. This system includes interconnected zones of vapor-dominated conditions and steam discharge beneath the summit region of the Lassen volcanic center and zones of lateral outflow of hot water that discharges in hot springs at lower elevations. Two such outflow zones have been delineated from test drilling and spring chemistry; one is fault-controlled and the other may occur along the contact between rocks from the Lassen volcanic center and underlying rocks from an older andesitic cone-building period. Water in the outflow conduits originates in a central upflow zone in which boiling occurs at temperatures near 240°C

and above which a parasitic vapor-dominated zone exists at pressures near 34 bars (Sorey and Ingebritsen, 1984). Similar systems could occur within the Cascade Range in Oregon; necessary conditions include: magmatic heat sources at mid to upper crustal depths, conduits for downward fluid circulation to depths of several kilometers, and permeable strata that allow lateral flow of heated water over distances of 15-25 km. Austin, Breitenbush, Belknap-Foley Hot Springs, and other lower temperature springs along the High Cascades - Western Cascades boundary may represent the discharge points for lateral outflow from High Cascade volcanoes to the east.

Constraints on the type of hydrothermal system(s) associated with these thermal-spring areas in Oregon are provided at present by chemical data from spring waters and thermal measurements in drill holes. Isotopic data from Mariner (this volume) suggest that the systems are recharged on or near the flanks of the High Cascade volcanoes rather than at lower altitudes along the High Cascade - Western Cascade boundary. Chloride flux measurements in streams yield estimates of the total flow of thermal water at Austin Hot Springs (110 L/s), Breitenbush Hot Springs (12 L/s), and Belknap-Foley Hot Springs (33 L/s). Corresponding estimates of reservoir temperatures based on sulfate isotope calculations are near 180°C for Austin and Breitenbush and 150°C for Belknap-Foley. Values of conductive heat flow and temperature gradient beneath the western part of the High Cascades in areas unaffected by vertical groundwater flow are consistently near 100 mWm<sup>-2</sup> and 65°C/km (Black, 1983). Thus, reservoir temperatures indicated for these spring systems could occur at depths of about 3 km,



possibly in marine sediments below the Cascade volcanics. A large component of marine origin in the hot spring waters is also indicated by chloride contents between 400 and 3400 mg/L.

Listed below are estimated convective heat flows, assumed reservoir areas, and conductive heat inputs required to sustain long-lasting regional hydrothermal systems that may be associated with each hot-spring area. The assumed reservoir areas approximate the areas of surface drainage between the springs and the axis of the High Cascades, and as such are probably overestimates of actual reservoir areas at depth.

Spring area	Convective heat flow	Reservoir area	Conductive heat input
	MW	km <sup>2</sup>	mWm <sup>-2</sup>
Austin	83	370	220
Breitenbush	9	170	53
Belknap-Foley	21	345	61

If, as implied by the models of Blackwell and others (1982), conductive heating of deeply circulating water occurs over about half the distance between the hot springs and the High Cascade axis, calculated rates of conductive heat input would increase to levels comparable to the estimated regional heat flow for two of the three spring systems. Only for the Austin Hot Springs system are required heat inputs too large to be supplied by regional heat flow, suggesting that for this system a more potent (magmatic) heat source or deeper levels of fluid circulation may be involved. Heat sources for fluid circulation at depth beneath the axis of the High Cascades could be restricted to youthful volcanoes with intermediate to silicic composition that may be

underlain by plutonic bodies, like Mt. Hood, Mt. Jefferson, and the South Sister.

Temperature data that has recently become available from the 2.4 km-deep exploration well drilled in 1981 near Breitenbush Hot Springs provide some needed detail on the nature of the regional hydrothermal systems in the Oregon Cascades. Bottom-hole temperature measurements in this well are consistent with the regional gradient of  $65^{\circ}\text{C}/\text{km}$  and heat flow of  $100 \text{ mWm}^{-2}$ . A heat flow of  $380 \text{ mWm}^{-2}$  is indicated above a depth of 750 m; this appears to result from lateral flow of water at about  $140^{\circ}\text{C}$  within the Western Cascade volcanic rocks encountered below that depth. This lateral flow could be fed by leakage from the upflow conduit(s) that supply hot water to Breitenbush Hot Springs. The Breitenbush well is located 2 km southeast of the springs, whereas temperature measurements in holes as close as 1 km to some other Cascade hot springs show no anomalous heat flow (Black, 1983).

The probable existence of regional-scale hydrothermal circulation systems within the Oregon Cascades offers promise for successful development of geothermal resources. However, several critical factors must first be evaluated. Confirmation that permeable reservoirs at temperatures near  $180^{\circ}\text{C}$  are present at depths of several kilometers is obviously needed. Although fluid temperatures significantly greater than  $180^{\circ}\text{C}$  may exist in these systems, such conditions may only occur in close proximity to heat sources along the High Cascade axis. The permeability distributions within these flow systems must also be delineated.

Low values for the average or effective permeability of each flow system between recharge and discharge areas are indicated by the rates of natural throughflow listed above, which are 1 to 2 orders of magnitude lower than throughflow rates in high permeability volcanic environments such as the Taupo graben in New Zealand. Regions of higher permeability may exist within the Cascade systems if throughflow rates are much greater than detected so far or are limited by zones of low permeability in regions of recharge or discharge. Such a possibility can only be evaluated by deep drilling.

#### References

- Black, G.L., 1983, Heat flow in the Oregon Cascades: in *Geology and geothermal resources of the central Oregon Cascade Range*, ed. Priest, G.R. and Vogt, B.F., Special Paper 15, Oregon Department of Geology and Mineral Industries, p. 69-76.
- Blackwell, D.D., Bowen, R.G., Hull, D.A., Riccio, J., and Steel, J.L., 1982, Heat flow, arc volcanism, and subduction in northern Oregon: *Journal of Geophysical Research*, v. 97, no. B10, p. 8735-8754.
- Priest, G.R., Hadden, M.M., Woller, N.M. and Brand, C.B., 1983 Preliminary soil-mercury survey of Newberry volcanoes: in *Survey of potential geothermal exploration sites at Newberry volcano, Deshutes County, Oregon*, Oregon Department of Geology and Mineral Industries Open-File Report 0-83-3.
- Sorey, M.L., and Ingebritsen, S.E., 1984, Quantitative analysis of the hydrothermal system in Lassen Volcanic National Park and Lassen Known Geothermal Resource Area: U.S. Geological Survey Water Resources Investigations Report 84-4278, 80 p.

## Mount Hood Geophysical Investigations

*Norman E. Goldstein*

Lawrence Berkeley Laboratory  
University of California  
Berkeley, CA

A number of geological, geochemical, and geophysical investigations were conducted around Mount Hood, Oregon (Williams et al., 1982) as part of a geochemical resource assessment jointly sponsored by DOE, U.S. Geological Survey, U.S. Department of Agriculture, Forest Service, and the State of Oregon. This particular Pleistocene-Holocene stratovolcano of the High Cascade Range was selected for study for several reasons:

- (1) its proximity to a major city and to nearby potential users of geothermal energy for space heating,
- (2) reasonably good road access, and
- (3) evidence for a thermal source beneath the summit.

Gravity, aeromagnetic, seismic and electromagnetic surveys were carried out by several research groups. We at LBL conducted (a) shallow controlled-source electromagnetic soundings, primarily on the accessible south and west flanks of the cone (Goldstein et al., 1982) and (b) deeper magnetotelluric and magnetotelluric-telluric soundings at station clusters at accessible locations around the cone (Mozley, 1982; Mozley et al., 1985). Our objectives were to determine whether we could discern any electrical resistivity anomalies, and to relate these to other geophysical anomalies and to possible features of geothermal interest. Because we were unable to occupy MT or telluric stations closer than 3 km from the summit, we discounted the possibility of detecting an anomaly from the conduit or a relatively shallow zone beneath the summit. We were extremely interested, however, in learning whether there might be a deeper conductor that could be related to a melt zone.

### **Thermal Gradient Drilling**

Geothermal data from 25 boreholes in a 20-km radius around the summit were collected and analyzed by Steele et al. (1982). They found that the thermal conditions are a complicated function of position around the volcano, which is in part related to rock age and rock type. A regional heat flow value of  $80 \text{ mW/m}^2$  was determined, the heat flow increasing towards the apex. Although holes could not be drilled sufficiently close to the apex to detect a neck-type magma chamber, the measured data and thermal modeling have placed certain constraints on the depth and size of the heat source. Steele et al. (1982) conclude that there cannot be a large, shallow subvolcanic melt zone less than 3 km deep and with a radius greater than 2-3 km.

### **Aeromagnetics**

Flanagan and Williams (1982) found a large, normally polarized magnetic high associated with the mountain. Removing the effects of a uniformly magnetized cone (the topographic anomaly), they found evidence in the residual anomaly for smaller, older volcanic cones largely concealed by the later eruptions, one cone under the north flank the other under the south flank. In addition, magnetic lows west and southwest of the mountain could be related to less magnetic Pliocene quartz diorite intrusives.

### **Subregional Gravity**

Couch and Gemperle (1979) found that Mount Hood is superimposed on a gravity low that suggests a north-south oriented graben-like structure. The cone itself is a Bouguer gravity high which, like other discrete highs in the survey area, is believed to be the denser core of the volcanic vent.

### **Detailed Gravity**

A detailed gravity survey revealed a dense body, presumably an intrusive, with a diameter of 5 km beneath the summit (Williams and Keith, 1982). The body appears to neck down to a diameter of 1 km with depth.

### **Seismic**

Results of a teleseismic P study indicated no velocity anomaly that one might expect if there were a large, shallow melt zone beneath the volcano (Weaver et al., 1982). The absence of a P-wave delay anomaly at Mount Hood and beneath Cascade volcanoes, in general, (Iyer et al., 1982) has been attributed to the possible small size of these melts (< 5 km in diameter) and the very complex vertical and lateral variations in velocity that makes identification difficult.

On the basis of a refraction survey, Kohler et al. (1982) found high seismic velocities on the west side of the volcano, wrapping around to the south. The high velocities seem to correlate with the magnetic low and an electrically resistive zone believed due to a broad area of Pliocene intrusives, only partially exposed by erosion.

### **Electromagnetics**

Several discrete conductive features were discerned after one-, two-, and three-dimensional modeling was done on the CSEM and MT data. Although the modeling did not produce a totally satisfying electrical model, the discrete anomalies could be assembled into a general composite model that seems to be consistent with other geophysical results.

### **Shallow Conductors**

Shallow conductors at depths of about 500 m were mapped on the east and south flanks. These conductors are believed due to the radial flow of meteoric water (mainly snowmelt) in permeable volcanic units. The waters are believed to be heated by hot rocks in the summit area (Wollenberg et al., 1979). The EM results agree with information from the hole at the base of the Pucci chairlift on the south flank.

### **Conduit Conductor**

Only weak evidence could be found for a neck-type conductor beneath the summit. Due to station locations, the presence or absence of a narrow conductor did not seem to affect the 2-D interpretation. However, some of the MT parameters in the 1 to 5 Hz bandwidth suggested a shallow conductor beneath the summit.

### **Resistive Anomalies**

Large areas of resistive rocks, extending from near the surface to 10+ km, were indicated on the south-southeast flank and beneath much of the western region of the volcano. There is a relatively strong correlation between the resistor, high velocities from a time-term analysis of refraction seismic data (Kohler et al., 1982), the aeromagnetics (Flanagan and Williams, 1982), and a few quartz diorite outcrops. The resistive nature of the rocks on the west flank and the occurrence of older intrusives was also confirmed by results from the Old Maid Flat well OMF-7A (Blackwell et al., 1982).

### **Deep Conductors**

An elongate conductor (1 ohm-m) striking N20°W was discerned at a depth of 12 km and with a depth extent of 10 km. The conductor is coincident with the trend of the High Cascades and it may be a zone of partial melt. Although this explanation has not been confirmed seismically, it is a reasonable one based on the corrected conductive thermal gradient of ~65°C/km measured in a number of holes around the volcano (Steele et al., 1982). That this conductor was resolved at all by MT came as something of a pleasant surprise in view of the limited spatial and frequency window available. However, the dimensions of the conductor could not be resolved, and this uncertainty admits another resistivity model, one with an additional deeper conductor at ~50 km.

## REFERENCES

- Blackwell, D.D., Murphy, C.F., and Steele, J.L., 1982, Heat flow and geophysical log analysis for OMF-7A geothermal test well: *in* Priest, G.R., and Vogt, B.F., eds., *Geology and Geothermal Resources of the Mount Hood Area*, Dep. Geol. Miner. Ind., Oregon, *Spec. Pap. 14*, 47-56.
- Couch, R.W. and Gemperle, 1979, Gravity measurements in the area of Mount Hood, Oregon: *in* *Geothermal Resource Assessment of Mount Hood, Open File Report O-79-8*, 137-189, Ore. Dep. of Geol. and Miner. Ind., Portland.
- Flanagan, G.F. and Williams, D.L., 1982, A magnetic investigation of Mount Hood, Oregon: *J. Geophys. Res.*, 87 (B4), 2804-2814.
- Goldstein, N.E., Mozley, E., and Wilt, M., 1982, Interpretation of shallow electrical features from electromagnetic and magnetotelluric surveys at Mount Hood, Oregon: *J. Geophys. Res.*, 87 (B4), 2815-2828.
- Iyer, H.M., Rite, A., and Green, S.M., 1982, Search for geothermal heat sources in the Oregon Cascades by means of teleseismic P-residual technique: paper presented at the 52nd Annual International Meeting, Soc. Expl. Geophys., Dallas Texas.
- Kohler, W.M., Healy, J.H., and Wegener, S.S., 1982, Upper crustal structure of the Mount Hood, Oregon, region as revealed by time term analysis: *J. Geophys. Res.*, 87 (B1), 339-355.
- Mozley, E.C., 1982, An investigation of the conductivity distribution in the vicinity of a Cascade volcano: Ph.D. Thesis, Univ. of Calif., Berkeley, Lawrence Berkeley Laboratory Rept. LBL-15671, 386 pp.
- Mozley, E.C., Goldstein, N.E., and Morrison, H.F., 1985, Magnetotelluric investigations at Mount Hood, Oregon: Lawrence Berkeley Laboratory Rept. LBL-19547 (in preparation).
- Steele, J.L., Blackwell, D.D., and Robison, J.H., 1982, Heat flow in the vicinity of the Mount Hood Volcano, Oregon: *in* Priest, G.R., and Vogt, B.F., eds., *Geology and Geothermal Resources of the Mount Hood Area*, Oregon, State of Oregon, Dep. Geol. Miner. Ind., *Spec. Pap. 14*, 31-42.
- Weaver, C.S., Green, S.M., Iyer, H.M., 1982, Seismicity of Mount Hood and structure as determined from teleseismic P wave delay studies: *J. Geophys. Res.*, 87 (B4), 2782-2792.
- Williams, D.L. and Keith, T.E.C., 1982, Aeromagnetic and Bouguer gravity maps of the Mount Hood Wilderness, Clackamas and Hood River Counties, Oregon: U.S. Geol. Survey Misc. Field Investigations Map MF 1379-D, scale 1:62,500.
- Williams, D.L., Hull, D.A., Ackermann, H.D., and Beeson, M.H., 1982, The Mt. Hood region: Volcanic history, structure, and geothermal energy potential: *J. Geophys. Res.*, 87 (B4), 2767-2781.
- Wollenberg, H.A., Bowman, R.G., Bowman, H.R., and Strisower, B., 1979, Geochemical studies of rocks, water and gases at Mt. Hood, Oregon: Lawrence Berkeley Laboratory Rept. LBL-7092, 57 pp.



CHARACTERIZATION OF GEOTHERMAL SYSTEMS IN THE CASCADE RANGE  
FROM GEOELECTRICAL STUDIES

by Adel A. R. Zohdy

U.S. Geological Survey, Denver, CO 80225

Several types of geoelectrical surveys have been made by the U.S. Geological Survey in the Cascades volcanic terrains. A list of the methods used, the areas studied, and the names of the investigators follows:

- (a) Magnetotellurics (MT) regional studies over the entire Cascade range, by W. D. Stanley.
- (b) Audiomagnetotellurics (AMT), near Lassen National Park and Mt. Hood, by D. B. Hoover.
- (c) Telluric profiling, in Medicine Lake volcano and near Lassen National Park, by K. Christopherson and D. B. Hoover.
- (d) Schlumberger soundings, in Medicine Lake volcano by A. A. R. Zohdy and R. J. Bisdorf, and in Newberry by R. J. Bisdorf, and south of Lassen by R. J. Bisdorf and A. A. R. Zohdy.
- (e) Time domain electromagnetic soundings, in Medicine Lake volcano, by W. L. Anderson and F. C. Frischknecht, and in Newberry by D. V. Fitterman.
- (f) Frequency domain electromagnetic soundings, in Medicine Lake volcano by W. L. Anderson and F. C. Frischknecht.
- (g) Airborne electromagnetics (DIGHEM), in Medicine Lake volcano and near Lassen National Park, by D. Frazier and D. B. Hoover.
- (h) Self potential (SP), in Mt. Hood, by D. B. Hoover.

Two crater volcanoes, namely Medicine Lake and Newberry, were studied more extensively by most of these methods. The geoelectric sections in these two areas are similar and may be described from top to bottom by four geoelectric units: (a) A very high resistivity unit of 15,000 to 150,000 ohm-m and of few tens of meters in thickness. This unit is comprised of loose pumice and dry volcanic rocks. (b) A medium-high resistivity unit of 500 to 5,000 ohm-m with thickness of 500 m to 1 km. This unit probably corresponds to Quaternary volcanic rocks that may be saturated in part with fresh, cold water. (c) A low resistivity unit of 5 to 30 ohm meters with a thickness of 500 meters to greater than 1 km. This unit probably represents altered volcanic rocks possibly saturated with circulating hot water. It represents the shallow geothermal resource target. (d) A high resistivity geoelectric basement of  $> 500$  ohm-m which is several kilometers in thickness. The shallowest depths to its top are about 1.5 to 2 km. It is believed to represent Tertiary intrusive rocks.

The exploration of these areas by the direct current resistivity method (Schlumberger soundings) proved to be a challenging task because of very high contact resistances, near absence of long straight roads, and off the road dense forest and fallen trees. Special formulas were derived for computing the proper geometric factors, digitizing the road maps, and correcting the field measurements. The mathematical solution to the winding road problem has opened the door to investigating many other areas with little concern to the presence of straight roads of sufficient length.

In the Medicine Lake area, a DIGHEM airborne electromagnetic survey was made to study the areal distribution of electrical resistivity at shallow depths of the order of few meters or tens of meters. The results showed three low resistivity anomalies (10 to 100 ohm-m), possibly representing near

surface hydrothermal alterations. The anomalies are near the northwest corner of Medicine Lake itself, southeast of the Hot Spot, and near Telephone Flat (slightly north of Bullseye Lake). The Schlumberger, time domain, frequency domain, and MT soundings showed that an extensive low resistivity layer of about 5 to 30 ohm-meters underlies a high resistivity cover, and that significant low resistivity anomalies are centered near Bullseye Lake and northwest of Medicine Lake. The results of the 51 Schlumberger soundings are illustrated by two cross sections and by four maps of interpreted-true resistivity at depths of 250 m, 500 m, 750 m, and 1,000 m.

At Newberry, an east-west geoelectric section constructed from the interpretation of Schlumberger soundings shows a low resistivity zone that corresponds to the increase in thermal gradient observed in a USGS test well which encountered 256°C water at a depth of 930 m. Another low resistivity zone about 600 m deep is present just to the west of the caldera boundary. A map of interpreted-true resistivity at a depth of 750 m shows that this western low resistivity zone has an areal extent of about 16 km<sup>2</sup>. This zone is interesting enough that at least one company has drilled it. Several time domain electromagnetic soundings also were made in the Newberry volcano area.

Regional east-west magnetotelluric cross sections show that the axis of the Quaternary volcanoes of the High Cascades, to the west of Medicine Lake and Newberry, coincides with an apparent structural trough. The sections also show that, beneath Medicine Lake and Newberry, there are high resistivity rocks believed to consist of multiple small intrusions forming structural highs underneath the low resistivity zones. These structural highs (which were detected by only very few of the Schlumberger and time domain soundings) at a depth of about 2 km or less could be considered as fundamental geothermal targets if they consist of multiple young intrusive units.

DEPARTMENT OF ENERGY DRILLING IN THE CASCADE RANGE

by Marshall Reed  
Geothermal and Hydropower Technologies Division  
U.S. Department of Energy  
Washington, D.C.

This year the U.S. Department of Energy, Geothermal and Hydropower Technologies Division (GHTD) will share with the geothermal exploration industry the cost of drilling several thermal-gradient holes to depths of 1 km or more in the Cascade Range. In return for its cost share, GHTD will receive copies of the drilling and geophysical logs, samples of cuttings and cores, and a period of access to the holes for further geophysical measurements. Participation in this prospect drilling will allow GHTD to evaluate the effectiveness of intermediate-depth thermal-gradient holes in locating geothermal resources in areas like the Cascades, where percolating surface water masks the thermal gradient at shallow depths. A total funding of \$1 million is available this year for the GHTD share of the drilling.

Several factors were considered in the planning for this project. Shallow holes in the older volcanic rocks of the Western Cascades provided good measurements of the conductive thermal gradient, but in the High Cascades thermal measurements in shallow holes were dominated by cold, moving ground water. Drilling at Mount Hood and Newberry Caldera indicated that thermal-gradient holes should be at least 1 km deep to penetrate the cold ground water and to reach conductive gradients. Much deeper holes would be required to penetrate the geothermal systems thought to exist at these two sites, but the drilling of intermediate-depth holes seemed to offer an attractive research tool. Since thermal-gradient holes can be drilled at a small diameter, they

are much less expensive than production wells. Sharing costs with industry is an attractive way of doubling the available experimental data from thermal-gradient sites.

Because of cost sharing, data from this project will be limited to geothermal prospects in the region of the Cascades and Newberry and Medicine Lake Calderas. Industry drilling is confined to those areas where geothermal development is permitted, and it is anticipated that the drilling will be on land with existing geothermal leases. An important aspect of this project from the industry viewpoint is that a reasonable chance of development exists if a resource is indicated.

In addition to sharing the drilling costs, GHTD will support research in and around these gradient holes. Stable temperature measurements will be made during the period of access, and thermal conductivities will be measured in cores from each hole to provide calculations of heat flow. This project requires the drilling of holes to 1 km or deeper, but the method of drilling is up to the bidder. If core holes are drilled, we will have a complete lithologic record that can be compared to surface surveys and provide constraints on geologic models. If larger-diameter holes are drilled, it may be possible to run several geophysical logs to aid in the interpretation of surface surveys. It is possible that thermal water will be intercepted in one or more holes, and if hole stability and environmental safety can be maintained there are plans to obtain water samples for geochemical analysis. We would appreciate your suggestions as to other research or exploration techniques that would enhance the data from this drilling project.

## POSSIBLE SITES FOR SCIENTIFIC DRILLING IN THE CASCADE RANGE

George R. Priest (Oregon Department of Geology and Mineral Industries)

### INTRODUCTION

A great deal has been learned about the Cascade Range from surface surveys and shallow temperature-gradient drilling. Nevertheless, a number of problems have stubbornly resisted solution. Among the most important of these are:

1. Mapping and characterization of hydrothermal systems in both the Western Cascades and the High Cascades.
2. Determination of background conductive heat flow in the High Cascade Range.
3. Mapping of the pre-Quaternary structure and stratigraphy of the High Cascades.
4. Accurate modeling of the source of the regional heat flow anomaly, including energy and mass balance calculations relative to rates of volcanism and subduction.

Drilling of intermediate-depth and deep drill holes is crucial to solution of these and related problems. Only drill holes can provide the necessary fluid and rock samples and down-hole geophysical measurements to place quantitative constraints on possible geologic and hydrologic models.

### WHERE SHOULD DRILLING BE FOCUSED?

Given that drilling should occur in the Cascades, the problem then is to decide which areas have the greatest potential for resolving the above problems. Most of the problems are related to the thermal regime of the Cascades, so picking sites can, to a first approximation, be accomplished utilizing a geothermal exploration strategy. The geothermal exploration industry has for decades utilized a phased approach to drilling programs. Surface geological, geochemical, and geophysical surveys are normally followed by shallow drilling of temperature-gradient holes which in turn are followed by drilling of deeper wells to test for hydrothermal fluids. Tables 1 and 2 are qualitative summaries of where various parts of the Cascade Range stand with respect to resource potential, access, and completion of temperature-gradient drilling. These tables take into account only publicly available data. The evaluation categories in the tables could be separated into a number of subcategories corresponding to the various exploration phases, but this would not materially affect the final ranking of sites. The point of this exercise is that there are only a few areas which are at present suitable targets for scientific drilling programs.

TABLE 1. Qualitative summary of the quality of the data base, access, and geothermal resource potential at possible sites for research drilling in the Cascades (0 = Poor; 1 = Moderately Poor; 2 = Moderate; 3 = Moderately Good; 4 = Good; 5 = Very Good). Soil geochemical surveys in the public domain are only available for Newberry and Medicine Lake volcanoes. Data from wells drilled to depths in excess of 600 m are only available for the Mount Hood area, the Breitenbush area, and Newberry Volcano.

Site	Geologic Coverage	Drilling (Heat Flow)	Geophysics (Non-heat flow)	Access	Resource Potential
<u>WASHINGTON</u>					
Mt. Baker	3	0	3	3	4
Mt. Rainier	3	0	2	1	4
Glacier Peak	2	0	2	1	4
Mt. Adams	5	0	4	1	4
Mt. St. Helens	5	0	4	1	4
N. & Cent. Wash. between volcanoes	2	0	2	2	0
Southern Wash. between volcanoes	2	2	3	4	2
<u>OREGON</u>					
Mt. Hood	4	2	3	1	4
Adjacent to Mt. Hood	3	4	3	3	2
Mt. Jefferson area	4	0	3	0	4
Breitenbush area	5	5	3	5	4
Santiam Pass area	5	2	3	4	4
North Sister area	5	0	3	0	4
Middle Sister area	5	0	3	0	4
McKenzie Pass	5	1	3	1	4
South Sister	5	0	3	0	5
Century Drive area	5	0	3	2	5
Newberry Volcano	5	2	5	3	5
Willamette Pass area	4	2	3	4	3
Crater Lake	5	1	3	1	5
Adjacent to Crater Lake	3	0	3	4	4
Oregon passes S. of Crater Lake	3	0	3	4	3
<u>CALIFORNIA</u>					
Mt. Shasta	5	0	5	1	4
Adjacent to Mt. Shasta	5	3	5	4	3
Medicine Lake	5	3	5	5	5
Mt. Lassen	5	2	4	0	4
Between Calif. stratocones	3	1	3	4	3

Table 2. Ranking of sites based on total scores from Table 1. Parentheses indicate site has poor or moderately poor accessibility. Brackets indicate site has poor resource potential.

Score	Site
22	Breitenbush Hot Springs area, Medicine Lake Volcano
20	Newberry Volcano,
19	Areas adjacent to Mt. Shasta
18	Santiam Pass area
16	Willamette Pass area
15	Areas adjacent to Mt Hood, Century Drive area, (Crater Lake), (Mt. Shasta), (Mt. Lassen)
14	(Mt. Adams), (Mt. St. Helens), (Mt. Hood), (McKenzie Pass), areas adjacent to Crater Lake, areas between northern California stratocones south of Mt. Shasta
13	Mt. Baker, (South Sister), Oregon passes south of Crater Lake, southern Washington Cascades between major volcanoes
12	(North Sister area), (Middle Sister area)
11	(Mt. Jefferson area)
10	(Mt. Rainier)
9	(Glacier Peak)
6	{Northern and central Washington between major volcanoes}

Areas which stand out as good potential drilling sites are:

1. Newberry and Medicine Lake volcanoes
2. The Breitenbush-Santiam Pass transect, central Oregon
3. Areas adjacent to Mt. Shasta, northern California
4. Willamette Pass-Century Drive transect, central Oregon
5. Areas adjacent to Mt. Hood, northern Oregon

The Newberry and Medicine Lake volcanoes, major silicic volcanic centers with large data bases, scored high in the ranking. It is now well known that these volcanoes have a high potential for shallow, high-temperature hydrothermal systems, as shown by the USGS Newberry 2 well (Sammel, 1981). However, the writer is not convinced that further drilling in these areas will tell us much about the High Cascade Range to the west, particularly areas characterized by andesitic to basaltic volcanism. What is needed is an east-west profile of the heat flow in the High Cascade Range and one or more wells deep enough to sample potential hydrothermal systems. Ultimately, it would be of immense scientific and economic interest to penetrate to 7-10 km into the source region for the Cascade heat flow anomaly (Blackwell and others, 1982).

The Breitenbush area, considering that it is in the volcanically inactive Western Cascade Range, ranks very high relative to other sites (Table 2). The abundance of geological, geophysical, and drilling data and the presence of a known, potentially exploitable hydrothermal system caused the Breitenbush area to score high. Sunedco Energy and Development released much of their data for the Breitenbush area including data from their 2.5 km well (Well No. 58-28), the deepest well in the U.S. Cascades. This well intersected a major hydrothermal aquifer at temperatures in excess of 136° C (Waibel, 1985), and it can be reentered for deepening and additional scientific studies (Edward Western, personal communication). Data from numerous shallow temperature-gradient holes are also available for the Breitenbush area. The temperature-gradient data indicate that the aquifer in Well No. 58-28 dips to the east where it might be intercepted at higher temperature (Waibel, 1985).

Well No. 58-28 is also close to the western margin of a regional graben structure inferred from gravity studies of Couch and others (1982) and Couch and Foote (1983). Referred to as the "Cascade graben" (Couch and Foote, 1983), this structure apparently extends from Crater Lake to just north of Mount Jefferson (Couch and others, 1982). The structure is inferred to have experienced active displacements since the mid- to late-Miocene (Couch and Foote, 1983). Deepening the Sunedco Well would test the hypothesis that this structure has a down-to-the east displacement in excess of 3 km (Couch and Foote, 1983).

Deepening the Sunedco well to about 4 km will allow examination of the pre-Cascade crust (Couch, unpublished calculated gravity cross section). This will help to resolve whether this part of the Cascade arc is built on oceanic crust, as postulated by Hamilton and Meyers (1966) and others.



Drilling in the Breitenbush area will help to resolve many questions about the nature of the Western Cascade structure and hydrothermal convection but will not elucidate the structure and thermal regime of the High Cascades. The proximity of the Santiam Pass area to the Breitenbush site, its accessibility, and excellent geological coverage (Taylor, 1967; Taylor, unpublished 1:24,000-scale mapping) make the Santiam Pass area a prime target for a profile across the High Cascades. Santiam Pass is also one of the few easily accessible places in the High Cascade Range where the axis of volcanism can be located with a great deal of confidence. It thus affords an unique opportunity to test the hypothesis that the current axis of mafic to intermediate volcanism may have high conductive heat flow and a high potential for hydrothermal resources. In addition, the area lies in a large Pliocene graben which formed after eruption of voluminous late Miocene to early Pliocene ash flows and lavas from the High Cascades (Taylor, 1980). Drilling could provide quantitative constraints on the amount of displacement on this graben, and the nature of the pre-graben volcanic arc (now buried by Quaternary volcanoes). Drilling would also help to test the hypothesis that the regional residual gravity high under the Three Sisters-Mt. Jefferson segment is caused by a large intrusive complex (R. Couch, 1985, personal communication).

A comprehensive study of the Breitenbush area and the adjacent Santiam Pass area is the most logical and cost-effective first step in a scientific drilling program. This would examine both of the major graben structures, the mid- to late-Miocene Cascade Graben and the Pliocene graben in the High Cascades; it also would take advantage of a wealth of previous studies, including the opportunity to reenter the deepest well in the U.S. Cascades. In future years other east-west transects should be studied to help extrapolate the Santiam-Breitenbush data to the north and south.

Drilling in the areas adjacent to Mount Shasta would be of great value both in terms of geothermal assessment and volcanologic research. Drilling would capitalize on the detailed geophysical and geological data base which has been developed for this area by the USGS. Completing a transect to Medicine Lake would also be of interest to study the transition zone between the two areas.

The Century Drive area adjacent to the South Sister is unique in that it has the only Holocene rhyodacite volcanic centers in the High Cascade Range which are also accessible by a major highway. The area is part of a long-lived silicic highland of regional extent (Taylor, 1980). It would be an ideal place to establish whether hydrothermal resources associated with silicic volcanism are present in the main High Cascade Range. Heat flow and geophysical profiles across the full width of the Willamette Pass-Century Drive area would be of interest, especially in comparison with profiles at Breitenbush-Santiam Pass where mafic and intermediate volcanism prevails. Extending the studies to Newberry Volcano would allow investigation of the relationship between the silicic highland at Century drive and silicic volcanism at Newberry.

Areas adjacent to Mount Hood scored well in the ranking because of the large available data base. However, extensive drilling in the area has not been successful in locating indications of major high-temperature hydrothermal systems. According to some interpretations of the geophysical surveys (Goldstein, personal communication), the drill holes may be too shallow and not in the best locations to intercept the hydrothermal systems. In any case, the gravity data indicate that, unlike many of the large High Cascade volcanoes to the south, Mount Hood may not lie above a large, shallow intrusive complex (R. Couch, 1985, personal communication), although Williams and Finn (1983) concluded that a shallow high density intrusion could be present. South of Mount Hood the regional background heat flow increases to values in excess of  $100 \text{ mW/m}^2$  (Blackwell and others, 1982), which would be additive to any local heat flow associated with shallow intrusions. This factor would increase the likelihood of large hydrothermal systems in the Cascades south of Mount Hood even if shallow plutons are locally present in both areas.

A transect of detailed temperature-gradient drilling and surface surveys across the southern Washington Cascades would be valuable. This would help to improve the meager heat flow data base in Washington and allow comparisons between this area of somewhat lower regional heat flow (Blackwell and Steele, 1983) and lower rates of Quaternary volcanism to areas with higher rates in Oregon and northern California. This is important for developing a comprehensive model for the Cascades as a whole.

## CONCLUSIONS

The Santiam Pass-Breitenbush area is the best site for the first phase of a scientific drilling program in the Cascade Range. This area has a known hydrothermal convection system, a high rate of volcanism, high background heat flow, and a large existing data base for siting drill holes. It is recommended that Sunedco Well No. 58-28 near Breitenbush Hot Springs be reentered, tested, and deepened to about 4 km. Drilling of an additional well to about 2 km depth east of the Sunedco site might cross the east-dipping hydrothermal aquifer at higher temperatures (Waibel, 1985, personal communication). An east-west transect of at least three, and preferably four, 1 km or deeper slim holes should be drilled across the Cascades at the latitude of Santiam pass to delineate the temperatures and heat flow below the zone of rapidly circulating cold ground water. A 2.7 km or deeper well should be drilled near the axis of Quaternary volcanism to test for potential hydrothermal fluids and to explore the nature of the pre-Quaternary stratigraphy. The drilling should be coupled with a program of detailed surface geophysical surveys and extensive radiometric dating of volcanic units to help site the wells and to allow maximum extrapolation of the drilling data. Stratigraphic thicknesses determined from mapping and drilling should be combined with the lateral extent and age of volcanic units to calculate rates of volcanism. This should then be compared to contemporaneous rates of subduction,

heat flow, hydrothermal circulation, and crustal deformation to examine causal relationships.

Drilling programs similar in scope to the Breitenbush-Santiam study should be accomplished in the southern Washington Cascades, the Century Drive-Willamette Pass area, and the Mount Shasta area in future years in order to develop a comprehensive hydrologic and geologic model of the Cascade Range. It would also be useful to link up east-west transects across the Cascades to similar transects across the Newberry and Medicine Lake volcanoes to examine the interrelationships between these silicic volcanic centers and the main Cascade arc. Ultimately, drilling to depths of 7-10 km will be necessary to investigate deep magmatic and metamorphic processes operative under the area of the regional heat flow anomaly associated with the Cascade volcanic arc.

#### REFERENCES CITED

- Blackwell, D.D., Bowen, R.G., Hull, D.A., Riccio, J., and Steele, J.L., 1982, Heat flow, arc volcanism, and subduction in northern Oregon: *Journal of Geophysical Research*, v. 87, no. B10, p. 8735-8754.
- \_\_\_\_\_, and Steele, J.L., 1983, A summary of heat flow studies in the Cascade Range: *Geothermal Resources Council, Transactions*, v. 7, p. 233-236.
- Couch, R.W., Pitts, G.S., Gemperle, M., Braman, D.E., and Veen, C.A., 1982, Gravity anomalies in the Cascade Range in Oregon: Structural and thermal implications: Oregon Department of Geology and Mineral Industries Open-File Report 0-82-9, 43 p.
- Couch, R.W., and Foote, R.W., 1983, Graben structures of the Cascade Range in Oregon: *Transactions of the American Geophysical Union, EOS*, v. 64, no. 45, p. 887.
- Hamilton, W., and Myers, W.B., 1966, Cenozoic tectonics of the Western United States: *Reviews of Geophysics*, v. 4, no. 4, p. 509-549.
- Sammel, E.A., 1981, Results of test drilling at Newberry Volcano, Oregon: *Geothermal Resources Council Bulletin*, v. 10, no. 11, p. 3-8.
- Taylor, E.M., 1967, Recent volcanism between Three Fingered Jack and North Sister, Oregon Cascade Range: Pullman, Wash., Washington State University doctoral dissertation, 84 p.
- \_\_\_\_\_, 1980, Volcanic and volcanoclastic rocks on the east flank of the central Cascade Range to the Deschutes River, Oregon, in Oles, K.F., Johnson, J.G., Niem, A.R., and Niem, W.A., eds., *Geologic field trips in western Oregon and*

southwestern Washington: Oregon Department of Geology and Mineral Industries Bulletin 101, p. 1-7.

Waibel, A.F., 1985, Summary of data from Sunedco well 58-28, Breitenbush area, Oregon: Oregon Department of Geology and Mineral Industries, Oregon Geology, in preparation.

Williams, D. L., and Finn, C., 1983, Gravity and aeromagnetic study of the Goat Rocks Wilderness, Washington: Transactions of the American Geophysical Union, EOS, v. 64, no. 45, p. 887.

# GEOHERMAL POTENTIAL OF THE GARIBALDI BELT

## Mount Meager and Mount Cayley

J.G. Souther

Geological Survey of Canada, Vancouver B.C.

The Garibaldi Volcanic Belt of southwestern British Columbia is a northern extension of the High Cascades. It comprises 18 Pliocene and younger eruptive centres that lie within a narrow northwest-trending zone that includes Glacier Peak and Mount Baker but diverges from the main Cascade trend. Within this zone, volcanoes of the Garibaldi Belt are aligned along three short, north-trending, en echelon segments. The most easterly segment includes Mount Garibaldi, the central segment includes Mount Cayley, and the western segment includes the Meager Mountain Complex.

Each segment of the Garibaldi Belt includes calc-alkaline and alkaline rocks. The calc-alkaline assemblage is dominant, and ranges in composition from basaltic andesite, through andesite and dacite to rhyodacite. The alkaline suite (alkali olivine basalt, mugearite) forms less than 5% of the lava volume and is confined to relatively small monogenetic centres.

Petrogenetic modelling has been done only for the volcanoes of the eastern segment. There, the alkaline basalts are believed to be fractionation products of an alkali picrite magma derived by partial melting of mantle pyrolite at ca 1400°C and 25 kb.

Major- and trace-element abundances indicate that the alkaline basalt is not parental to andesites of the calc-alkaline series. The latter are believed to be the product of multistage fractionation of water-undersaturated tholeiite magma generated by partial melting of hydrous mantle peridotite at 60-85 km depths. Ascent of this magma to the surface was interrupted at several different depths where it underwent crystal fractionation leading to the dacite and rhyodacite magmas.

Geothermal exploration in the Garibaldi Belt has focused on Meager Mountain and Mount Cayley - the only two volcanoes that are associated with thermal springs.

Meager Mountain, with a total volume of -----km<sup>2</sup>, is a composite of at least nine overlapping volcanic assemblages that display a regular age progression from 1.9 Ma in the south to 2400 years B.P. in the north. Andesite and associated explosion breccia constitute most of the older units which are overlain and intruded by younger dacite flows and domes. The volcanic complex rests on a surface of high relief eroded into metamorphic and plutonic rocks of the late Mesozoic to early Tertiary Coast Plutonic Complex and late Miocene (7.9 Ma) epizonal plutons of the Pemberton Belt.

Pebble Creek hot spring (59°C) issues from basement north of the volcanic complex, and Meager Creek hot spring (58°C) issues from thick alluvial fill south of the complex. Chemical geothermometry on these waters predicts a range of subsurface temperatures from 78 to 168°C. The absence of tritium indicates a residence time in excess of 25 years. Oxygen isotope

studies show that Pebble Creek water lies on the meteoric line whereas Meager Creek water has shifted toward the heavy isotope.

Resistivity anomalies of 10 to 100  $\Omega$ -m were defined along the north and south flank of the Meager Mountain Complex in crystalline basement rocks with normal background values of 1500 to 6000  $\Omega$ -m. Diamond drilling in the northern anomaly encountered extensive zones of argillic alteration but failed to identify an active hydrothermal system. Diamond drilling in the south anomaly encountered temperatures above 200°C and confirmed the presence of a near-surface outflow plume of hot saline water issuing from basement fractures near the south slope of the complex. Subsequent drilling of three, 3000 m pre-production holes recorded a maximum temperature of 270°C. Low porosity and permeability precluded any production from two of the holes but the third has sustained a flow of more than 20 tonnes of mixed steam and water with an inflow temperature of 194°C for periods of 4 to 6 months. Production is from the Meager Creek fault zone which defines the southern edge of the hydrothermal system and dips north beneath the volcanic complex.

Large-array D.C. resistivity and A.M.T surveys in the south reservoir indicate a strong east-west polarization to depths of 1 to 4 km that probably reflect saline fluid and alteration along fractures that mimic the Meager Creek Fault. Low-frequency M.T. soundings record a strong N-S anisotropy that extends to 30 km and runs parallel to the alignment of eruptive centres. This may be interpreted as evidence of crustal rifting or dyke swarms. An apparent MT resistivity of 300 to 600  $\Omega$ -m at

a depth of 25 km may define a magmatic heat source. The mechanism of heat transfer from such a source to the near-surface hydrothermal system is unresolved. Geothermometry of spring and borehole waters suggests that circulation never penetrated significantly below the 200° isotherm. However, mixing models suggest that small amounts of brine have leaked from a deeper and much hotter hydrothermal system that is transferring heat from a magmatic source.

Mount Cayley, the largest volcano in the Central Garibaldi Belt, is a composite pile of dacite, rhyodacite and minor andesite that formed during at least three distinct stages of activity between 3 to 0.3 Ma. The main edifice, with a volume of about 15 km<sup>3</sup>, is a multiple plug dome underlain by a thick pyroclastic unit and overlain by remnants of a once extensive pyroclastic cone. The volcano rests on a steep, west-dipping surface underlain by crystalline rocks of the Coast Plutonic Complex. Deeply incised valleys were eroded through the central pile and into underlying basement rocks prior to eruption of two small exogenous domes on the south flank. Low resistivity anomalies appear to be associated with alteration and seeps of warm saline water in the vicinity of these domes. Diamond drilling has confirmed conductive gradients of about 100°C/km. The hydrothermal system appears to be controlled by north-trending regional fractures and porous breccia zones surrounding glass-rimmed dacite cupolas in basement rock beneath the central edifice.

Recoverable energy from Meager Mountain is estimated to be 9000 Mw-years and from Mount Cayley 4000 Mw-years.



**AGENDA**  
**WORKSHOP ON GEOTHERMAL RESOURCES OF THE CASCADE RANGE**

May 22

8:00-8:45 Patrick Muffler -- Undiscovered geothermal resources of the United States  
Marianne Guffanti -- Previous USGS estimates of geothermal resources of the Cascade Range

**SESSION I What is the regional geologic and tectonic setting in which geothermal systems of the Cascades occur?**

Objectives: To describe the Cascades as a regional feature and to provide the context in which to discuss specific magmatic/geothermal systems.

Moderator: Robert Christiansen

*Tape #1* 8:45- 9:05 ~~Craig Weaver~~ *Dal Stanley* -- Segmentation of the Juan de Fuca Plate  
9:05- 9:35 Related short, informal communications and discussion

*Tape #1* 9:35- 9:55 Norm MacLeod -- Volcanism in the Cascade Range  
9:55-10:25 Related short, informal communications and discussion

10:25-10:45 Break

*Tape #1* 10:45-11:05 David Blackwell -- Oregon heat flow  
11:05-11:35 Related short, informal communications and discussion

11:35-12:00 Open discussion period

12:00- 1:30 Lunch

**SESSION II What are the characteristics of magmatic systems as heat sources in the Cascades?**

Objectives: To summarize what is known about the volume, depth, age, composition, evolution, and eruptive history of Cascades magmatic systems and relate these characteristics to their geothermal potential.

Moderator: Wes Hildreth

1:30- 3:00 Characteristics of Cascades magmatic systems determined from geologic studies

*tape #2* { Robert Christiansen: Mt. Shasta  
Michael Clyne: Lassen volcanic center  
Julie Donnelly-Nolan: Medicine Lake volcano  
J. Donnelly-Nolan (for C. Bacon): Mt. Mazama & Crater Lake  
Norman MacLeod: Newberry volcano  
Ed Taylor: 3 Sisters to Jefferson area  
Wes Hildreth: Mt. Adams  
Jack Souther: Meager Mountain and Mt. Cayley

3:00- 3:30 Discussion and summary of geologic studies

3:30- 4:00 Break

*tape 2  
side 2* 4:00- 4:20 H. M. Iyer -- Characteristics of Cascades magmatic systems  
determined from seismic studies

4:20- 4:45 Related short, informal communications and discussion

4:45- 5:05 Carol Finn -- Gravity and magnetic studies in the Cascade  
Range

5:05- 5:30 Related short, informal communications and discussion

5:30 Adjourn

May 23

**SESSION III What is the hydrothermal expression of magmatism in the Cascades?**

Objectives: To discuss the nature of hydrothermal systems in the Cascades (i.e., their distribution, geometry, temperature, evolution, hydrodynamics) and their interaction with underlying magmatic systems, in order to provide a basis for assessing hydrothermal resources of the region.

Moderator: Alfred Truesdell

*tape 3  
side 1* 8:00- 8:20 Robert Mariner -- Geochemical features of Cascades  
hydrothermal systems

8:20- 8:50 Related, short informal communications and discussion

*side 1* 8:50- 9:10 Michael Sorey -- Types of hydrothermal convection systems  
in the Cascade Range of California and Oregon

9:10- 9:40 Related short, informal communications and discussion

9:40-10:10 Break

10:10-10:30 Norman Goldstein -- Mt. Hood geophysical investigations

10:30-11:00 Related short, informal communications and discussion

*tape 3  
side 2* 11:00-11:20 Adel Zohdy -- Characterization of geothermal reservoirs in  
the Cascade Range from geoelectric studies

11:20-11:50 Related short, informal communications and discussion

11:50 1:30 Lunch

(continued)

**SESSION IV What conceptual models of geothermal resources should guide drilling programs in the Cascades?**

Objectives: To outline, in general terms, geologic and hydrologic models of Cascades geothermal systems, as a basis for making recommendations about strategy for geothermal drilling in the Cascades.

Moderator: Clay Nichols

- Topic # 4* 1:30- 2:30 Panel on conceptual models of Cascades geothermal resources  
Patrick Muffler  
Robert Christiansen  
Norman MacLeod  
Alfred Truesdell  
Donald White  
Michael Sorey  
Craig Weaver
- 2:30- 3:00 Break
- 3:00- 3:20 Marshall Reed -- DOE drilling in the Cascade Range
- 3:20- 3:40 George Priest -- Possible sites for scientific drilling in the Cascade Range
- 3:40- 4:45 Discussion on drilling
- 4:45- 5:00 Patrick Muffler -- Closing remarks
- 5:00 Adjourn

**UNITED STATES GEOLOGICAL SURVEY  
GEOTHERMAL RESEARCH PROGRAM  
Fiscal Year 1985**

The goal of the Geothermal Research Program of the U.S. Geological Survey (USGS) is to improve our understanding of the nature, distribution, and energy potential of the Nation's geothermal resources, as a basis for determining the proper role of geothermal energy as one of the alternative energy sources in national energy policy. This goal is accomplished through multidisciplinary research designed to (1) determine the geological and hydrological factors that control the characteristics, occurrence, longevity, and size of all types of geothermal systems, (2) estimate the location and magnitude of the Nation's geothermal resources, periodically updating these assessments as new information and assessment methodologies are refined, and (3) evaluate geoenvironmental effects that may result from extraction and injection of geothermal fluids during development.

**ORGANIZATION**

The Geothermal Research Program comprises geologic, geochemical, geophysical, and hydrologic studies conducted within the Geologic and Water Resources Divisions. The program is administered by the Geologic Division under the direction of the Chief of the Office of Earthquakes, Volcanoes, and Engineering (John Filson, Chief, Reston, Virginia). The overall program is managed by a scientific coordinator (Patrick Muffler, Menlo Park, California) who is also directly responsible for coordinating geothermal investigations carried out within the Geologic Division. Studies conducted by the Water Resources Division are coordinated by Franklin H. Olmsted (Menlo Park). Donald E. White (Menlo Park) is the program's senior scientist and advisor. Donald W. Klick (Reston, Virginia) carries out program management duties associated with budgetary, administrative, and interagency liaison matters.

The geothermal research activities of the USGS are closely coordinated with those of the U.S. Department of Energy (DOE). Some activities of the Survey's Geothermal Research Program directly support certain objectives of the geothermal energy program of DOE. Accordingly, DOE has provided the USGS with supplemental funds occasionally to increase the timeliness of such activities. Since 1975, the Survey has provided a scientist (currently Raymond H. Wallace, Jr., Washington, D.C.) to DOE's Geothermal and Hydropower Technologies Division on a rotating basis to serve as liaison and advisor.

Since reorganization within the Department of the Interior (DOI) in 1982, the USGS no longer has the responsibility for evaluation and administration of Federal lands for geothermal leasing that had resided in the Conservation Division. That division was abolished, and its functions were transferred to the new DOI Minerals Management Service in January 1982, then to DOI's Bureau of Land Management in December 1982.

## BUDGET

Until 1971 the Survey did not have a separately organized and funded program of geothermal research and resource assessment. Limited investigations of hot springs, geysers, and hydrothermal systems had been conducted before 1971 as part of the Geologic Division's continuing study of the Nation's energy and mineral resources. In recognition of a need to examine alternate sources of energy, Congress authorized a specific program of geothermal research in fiscal year (FY) 1972. The fiscal history of the program is summarized below:

FY-71	\$205,000 (part of ongoing Geologic Division program)
FY-72	\$665,000
FY-73	\$2,255,000
FY-74	\$2,555,000 (also \$300,000 from NSF and \$120,000 from AEC)
FY-75	\$8,966,000 (also \$343,000 from ERDA)
FY-76	\$9,114,000 (also \$320,000 from Conservation Division, USGS, and \$315,590 from ERDA)
FY-77	\$9,243,000 (also \$130,705 from Conservation Division, USGS, and \$532,000 from ERDA)
FY-78	\$9,589,000 (also \$116,731 from Conservation Division, USGS, and \$1,011,832 from DOE)
FY-79	\$11,863,000 (also \$120,080 from Conservation Division, USGS, and \$1,499,000 from DOE)
FY-80	\$10,047,000 (also \$33,000 from Conservation Division, USGS, and \$1,474,831 from DOE)
FY-81	\$7,718,000 (also \$1,255,415 from DOE and \$135,000 from Conservation Division, USGS)
FY-82	\$6,771,000 (also \$575,000 from DOE and \$70,000 from the National Park Service)
FY-83	\$6,953,000 (also \$344,964 from DOE and \$98,000 from the National Park Service)
FY-84	\$7,284,000 (also \$240,000 from DOE, \$25,000 from the Agency for International Development, and \$52,169 from the National Park Service.)
FY-85	\$7,190,000 (funds from other Federal agencies not yet determined)

Additional information on the program's budget and objectives is available in Duffield and Guffanti (1981).

## ACCOMPLISHMENTS

Major accomplishments of the Geothermal Research Program over the past 12 years include (a) three major nationwide assessments of geothermal resources of the United States, (b) comprehensive multidisciplinary studies of major geothermal regions, including The Geysers, Long Valley, and Coso in California, Yellowstone National Park, and the Snake River Plain in Idaho, (c) development of methods to interpret water and gas analyses to give subsurface temperatures and to elucidate subsurface processes, (d) development of electrical and electromagnetic techniques for determining the geometry of geothermal systems, (e) development and refinement of various passive and active seismic techniques for investigation of hydrothermal systems and

underlying intrusive complexes, and (f) development of hydrologic models to describe mathematically the behavior of geothermal systems.

Specific recent accomplishments of the Geothermal Research Program include (1) publication of a nationwide assessment of geothermal resources at temperatures less than 90°C, (2) completion of a series of six 1:1,000,000 maps that show the distribution, rock type, and age of young volcanic centers in the United States, (3) completion of a 932-m drillhole that discovered a 265°C geothermal system at Newberry Volcano, Oregon, (4) development of a comprehensive geochemical and hydrologic model to describe the parasitic vapor-dominated geothermal system at Lassen Volcanic National Park, (5) development of geochemical and isotopic methods to analyze the behavior of geothermal reservoirs under exploitation, with specific application to Cerro Prieto in Mexico, Larderello in Italy, and The Geysers in California, (6) execution of a seismic refraction experiment in the Imperial Valley of southern California that describes crustal structures and rock types associated with the many geothermal reservoirs throughout the area, (7) drilling of a 331-m research borehole in lake sediments at Tulelake, California, for core recovery and study of detailed volcanic history of the southern Cascade Range, (8) execution of a comprehensive set of geological and geophysical studies at Newberry and Medicine Lake Volcanoes of the Cascade Range, (9) undertaking of regional geophysical surveys, including seismic refraction, magnetotelluric, aeromagnetic, and gravity studies, of the Cascade Range, and (10) delineation of the complex geometry of the Juan de Fuca plate beneath the Cascade Range, based on analysis of seismic data.

#### FY 1985 ACTIVITIES

In FY 1985, a wide variety of studies is continuing in the Cascade Range of northern California, Oregon, and Washington. The goal of this effort is to understand the active volcanic, tectonic, and hydrothermal processes of the Cascades as a framework for characterizing and quantifying geothermal energy resources. The Cascades studies include geologic mapping, geochronology, petrology of fresh and hydrothermally altered rock, fluid geochemistry, hydrology, and both regional and local geophysical surveys. Aeromagnetic and gravity coverage is complete for the entire Range. Results from ongoing, detailed geologic mapping is being compiled for a geologic map of the Cascades. The southern part of the Cascades, in particular the Newberry, Oregon, and Shasta/Medicine Lake, California, regions, is an area of comprehensive geological, geophysical, and hydrological studies

The current trend for the Geothermal Research Program is increasing emphasis on understanding the interaction of magmatic heat sources and geothermal systems and assessing all types of geothermal energy sources associated with young igneous systems. For example, the Questa caldera in northern New Mexico is being investigated to characterize magmatic/-hydrothermal processes in the roots of an older, well-exposed igneous system. Advanced seismic techniques are being refined to identify and characterize subsurface magma bodies. Geophysical studies on the island of Hawaii are delineating subsurface magmatic conduits of the active volcanoes Kilauea and Mauna Loa, and geologic mapping is being compiled into a new geologic map of the entire island. Geologic mapping of young silicic volcanic centers is underway in Alaska.

An important component of the FY 1985 program continues to be research to understand the dynamics and evolution of hydrothermal systems. Isotopic and chemical studies are carried out, particularly at the Cerro Prieto field in Mexico, Larderello in Italy, The Geysers in California, and Yellowstone National Park to refine geothermometers, elucidate hydrothermal processes, and determine the origin of chemical constituents in geothermal fluids. Geoelectrical techniques are developed and applied to determine the location and geometry of hydrothermal convection systems. Mineralogical studies are designed to characterize chemical interactions of geothermal fluids and the rocks through which they circulate, as exemplified by a study of hydrothermal alteration in core from a hole drilled in 1981 by the Geothermal Research Program at Newberry Volcano, Oregon.

The Geothermal Research Program also supports projects that evaluate selected geoenvironmental concerns of geothermal development. Monitoring and analysis of earthquakes continues at The Geysers geothermal field. Numerical modeling and seismic and geochemical monitoring are used to examine the extent to which thermal features in National Parks might be affected by possible geothermal development in adjacent areas.

Many other specific topics are investigated in projects listed below.

#### REFERENCES CITED

Duffield, W.A, and Guffanti, Marianne, 1981, The Geothermal Research Program of the Geological Survey: U.S. Geological Survey Open-File Report 81-504, 108p.

#### FY 1985 PROJECTS

Geologic Division -- \$6,052,000

Project No.	Title	Project Chief
9310-00839	Alaska Geothermal Studies	Miller, T. P.
9320-03305	Geologic Map of Cascade Range	Smith, J. G.
9380-00342	Resistivity Interpretation	Zohdy, A. A. R.
9380-01699	Geothermal Petrophysics	Olhoeft, G. R.
9380-01746	Electrical Techniques for Shallow to Medium Depth Exploration for Geothermal Systems	Hoover, D. B.
9380-01748	Hawaiian Geothermal Studies	Kauahikaua, J.
9380-01750	Electromagnetic Modeling & Inversion of Controlled-Source Measurements	Anderson, W. L.
9380-01752	Physical Properties of Crust/Upper Mantle	Fitterman, D. V.
9380-03057	Infrared Aerial Surveillance of Volcanoes of Cascade Range, Hawaii, & Alaska	Friedman, J. D.
9380-03071	Geophysical Characterization of Young Silicic Volcanic Fields	Williams, D. L.
9380-03085	Geoelectric Sounding Studies	Stanley, W. D.
9380-03398	Aeromagnetic Surveys	Blakely, R. J.
9380-03830	Gravity & Magnetic Modeling	Gettings, M. E.
9450-02739	Lake City Caldera Studies	Grauch, R. I.

9460-03408	Marine Metals	Clague, D. A.
9530-02171	Roots of Calderas	Lipman, P. W.
9540-01943	Sacramento Valley & No. Sierra	Harwood, D. S.
9540-01947	Tephrochronology	Sarna-Wojcicki, A.
9540-03234	Northwest California Framework	McLaughlin, R. J.
9540-03512	Sauk Sheet (Glacier Peak)	Tabor, R. W.
9540-03770	Cima Volcanic Field	Wilshire, H. G.
9570-00374	Radiocarbon	Rubin, M.
9570-01140	Geothermal Geochronology	Fleck, R. J.
9570-01376	Gaseous Emanation Detection	Reimer, G. M.
9570-01474	Stable Isotopes and Ore Genesis	Rye, R. O.
9570-01763	Obsidian Hydration Dating	Friedman, I.
9570-01764	Radiocarbon Geochronology	Robinson, S. W.
9590-03772	Paleoclimatic History of Tulelake, CA	Adam, D. P.
9920-03379	Socorro Magma Bodies	Jaksha, L. A.
9930-01160	N. Calif. Seismicity Studies	Lester, F. W.
9930-01170	Field Experiment Operations	VanSchaack, J. R.
9930-01172	Teleseismic & Microearthquake Studies in Geothermal Areas	Iyer, H. M.
9930-01496	Active Seismic Studies of Recent Volcanic Systems	Hill, D. P.
9930-01499	Data Processing Center Operations	VanSchaack, J. R.
9930-01891	Central Calif. Net Operations	Hall, W. D.
9930-02097	Geothermal Seismotectonic Studies	Weaver, C. S.
9930-02102	Active Seismic Studies of Fault Zones	Mooney, W. D.
9930-02106	Earthquake Studies in The Geysers/ Clear Lake Area, CA	Oppenheimer, D. H.
9930-03354	Seismological Data Processing	Bekins, B. A.
9930-03412	Consolidated Digital Recording	Stewart, S. W.
9930-03563	Hawaiian Seismic Studies	Klein, F.
9930- NEW	Seismic Analysis, N. Calif.	Cockerham, R. S.
9960-01176	Geothermal Studies (Heat Flow)	Sass, J. H.
9980-00283	Geologic Thermometry	Huebner, J. S.
9980-00292	Hydrothermal Fluids	Fournier, R. O.
9980-00299	Thermodynamic Properties of Minerals	Robie, R. A.
9980-00304	Geochemical Indicators	Truesdell, A. H.
9980-00314	Geology & Petrology, Long Valley-Mono Basin	Bailey, R. A.
9980-00322	Active & Fossil Geothermal Systems	White, D. E.
9980-01379	Mineralogical Crystal Chemistry	Evans, H. T.
9980-01772	Volcanology & Geothermal Potential of Lassen Region, CA	Muffler, P.
9980-01875	Geothermal Reservoirs	Nathenson, M.
9980-01922	Thermodynamic Data	Haas, J. L.
9980-02321	Petrology of Clear Lake Volcanics	Hearn, B. C.
9980-02322	Eruptive History of Mauna Loa Volcano, HI	Lockwood, J.
9980-02344	Hydrothermal Alteration in the Cascades	Bargar, K. E.
9980-02481	Volcanic Evolution of Crater Lake	Bacon, C. R.
9980-02500	Medicine Lake Volcano, CA	Donnelly-Nolan, J.
9980-02649	Multiphase Fluid Flow in Geothermal Systems	Kieffer, S. W.
9980-02797	Geology of Hualalai Volcano, Hawaii	Moore, R. B.
9980-02798	Systems Analysis of Geologic Rate Processes	Shaw, H. R.
9980-02900	Hawaiian Volcano Observatory	Wright, T. L.
9980-02906	Volcanology & Petrology of Late Cenozoic Magmatic Systems	Christiansen, R. L.



9980-03388	Mono-Inyo Craters, CA	Miller, C. D.
9980-03646	Mineralogy & Geochemistry of Hydrothermal Alteration	Keith, T. E. C.
9980-03648	Geologic Map of the Island of Hawaii	Wolfe, E. W.
9980-03719	Mineralogical Investigations	Erd, R. C.
9980-03862	Internal Structure of Magmatic Systems	Ryan, M. P.
9980- NEW	Geothermal Volcanology	Duffield, W. A.

Water Resources Division -- \$1,138,000

<u>Project No.</u>	<u>Title</u>	<u>Project Chief</u>
CR 80-257	Geothermal Subsidence Research	Riley, F.
CR 140-FG	Borehole Geophysics Applied to Geothermal Research	Paillet, F. L.
NR 75-064	Physical Chemistry of Stable Isotope Fractionation in Hydrologic Processes	Coplen, T.
WR 72-082	Hydrologic Reconnaissance of Geothermal Areas in Nevada & California	Olmsted, F. H.
WR 73-102	Modeling of Geothermal Systems	Sorey, M. L.
WR 74-121	Reconnaissance Studies and Numerical Modeling of Geothermal Areas in Oregon	Sammel, E. A.
WR 75-127G	Energy Transport in Groundwater	Moench, A. F.
WR 76-139	Geochemistry of Geopressured Systems	Kharaka, Y. K.
4387-16500	Western U.S. Geochemistry	Mariner, R. H.

**UNITED STATES GEOLOGICAL SURVEY  
VOLCANO HAZARDS PROGRAM  
Fiscal Year 1985**

**OBJECTIVES AND ORGANIZATION**

The Disaster Relief Act of 1974 (PL 93-288) assigns to the U.S. Geological Survey (USGS) the responsibility of providing timely warnings of volcanic eruptions and related activity. The ultimate goal of the USGS Volcano Hazards Program is, therefore, to reduce the loss of life, property, and natural resources that can result from volcanic eruptions and related consequences. Progress toward this goal is being achieved by a broad range of work with the objectives of (1) preparing assessments of the potential geologic and hydrologic hazards associated with eruptions of volcanoes in the United States, (2) monitoring behavior of active and potentially hazardous volcanoes, (3) conducting fundamental volcanological and related hydrological research, and (4) providing hazards information to other Federal, state, and local agencies and officials for land-use and emergency-response planning. The hazards assessments and the monitoring projects are carried out with primary emphasis on those volcanoes that appear to be most likely to erupt in the near future and that are major threats to life and economic well-being of those areas. A more detailed description of the objectives and long-range plans of the Volcano Hazards Program is available in Bailey and others (1983).

The current Volcano Hazards Program (VHP) is an interdivisional activity involving the cooperative efforts of the Geologic, Water Resources, and National Mapping Divisions. The Geologic Division is primarily responsible for seismic, geodetic, geochemical, and observational monitoring of volcanoes and for volcanic hazards assessments and related research. The Geologic Division component of the VHP is administered out of the Office of Earthquakes, Volcanoes, and Engineering (John Filson, Chief) and managed by a scientific coordinator (Patrick Muffler, Menlo Park, California). The Water Resources Division conducts related hydrologic monitoring and hazards assessments and research. The coordination for these activities is administered out of the Office of the Assistant Chief Hydrologist for Research and Technical Coordination (Gordon Bennett, Reston, Virginia). The National Mapping Division prepares a wide range of cartographic and geographic data and products essential for the above activities, as well as for emergency-response planning of other Federal, state, and local agencies.

In order to achieve program objectives and to maintain a competent and experienced staff of volcanologists available for emergencies, the USGS operates two volcano observatories: the Hawaiian Volcano Observatory (HVO) at Kilauea volcano, Hawaii, and the Cascades Volcano Observatory (CVO) at Vancouver, Washington, near Mount St. Helens. These two facilities serve not only as bases for monitoring the active and dormant volcanoes of Hawaii and the Cascades, but also are centers for the training of volcanologists and for research and development of new monitoring techniques and instruments. The Hawaiian Volcano Observatory has been in operation since 1912, and has pioneered the development of most of the monitoring methods and instruments now used worldwide. It has a staff of about 30 volcanological scientists and technicians. The Cascades Volcano Observatory was established in 1980 to monitor the current activity of Mount St. Helens, which is expected to

continue for as long as several decades, and to serve as a base for periodic monitoring of the other, currently less active but potentially dangerous, Cascade volcanoes. It presently has a staff of about 60, including volcanologists, geophysicists, geochemists, hydrologists, technicians, and other support personnel.

#### FUNDING HISTORY

The funding history of the USGS Volcano Hazards Program for fiscal years 1978 through 1985 is shown below. Funding in 1978 and 1979 reflects the older, modest program which covered only monitoring studies at the Hawaiian Volcano Observatory and very limited volcanic hazards assessments in the Cascades. No hydrologic studies were funded during this period. Funding in 1980 and 1981 reflects the increases resulting from the Mount St. Helens eruption and the establishment of a Cascades Volcano Observatory in Vancouver, Washington, late in 1980.

1978 -- \$ 1,055,000	1982 -- \$ 9,664,000
1979 -- \$ 1,078,000	1983 -- \$10,803,000
1980 -- \$ 4,387,000	1984 -- \$11,651,000
1981 -- \$12,656,000	1985 -- \$11,273,000

#### FY 1985 Activities

For many volcanoes, the nature of their activity remains similar over relatively long periods of time; thus, the study of the past activity of a given volcano provides a means of forecasting the nature of its future activity. Hazards assessments are based on detailed studies of the size, distribution, and age of various historic and prehistoric volcanic deposits in order to estimate the kinds, magnitudes, timing, and effects of possible future eruptions. Such studies were pioneered by USGS geologists beginning in the 1950's and have been demonstrated to be extremely useful worldwide.

Volcanic hazard studies in recent years have been concentrated on the Cascade Range volcanoes of the Pacific Northwest which tend to erupt relatively frequently and often violently. Volcanic hazards maps and assessment reports have been published (some preliminarily) for Mt. Baker, Glacier Peak, Mt. Rainier, and Mount St. Helens in Washington; Mt. Hood, Oregon; Mt. Shasta and the Long-Valley-Mono Lake area in California; and the islands of Maui and Hawaii. In addition, hydrologic hazards studies have been completed for Mt. Hood and Mt. Shasta, and glacier-ice volume studies have been completed for Mt. Rainier and Mt. Hood. Long-range plans include preparation of hazards assessments for all potentially hazardous U.S. volcanoes. During FY85, assessment efforts are focused on the Long Valley area and Lassen Volcanic National Park in California, Mt. Rainier, Mt. St. Helens, and Mt. Baker, Washington, Three Sisters volcanic center and Newberry volcano in Oregon, on the island of Hawaii, and at selected Alaskan volcanoes.

The basic objectives of volcano monitoring are (1) to detect and track as early as possible the rise of magma from depth into the shallow crust so that warnings can be issued before the magma erupts at the surface and (2) to

provide early warning of hazardous floods and mudflows that can be generated from an eruption. During FY84, monitoring activities were carried out in three principal locations: Long Valley-Mono Lake area in California, the Cascade Range of Washington, Oregon, and California, and the Island of Hawaii.

Monitoring studies in the Long Valley-Mono Lake volcanic area in California, were begun in 1980 after a renewal of seismic activity. Seismic, geodetic, and other geophysical monitoring has indicated the presence of a partly molten, shallow magma reservoir beneath the Long Valley caldera, and has revealed that nearly 18 inches of domical ground uplift occurred above it between 1975 and 1984. Earthquakes localized in an epicentral area near the town of Mammoth Lakes suggest that magma may have been injected to crustal depths of a few miles. On the basis of these discoveries and particularly in response to an earthquake swarm in January 1983, the USGS has intensified its hydrologic and geologic monitoring activities in the area.

In FY 1985, the Volcano Hazards Program continued monitoring of seismic networks at Mount St. Helens and other northern Cascades volcanoes by a joint University of Washington/USGS team; seismic networks are also maintained by the USGS at volcanoes in the southern Cascades. The USGS is primarily responsible for geodetic monitoring throughout the Cascades. At Mount St. Helens, the USGS has established a telemetered hydrologic monitoring system for Spirit Lake (now dammed by debris from the 1980 eruption) in order to transmit real-time water-level data to provide immediate warning of any failure of the debris blockage. A collection of articles was published summarizing the results of eruption prediction and geophysical, geochemical, and geological monitoring at Mount St. Helens through 1982 (Science, v. 221, September 1983).

Extensive seismic, geodetic, geochemical, and geologic monitoring is conducted by HVO at Kilauea and Mauna Loa volcanoes on the Island of Hawaii. Both volcanoes erupted in 1984, and in late March 1984 they were in concurrent eruption for the first time since 1919. The east-rift eruption of Kilauea began in January 1983 and has continued episodically into 1985, occasionally threatening communities on the southeast coast of Hawaii. Mauna Loa erupted for the first time since 1975 on March 25 to April 15, 1984. Lavas erupted from Mauna Loa's northeast rift zone traveled about 17 miles from the vent, approaching to within 5 miles of the populated outskirts of the town of Hilo. HVO carefully monitored the progress of these eruptions and provided short-term warnings to the public as needed.

Another area of concern to the USGS is Yellowstone National Park where recent geologic activity is similar in many respects to the current unrest at Long Valley. The Yellowstone region is one of extensive young volcanism and has experienced several strong earthquakes, numerous earthquake swarms, significant hydrothermal changes, and almost 35 inches of crustal uplift during the last century. In order to document continuing changes, the USGS increased its monitoring effort there in 1983.

Eruption research carried out within the Geologic Division has the general goals of explaining the distribution of volcanoes in space and time and of understanding eruptive processes. This research includes such studies as the correlation of volcanic activity with other possibly related geophysical phenomena, the physical properties and chemical evolution of magma, and the

mechanisms of magma generation at depth, transport through the crust, and eruption at the surface. Results of such studies are vital to our understanding of the mechanisms and causes of volcanism and provide the only sound basis for predicting eruptions and assessing volcanic hazards. Hydrologic research is directed toward understanding the impacts of volcanic eruptions on rivers and lakes to aid in the planning of effective mitigation. The principal efforts include the study of the geomorphology of the sediments of volcanic origin, the study of the process of transport of extremely high sediment loads, the study of the processes of mudflow movement and the physical properties of the flowing material, and the modeling of rainfall-runoff events in order to document the effects of ashfall and the return of the drainage to stable conditions. A significant research accomplishment of the Volcano Hazards Program is the publication by the USGS of an 844-page document (Lipman and Mullineaux, 1981) containing 64 reports that describe early results of studies of the volcanic activity and eruptive products at Mount St. Helens in 1980.

#### REFERENCES CITED

- Bailey, R. A., Beauchemin, P. R., Kapinos, F. P., and Klick, D. W., 1983, The Volcano Hazards Program -- objectives and long-range plans: U.S. Geological Survey Open-File Report 83-400, 33p.
- Lipman, P. W., and Mullineaux, D. R. (eds.), 1981, The 1980 eruptions of Mount St. Helens, Washington: U.S. Geological Survey Professional Paper 1250, 844p.

#### FY 1985 PROJECTS

The studies of the Survey's Volcano Hazards Program are carried out in individual projects. Of the total budget of \$11,273,000, the Geologic Division was allocated \$7,909,000, and the Water Resources Division was allocated \$3,264,000. National Mapping Division received \$100,000.

#### Geologic Division

Project No.	Title	Project Chief
9310-03049	Volcanic Hazards in Alaska	Miller, T. P.
9380-01748	Hawaiian Geothermal Studies	Kauahikaua, J.
9380-03057	Infrared Aerial Surveillance of Volcanics of the Cascade Range, Hawaii, and Alaska	Friedman, J. D.
9380-03069	Gravity Studies, Lassen National Park	Jachens, R. C.
9540-01947	Tephrochronology	Sarna-Wojcicki, A.
9570-00374	Carbon-14 Geochronology	Rubin, M.
9570-00377	Geochronology	Fleck, R. J.
9570-00382	Light Stable Isotopes	Friedman, I.
9570-01376	Gaseous Emanation Detection	Reimer, M. G.
9570-01568	Radiocarbon and U/Th Geochronology	Robinson, S.W.
9570-03066	Geomagnetic Secular Variation & Volcanic Timing	Champion, D. E.
9570- NEW	Photogrammetry (Mauna Loa)	Wu, S.
9590-03444	Quaternary Climate History, S-central AK	Ager, T. A.

9910-03589	Spirit Lake Hazards Evaluation- Liquefaction Studies	Chen, A. T. F.
9930-01160	Northern California Seismicity Studies	Lester, F. W.
9930-01170	Field Experiment Operations	VanSchaack, J. R.
9930-01496	Active Seismic Studies of Recent Volcanic Systems	Hill, D. P.
9930-01499	Data Processing Center Operations	VanSchaack, J. R.
9930-01891	Central California Network Operations	Hall, W. D.
9930-02097	Geothermal Seismotectonic Studies	Weaver, C. S.
9930-03353	Crustal Changes North of San Francisco Bay	Ward, P. L.
9930-03354	Seismic Data Processing	Bekins, B. A.
9930-03563	Hawaiian Seismic Studies	Klein, F.
9930- NEW	Seismic Analysis, N. Calif.	Cockerham, R. S.
9930- NEW	Sesimological Investigation of Magmatic Injection in Volcanic Terranes	Chouet, B.
9950-03024	Engineering Implications of the Mt. St. Helens Eruptions	Schuster, R. L.
9960-01182	Mechanics of Earthquake Faulting	Dieterich, J. H.
9960-01187	Crustal Strain	Prescott, W. H.
9960-01801	Experimental Tilt & Strain Instrumentation	Mortenson, C. E.
9960-02112	Mechanics of Geologic Structures Associated with Faulting	Segall, P.
9960-02114	Strain & Magnetic Field Operations	Johnston, M.
9960-02156	Geodetic Strain Monitoring	Langbein, J. O.
9960-03718	Long Valley Deformation	Savage, J. C.
9960-03815	Dilatometer Network Operations	Myren, G. D.
9980-00314	Geology & Petrology of the Long Valley- Mono Basin California	Bailey, R. A.
9980-00647	Volcanic Hazards	Hoblitt, R.
9980-02322	Eruptive History of Mauna Loa Volcano, HI	Lockwood, J. P.
9980-02431	Tephra Hazards from Cascade Range Volcanoes	Mullineaux, D. R.
9980-02649	Multiphase Fluid Flow in Geothermal Systems	Keiffer, S. W.
9980-02773	Geochemistry of Gas-Forming Elements	Sato, M.
9980-02789	System Analysis of Geologic Rate Processes	Shaw, H. R.
9980-02900	Hawaiian Volcano Observations	Wright, T. L.
9980-02992	Cascade Volcano Observations	Peterson, D. W.
9980-03385	Geology of Lassen Volcanic National Park	Clynne, M. A.
9980-03388	Volcanic Hazards of Mono-Inyo Craters, CA	Miller, C. D.
9980-03401	Hawaiian Eruptive Processes	Tilling, R. I.
9980-03656	Volcanic Processes	Decker, R. W.
9980-03664	Processes in Large Silicic Systems	Dzurisin, D.
9980-03721	Metals in Volcanic Exhalations	Hinckley, T.
9980-03860	Volcanic Crisis Assistance Team	Banks, N. G.
9980-03862	Internal Structure of Magmatic Systems	Ryan, M. P.
9980-03863	Short-term Volcanic Forecasting	Newhall, C. G.
9980-70029	Volcanic Hazards Support	Peterson, D. W.
9980-02914	Recent Volcanic Processes	Moore, J. G.

(continued)

Water Resources Division

Project No	Title	Project Chief
AK 84-158	Volcano-Glacier Interactions at Mt. Veniaminoff, AK	Emanuel, R. P.
CA 81-392	Hydrologic Hazards Assessment of Mt. Shasta, CA	Blodgett, J. C.
CA 82-431	Hydrologic Studies Related to Volcanic Activity in Long Valley, CA	Anttila, P. W.
CR 77-228	Precipitation-Runoff Modeling of Watershed Systems	Leavesley, G. H.
NR 79-099	Carbon Fluxes in Hydrologic & Geologic Processes	Sundquist, E. T.
OR 81-108	Hydrologic Hazards Assessment of Mt. Hood, OR	Hubbard, L. L.
OR 81-109	Hydrologic Hazards Assessment for Three Sisters Area, OR	Laenen, A.
SR 83-074	Fluvial Processes & River Mechanics	Chen, C.
WA 80-243	Hydrologic Data Collection in the Mt. St. Helens Area, WA	Childers, D.
WA 81-247	Sediment Deposition in the Cowlitz River, WA	Swift, C. H.
WA 81-250	New Lakes of North Fork Toutle River, Mt. St. Helens, WA	Carpenter, P. J.
WA 81-256	Sediment Transport - Mt. St. Helens, WA	Onions, C. A.
WA 82-272	Sedimentology of Lahars at Mt. St. Helens, WA	Scott, K. M.
WA 82-273	Mudflow/Debris Flow Rheology & Initiation	Pierson, T. C.
WA 82-274	Immediate & Long-Term Response to the 1980 Eruption of Mt. St. Helens, Toutle River Basin, WA	Janda, R. J.
WA 83-289	Sediment Transport Model Development & Preliminary Application to Columbia River	Sikonia, W. G.
WA 84-299	Hydrologic Hazards Assessment of Mt. Baker, WA	Dinicola, R. S.
WA 85-310	South Coldwater Geomorphology, WA	Meyer, D. F.
WA 85-311	Mss-Movement Dynamics -- Interaction with Groundwater and Streams	Iverson, R. M.
WD 80-157	Volcano Hazards Program Coordination	Kapinos, P.
WR 72-080	Volcanic Volatiles Barnes, I.	
WR 73-102	Modeling Geothermal Systems	Sorey, M. L.

23 May 85

## Cascades Cost-shared Drilling Meeting

Attendance: Marshall Reid, Clay Nichols, Marty Holloy, Norm Goldstein, Patrick Muffler, Bob Marinus, Mike Soreq, George Priest, Mike Wright, Dave Blackwell

Purpose: To obtain input re cascades drilling for possible inclusion in ID negotiations w/ proposers, especially on data gathering activities desired down-hole.

Preamble:

1. Proposals are in
2. Core holes to typical depths 4000 ft
3. Proposed completion -- iron pipe, sealed at bottom. Hole dia at bottom NQ
4. Limited period after drilling when rig will be on site for experiments
5. GP logs are proposed

Questions:

1. IS completion satisfactory?
2. How can water samples be obtained?
3. How can permeability data be obtained? and/or productivity?
4. How long after drilling should last temp log be obtained for equilibrium?
5. What are priorities for data collection down hole?
6. How long should/can we maintain access for science?



Completion

1 - better for A/F to cement core

every downlogs prob = before cementing

centering each of further core

2. Injection for permeability?

- hard to rock de materials; temp contrast will  
- disturb results.

3. ~~Injection~~ Drill stem tests? need 1 day - how deep dia?

4. Set up a steering communitas?

5. Drill w/ water? this is preferred


6. After samples -

down-hole sample area fine -  
pay for my standing in  
ahead counting paper in

7. BATT at each stop point could be expected  
to plan further logging / completion

8. drill ahead w/ fast returns when possible

9.



U.S. Geological Survey, Geothermal Research Program  
Workshop on the Geothermal Resources of the Cascade Range  
May 22-23, 1985 Menlo Park, CA

LIST OF PARTICIPANTS

Adam, David U.S. Geological Survey, MS 915  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 328-8111

✓ Bain, Desmond U.S. Forest Service  
630 Sansome Street  
San Francisco, CA 94111  
(415) 556-9768

Bargar, Keith U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓ Beall, Joe Geysers Geothermal Company  
8569 Sonoma Avenue  
Sebastopol, CA 95472

✓ Beilin, Phil U.S. Geological Survey  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Berge, Patricia U.S. Geological Survey, MS 977  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓ Blackwell, David Department of Geological Sciences  
Southern Methodist University  
Dallas, TX 75275  
(214) 692-2745

✓ Bodell, John Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

✓ Brophy, Paul California Energy Company  
3333 Mendocino Ave, #100  
Santa Rosa, CA 95401  
(707) 526-1000

✓ Casterson, Mark Geothermal Resources International  
545 Middlefield Road  
Menlo Park, CA 94025  
(415) 326-5470

✓ Chitwood, Larry

U.S. Forest Service  
Deschutes National Forest  
1645 Highway 20 E.  
Bend, OR 97701  
(503) 389-2373

✓ Christiansen, Robert

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓ Ciancanelli, Eugene

Cascadia Exploration Corporation  
3358 Apostol Road  
Escondido, CA 92025  
(619) 489-0969

✓ Collier, John

U.S. Forest Service  
Pacific Northwest Region  
P.O. Box 3623  
Portland, OR 97208  
(503) 221-2921

✓ Combs, James

Geothermal Resources International  
545 Middlefield Road, Suite 200  
Menlo Park, CA 94025  
(415) 326-5470

✓ Dansart, William

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

✓ DeBunce, Holly

Geology Department  
University of Oregon  
Eugene, OR 97403  
(503) 686-4573

✓ D'Olier, William

Thermal Power Company  
3333 Mendocino Avenue  
Santa Rosa, CA 95401  
(707) 576-7040

✓ DeWitt, David

Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

✓ Dondanville, Richard

Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

Erlich, Edward

U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1020

Fierstein, Judy U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Finn, Carol U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1345

Fitterman, David U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1382

Fournier, Robert U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Frischknecht, Frank U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1212

Fujimoto, Robert Bureau of Land Management  
Oregon State Office  
P.O. Box 2965  
Portland, OR 97208  
(503) 231-6946

Gaal, Robert Calif. State Lands Commission  
245 W. Broadway, Suite 425  
Long Beach, CA  
(213) 590-5233

Galanis, Peter U.S. Geological Survey, MS 23  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Geyer, John Bonneville Power Administration, *PRSB*  
P.O. Box 3621  
Portland, OR 97208  
(503) 230-5327

Goldstein, Norman Lawrence Berkeley Laboratory  
Earth Science Division, Bldg. 50-E  
Berkeley, CA 94720  
(415) 486-5961

Guffanti, Marianne U.S. Geological Survey  
922 National Center  
Reston, VA 22092  
(703) 860-7481

Hagood, Michael

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

Haizlip, Jill

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

Hammond, Paul

Department of Geology  
Portland State University  
P.O. Box 751  
Portland, OR 97207  
(503) 229-3022

Hansen, Peter

Geothermal Resources International  
545 Middlefield Road  
Menlo Park, CA 94025  
(415) 326-5470

Hebein, Jeffrey

Thermal Power Company  
3333 Mendocino Ave, Suite 120  
Santa Rosa, CA 95401  
(707) 576-1398

Higgins, Chris

California Division of Mines & Geology  
610 Bercut Drive  
Sacramento, CA 95814  
(916) 322-9997

Hildreth, Wes

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Holman, William

Department of Energy  
1333 Broadway  
Oakland, CA 94612  
(415) 273-6370

Hoover, Don

U.S. Geological Survey, MS 864  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1326

Ingebritsen, Steven

U.S. Geological Survey, MS 439  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-1111

Iovenitti, Joe

Thermal Power Company  
3333 Mendocino Ave., Suite 120  
Santa Rosa, CA 95401  
(707) 576-7232

Iyer, H. M.

U.S. Geological Survey, MS 977  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

James, Earl

Chevron Resources  
P.O. Box 7147  
San Francisco, CA 94120-7147  
(415) 894-2882

Johnson, Keith

Cascadia Exploration Corporation  
3358 Apostol Road  
Escondido, CA 92025  
(619) 489-0969

Johnson, Michael

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

Keith, Terry

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Klick, Donald

U.S. Geological Survey  
905 National Center  
Reston, VA 22092  
(703) 860-6471

LaFleur, Joe

California Energy Company  
3333 Mendocino Avenue, #100  
Santa Rosa CA 95401  
(707) 526-1000

Lippmann, Marcelo

Earth Sciences Division  
Lawrence Berkeley Laboratory  
Berkeley, CA 94720  
(415) 486-5035

MacLeod, Norman

U.S. Geological Survey  
Cascades Volcano Observatory  
5400 MacArthur Blvd.  
Vancouver, WA 98661  
(206) 696-7806

Mariner, Robert

U.S. Geological Survey, MS 434  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

McNitt, James

GeothermEx  
5221 Central Avenue, Suite 201  
Richmond, CA 94804  
(415) 527-9876

Michaelson, Caryl

U.S. Geological Survey, M.S. 977  
345 Middlefield Road  
Menlo Park, CA 94025  
(206) 545-3813

✓ Molloy, Martin

Department of Energy  
1333 Broadway  
Oakland, CA 94612  
(415) 273-7945

Mooney, Walter

U.S. Geological Survey, MS 977  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓ Moore, Joe

University of Utah Research Institute  
Earth Science Laboratory  
391 Chipeta Way, Suite C  
Salt Lake City, UT 84108  
(801) 524-3428

✓ Muffler, Patrick

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Nathenson, Manuel

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓ Nichols, Clay

Department of Energy  
550 Second Street  
Idaho Falls, Idaho 83401  
(208) 526-0535

✓ Nielson, Dennis

University of Utah Research Institute  
Earth Science Laboratory  
391 Chipeta Way, Suite C  
Salt Lake City, UT 84108  
(801) 524-3422

Nordquist, Gregg

Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

✓ Olmsted, Frank

U.S. Geological Survey, MS 467  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Olson, Harry J.

Steam Reserve Corporation  
1707 Cole Boulevard  
Golden, CO 80401  
(303) 231-0659



Pilkington, Dean

Steam Reserve Corporation  
1707 Cole Boulevard  
Golden, CO 80401  
(303) 231-0661

Priddy, Charles

California State Lands Commission  
1807 13th Street  
Sacramento, CA 95814  
(916) 323-7210

Priest, George

Department of Geology & Mineral Industries  
1005 State Office Building  
Portland, OR 97201  
(503) 229-5580

Prucher, Libby

Geology Department  
University of Oregon  
Eugene, OR 97403  
(503) 686-4573

Randall, Walter

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, California 95401  
(707) 523-4272

Reed, Marshall

Department of Energy  
CE-324  
Washington, D.C. 20585  
(202) 252-8017

John Rowley

Los Alamos National Laboratory, D-462  
P.O. Box 1663  
Los Alamos, NM 87545  
(505) 667-1378

Schriener, Alex

Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

Sherrod, David

U.S. Geological Survey, MS 901  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Sibbett, Bruce

University of Utah Research Institute  
Earth Science Laboratory  
391 Chipeta Way, Suite C  
Salt Lake City, UT 84108  
(801) 524-3424

Smith, James

U.S. Geological Survey, MS 901  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓Sorey, Michael U.S. Geological Survey, MS 439  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

✓Souther, J. G. Geological Survey of Canada  
100 West Pender St.  
Vancouver, B.C.  
Canada V6B 1R8  
(604) 666-0528

Stanley, Dal U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, Colorado 80225  
(303) 236-1328

Stephenson, Brian Union Geothermal Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

Sternfeld, Jeffry GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

✓Siemnich, Gene Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95406  
(707) 542-9543

Sun, Stanley Bureau of Land Management  
2800 Cottage Way  
Sacramento, CA 95825  
(916) 484-4515

✓Swanberg, Chandler Geothermal Resources International  
545 Middlefield Rd.  
Menlo Park, CA 94025  
(415) 321-5662

✓Swift, Charles Chevron Resources  
P.O. Box 7147  
San Francisco, 94120  
(415) 894-4037

Taylor, Edward Department of Geology  
Oregon State University  
Corvallis, OR 97331  
(503) 754-2484

Temko, Heidi Bureau of Land Management  
555 Leslie Street  
Ukiah, CA 95482  
(707) 462-3873

Thomas, Richard

California Division of Oil and Gas  
1916 9th St., Rm 1310  
Sacramento, CA 95826 95-814  
(916) 323-1788

Thompson, Michael

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Towne, E.B.

Geothermal Exploration  
582 Market Street, #1508  
San Francisco, Ca 94104  
(415) 434-0700

Truesdell, Alfred

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Waibel, Al

Columbia Geoscience  
22495 NW Quatama Road  
Hillsboro, OR 97124  
(503) 640-9877

Walker, George

U.S. Geological Survey, MS 901  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Walter, Stephen

U.S. Geological Survey, MS 977  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Walters, Mark

GEO Operator Corporation  
2300 County Center Drive  
Santa Rosa, CA 95401  
(707) 523-4272

White, Donald

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

White, Thomas

Bonneville Power Administration  
825 N.E. Multnomah  
Portland, OR 97208  
(503) 230-4386

Williamson, Ken

Union Oil Company  
P.O. Box 6854  
Santa Rosa, CA 95401  
(707) 542-9543

Winnett, Terrie

U.S. Geological Survey, MS 910  
345 Middlefield Road  
Menlo Park, CA 94025  
(415) 323-8111

Witham, Rodger

Buureau of Land Management  
2800 Cottage Way  
Sacramento, CA 95825  
(916) 484-4515

Wright, Mike

University of Utah Research Institute  
Earth Science Laboratory  
391 Chipeta Way, Suite C  
Salt Lake City, UT 84108  
(801) 524-3422

Zohdy, Adel

U.S. Geological Survey, MS 964  
Denver Federal Center  
Denver, CO 80225  
(303) 236-1222

Seismic Studies in the Cascade Range: Washington, Oregon, Northern California

- \*S Berg, J.W., Jr., and Baker, C.D., 1963, Oregon Earthquakes, 1941 through 1958: Bulletin of the Seismological Society of America, v. 53, p. 95-108.
- \*R Berg, J.W., Jr., Trembly, L., Emilia, D.A., Hutt, J.R., King, J.M., Long, L.T., McKnight, W.R., Sarmah, S.K., Souders, R., Thiruvathukal, J.V., and Vossler, D.A., 1966, Crustal refraction profile, Oregon Coast Range: Bulletin of the Seismological Society of America, v. 56, p. 1357-1362.
- \*R Berge, P.A., and Stauber, D.A., 1985, Seismic refraction study of the Lassen Peak area in California: (in review).
- \*T Berge, P.A., 1985, Teleseismic P-wave traveltime residuals across the Cascade Range in southern Oregon: U.S. Geological Survey Open-File report (in review).
- \*S Bolt, B.A., and Miller, R.D., 1971, Seismicity of northern and central California, 1965-1969: Bulletin of the Seismological Society of America, v. 61, p. 1831-1847.
- \*R Catchings, R.D., 1983, Crustal structure from seismic refraction in the Medicine Lake area of the Cascade Range and Modoc Plateau, northern California: Master of Science thesis, University of Wisconsin, Madison, Wisconsin, 97 p.
- S Chandra, U., 1974, Seismicity, earthquake mechanisms, and tectonics along the western coast of North America, from 42 degrees N to 61 degrees N: Bulletin of the Seismological Society of America, v. 64, p. 1529-1549.
- \*S Chiburis, E.F., 1966, Crustal structures in the Pacific Northwest states from phase-velocity dispersion of seismic surface waves: Phd. dissertation, Oregon State University, Corvallis, 170 p.
- \*S Crosson, R.S., 1976, Crustal structure modeling of earthquake data 2. Velocity structure of the Puget Sound region, Washington: Journal of Geophysical Research, v. 81, n. 17, p. 3047-3054.
- \*S+ Dehlinger, P., Chiburis, E.F., and Colver, M.M., 1965, Local travel-time curves and their geologic implications for the Pacific Northwest states: Bulletin of the Seismological Society of America, v. 55, p. 587-607.
- \*R Eaton, J.P., 1966, Crustal structure in northern and central California from seismic evidence, in Geology

of northern California: California Division of Mines and Geology Bulletin 190, p. 419-426.

- \*T Evans, J.R., 1982, Compressional-wave velocity structure of the Medicine Lake volcano and vicinity from teleseismic relative travelttime residuals: Technical Program of the Society of Exploration Geophysicists Fifty-Second Annual International Meeting and Exposition, Dallas, Texas, p. 482-485.
- \*R Hill, D.P., 1972, Crustal and upper mantle structure of the Columbia Plateau from long range seismic-refraction measurements: Geological Society of America Bulletin, v. 83, p. 1639-1648.
- RS Hill, D.P., 1978, Seismic evidence for the structure and Cenozoic tectonics of the Pacific Coast states, in Smith, R.B., and Eaton, G.P., eds., Cenozoic tectonics and regional geophysics of the Western Cordillera: Geological Society of America Memoir 152, p. 145-174.
- Hughes, J.M., Stoiber, R.E., and Carr, M.J., 1980, Segmentation of the Cascade volcanic chain: Geology, v. 8, p. 15-17.
- \*T Iyer, H.M., and Rite, A., 1981, Teleseismic P-wave delays in the Oregon Cascades: Transactions of the American Geophysical Union (EOS), abs., v. 62, p. 1089.
- \*R Johnson, S.H., and Couch, R.W., 1970, Crustal structure in the north Cascade Mountains of Washington and British Columbia from seismic-refraction measurements: Seismological Society of America Bulletin, v. 60, p. 1259-1270.
- \*S Klein, F.W., 1979, Earthquakes in Lassen Volcanic National Park, California: Bulletin of the Seismological Society of America, v. 69, p. 867-875.
- \*R Kohler, W.M., Healy, J.H., and Wegener, S.S., Upper crustal structure of the Mount Hood, Oregon, region as revealed by time term analysis: Journal of Geophysical Research, v. 87, p. 339-355.
- \*S Kollmann, A., 1984, Oregon Seismicity-- August 1980 to October 1982: U.S. Geological Survey Open-File Report 84-832, 32 p.
- \*S+ Langston, C.A., 1977, Corvallis, Oregon, crustal and upper mantle receiver structure from teleseismic P and S waves: Bulletin of the Seismological Society of America, v. 67, p. 713-724.
- \*S Langston, C.A., 1981, Evidence for the subducting lithosphere under Southern Vancouver Island and Western Oregon from

teleseismic P-wave conversion: Journal of Geophysical Research, v. 86, p. 3857-3866.

- \*RS Leaver, D.S., 1982, A refraction study of the Oregon Cascades: Master of Science Thesis, University of Washington.
- \*RS Leaver, D.S., Mooney, W.D., and Kohler, W.M., 1984, A seismic refraction study of the Oregon Cascades: Journal of Geophysical Research, v. 89, p. 3121-3134.
- McBirney, A.R., 1978, Volcanic evolution of the Cascade Range: Annual Review of Earth and Planetary Sciences, v. 6, p. 437-456.
- \*S McCollom, R.L., and Crosson, R.S., 1975, An array study of upper mantle velocity in Washington State: Seismological Society of America Bulletin, v. 65, p. 467-482.
- \*S McKenzie, D., and Julian, B., 1971, Puget Sound, Washington, earthquake and the mantle structure beneath the northwestern United States: Geological Society of America Bulletin, v. 82, p. 3519-3524.
- T Michaelson, C.A., 1983, Three-dimensional velocity structure of the crust and upper mantle in Washington and northern Oregon: M.S. thesis, University of Washington, 100 p.
- \*T Michaelson, C.A., and Weaver, C.S., 1985, Upper mantle structure from teleseismic P-wave arrivals: Journal of Geophysical Research (in review).
- \*T Monfort, M.E., 1980, P-wave residuals in the Mount Lassen volcanic area: Transactions of the American Geophysical Union (EOS), abs., v. 61, p. 1025.
- \*S Monfort, M.E., 1982, Seismic structure of the Lassen volcanic area: Transactions of the American Geophysical Union (EOS), abs., v. 63, p. 174.
- \*R Prodehl, D., 1970, Seismic refraction study of crustal structure in the Western United States: Geological Society of America Bulletin, v. 81, p. 2629-2646.
- \*S Rite, A., and Iyer, H.M., 1981, Oregon Seismicity: 1981 (abs.): American Geophysical Union Transactions (EOS), v. 62, p. 966.
- \*S Robinson, H.B., and Byerly, D., 1948, Earthquake swarm in Lassen Volcanic National Park: Bulletin of the Seismological Society of America, v. 38, p. 179-193.
- R Rohay, A., 1979, A refraction study of the North Cascades and northeastern Washington, in Malone, S., ed., Annual Technical Report on Earthquake Monitoring of the Hanford Region, Eastern Washington: Geophysics Program, University

of Washington, Seattle.

- R Shor, G.G., Jr., Dehlinger, P., Kirk, H.K., and French, W.S., 1968, Seismic refraction studies off Oregon and northern California: *Journal of Geophysical Research*, v. 73, p. 2175-2194.
- \*T Stauber, D.A., and Berge, P.A., 1984, P-velocity structure of Mt. Shasta, CA, and Newberry Volcano, abs.: *Transactions of the American Geophysical Union (EOS)*, v. 66, p. 25.
- \*T Stauber, D.A., Green, S.M., and Iyer, H.M., 1985, Three-dimensional P-velocity structure of the crust below Newberry Volcano, Oregon: submitted to the *Journal of Geophysical Research*.
- \*R Stauber, D.A., Iyer, H.M., Mooney, W.D., and Dawson, P.B., 1985, Three-dimensional P-velocity structure of the summit caldera of Newberry Volcano, Oregon: submitted to the *Geothermal Resources Council*.
- \*S Taber, J.J., and Smith, S.W., 1985, Seismicity and focal mechanisms associated with the subduction of the Juan de Fuca plate beneath the Olympic Peninsula, Washington: *Bulletin of the Seismological Society of America*, v. 75, p. 237-249.
- S Tobin, D.G., and Sykes, L.R., 1968, Seismicity and tectonics of the northeast Pacific Ocean: *Journal of Geophysical Research*, v. 73, p. 3821-3845.
- \*S Walter, S.R., Rojas, V., and Kollmann, A., 1984, Seismicity of the Lassen Peak area, California: *Geothermal Resources Council Transactions*, v. 8, p. 523-527.
- \*TS Weaver, C., Green, S.M., and Iyer, H.M., 1979, Seismic studies at the Mt. Hood volcano, northern Cascade Range, Oregon: *U.S. Geological Survey Open-File Report 1691*.
- \*TS Weaver, C., Green, S.M., and Iyer, H.M., 1982, Seismicity of Mount Hood and structure as determined from teleseismic P wave delay studies: *Journal of Geophysical Research*, v. 87, p. 2782-2792.
- S Weaver, C.S., and Smith, S.W., 1983, Regional tectonic and earthquake hazard implications of a crustal fault zone in southwestern Washington: *Journal of Geophysical Research*, v. 88, p. 10371-10383.
- S+ Weaver, C.S., and Michaelson, C.A., 1985, Seismicity and volcanism in the Pacific Northwest: Evidence for the segmentation of the Juan de Fuca plate: *Geophysical Research Letters*, v. 12, p. 215-218.
- \*R White, W.R., and Savage, J.C., 1965, A seismic refraction and gravity study of the Earth's crust in British Columbia: *Seismological Society of America Bulletin*, v. 55, p. 463-486.
- \*R Zucca, J.J., Catchings, R.D., Fuis, G.S., and Mooney, W.D., 1981,



Preliminary report on a seismic refraction study in the Mt. Shasta region of northern California, abs.: Transactions of the American Geophysical Union (EOS), v. 62, p. 1089.

- \*R Zucca, J.J., Fuis, G.S., Milkereit, B., Mooney, W.D., and Catchings, R.D., 1985, Crustal structure of northeastern California from seismic refraction data: Journal of Geophysical Research (in press).

KEY:

R = Refraction

T = Teleseismic P-delay

S = Seismicity, P-to-S conversion, Phase-velocity dispersion, or any similar studies. Not all the Washington and northern California network stations have been plotted on the poster.

\* = stations plotted on poster

+ = see author's bibliography for other studies