

Figure 1. Generalized geologic map of the Dun Glen quadrangle, Pershing County, Nev.

- EXPLANATION
- Qa ALLUVIUM (Quaternary)
 - Tv VOLCANIC ROCKS (Tertiary)
 - Ts SEDIMENTARY ROCKS (Tertiary)
 - Ji GRANITIC INTRUSIVE ROCKS (Jurassic)
 - Ra AULD LANG STYNE GROUP (Triassic)—Argillite, siltstone, quartzite, limestone, and dolomite
 - Ksp STAR PEAK GROUP (Triassic)—Limestone, dolomite, and minor conglomerate and argillite
 - Kk KOIPATO GROUP (Triassic)—Rhyolitic tuffs, flows, and breccias
 - Ms METASEDIMENTARY ROCKS OF UNDETERMINED AGE
 - PPh HAVALLAH SEQUENCE OF SILBERLING AND ROBERTS (1962) (Permian, Pennsylvanian, and Mississippian)—Argillite, quartzite, chert, greenstone, and limestone
 - Mdi INSKIP FORMATION (Mississippian? and Devonian?)—Phyllite, schists, quartzite, conglomerate, metavolcanic rocks, and limestone
 - Ov VALMY FORMATION (Ordovician)—Argillite, chert, greenstone, quartzite, and minor limestone
 - Ch HARMONY FORMATION (Cambrian)—Feldspathic and micaceous quartzite, argillite, and minor limestone
- Contact
- - - Fault—Dashed where approximately located
- - - Thrust fault—Dashed where approximately located; dotted where concealed
x Mine

DISCUSSION

The nine geochemical maps show the distribution and abundance of gold, silver, mercury, arsenic, antimony, copper, lead, zinc, and molybdenum in 114 veins and 45 fissure fillings in the Dun Glen quadrangle, Pershing County, Nev. Samples were collected to determine the distribution and abundance of metals and to outline those areas most favorable for exploration. The samples represent the most highly mineralized rock units at each locality.

The silver and gold elements were determined by semi-quantitative spectrographic analyses by R. F. Godey, K. D. Criss, G. W. Hays, M. J. Hopkins, R. E. Mays, J. M. Hoots, J. H. Berolico, and R. F. Sizem. Determinations of gold, silver, mercury, copper, lead, and zinc by atomic absorption methods and of arsenic and antimony by colorimetric methods were by H. R. Babcock, C. A. Curtis, M. E. Erickson, J. E. Hesser, S. L. Hoffman, R. W. Jones, J. T. Loomis, N. D. McElroy, C. L. Miller, R. W. O'Leary, T. S. Rickard, S. A. Vintola, and A. W. Wells.


All the samples were analyzed for gold and silver by atomic absorption, and spectrographic analyses were obtained for 135 samples. During periods in which mobile field laboratories of the U.S. Geological Survey were operating in northeastern Nevada, determinations of mercury, arsenic, and antimony by atomic absorption and colorimetric methods also was made on 112 of the samples, and determination of copper, lead, and zinc by atomic absorption was made on 74 samples. Lower limits of detection by the spectrographic method are: Au, 200 ppm; Cu, 5 ppm; Pb, 5 ppm; Fe, 10 ppm; Sb, 100 ppm; and Zn, 200 ppm. Lower limits of detection by atomic absorption or colorimetric methods are: Au, 0.02 ppm; Cu, 0.2 ppm; Fe, 0.2 ppm; Pb, 10 ppm; Sb, 5 ppm; and Zn, 5 ppm. Because the lower limits of detection for arsenic, antimony, and zinc are significantly higher in samples analyzed by spectrographic methods, a symbol is used on the geochemical maps of these elements to denote those samples analyzed only spectrographically that contain less than detectable amounts. In the case of lead, the symbol is used on the maps of lead only when the lead content of the sample is less than 10 ppm. In the case of copper, the symbol is used on the maps of copper only when the copper content of the sample is less than 10 ppm. In the case of mercury, the symbol is used on the maps of mercury only when the mercury content of the sample is less than 0.1 ppm. In the case of arsenic, the symbol is used on the maps of arsenic only when the arsenic content of the sample is less than 0.1 ppm. In the case of antimony, the symbol is used on the maps of antimony only when the antimony content of the sample is less than 0.1 ppm. In the case of zinc, the symbol is used on the maps of zinc only when the zinc content of the sample is less than 10 ppm. In the case of molybdenum, the symbol is used on the maps of molybdenum only when the molybdenum content of the sample is less than 5 ppm. The median values for Au, Ag, Cu, Fe, Pb, Sb, Zn, and W are less than the lower limits of detection by spectrographic methods.

Silver's main correlation method was used to indicate the relative strength of association of the various elements with gold and silver. In the veins, lead, zinc, antimony, silver, arsenic, and mercury show the highest positive correlation with gold, and antimony, lead, copper, silver, mercury, and gold show the highest correlation with silver. In the fractures, silver, mercury, iron, antimony, arsenic, and lead show the highest positive correlation with gold, and mercury, gold, lead, manganese, and molybdenum show the highest correlation with silver. Apparent high positive correlation coefficients between tungsten and silver in veins and between lanthanum and both gold and silver in fractures seems to be significant because they are based on relatively few samples.


The geology of the Dun Glen quadrangle has been mapped by Whitebread (1978), and the reader is referred to his map for a more detailed version of the geology. Gold and silver veins in the Dun Glen quadrangle are in the north half of the quadrangle and are mostly in quartz veins in the halos and in the lower units of the Triassic and Permian. Samples of additional small veins in the Dun Glen quadrangle are scattered throughout the area. Some of these veins are from around the perimeters of several granitic intrusive bodies or are from near the Willow Creek thrust. Small quartz and silver veins in the upper plate of the Clinch are the main source of the placer gold in that area. Many samples near the southern trace of the Willow Creek thrust contain arsenic in amounts greater than median values in the lower plate. Many all right samples that contain detectable tungsten are from the Star Peak Group to the northern part of the quadrangle.

REFERENCES


Silberling, M. J., and Roberts, R. J., 1962, Pre-Tertiary stratigraphy and structure of northeastern Nevada, Geol. Soc. America, Special Paper 77, 58 p.
Whitebread, Donald H., 1978, Preliminary geologic map of the Dun Glen quadrangle, Pershing County, Nevada, U.S. Geol. Surv. Open-File Map 78-607.




Gold, in parts per million
+ □ ○ ●
<0.02 0.02-0.1 0.1-1 1.5-10 >10
Figure 2. Distribution of gold in 189 samples from veins and fractures.




Silver, in parts per million
+ □ ○ ●
<0.5 0.5-3 5-100 >100
Figure 3. Distribution of silver in 189 samples from veins and fractures.




Mercury, in parts per million
+ □ ○ ●
<0.02 0.02-0.1 0.1-1 1.5-10 >10
Figure 4. Distribution of mercury in 132 samples from veins and fractures.




Arsenic, in parts per million
+ □ ○ ●
<20 20-50 50-200 >200
Figure 5. Distribution of arsenic in 155 samples from veins and fractures.




Antimony, in parts per million
+ □ ○ ●
<10 10-30 30-200 >200
Figure 6. Distribution of antimony in 155 samples from veins and fractures.




Copper, in parts per million
+ □ ○ ●
<5 5-30 30-200 200-1500 >1500
Figure 7. Distribution of copper in 155 samples from veins and fractures.




Lead, in parts per million
+ □ ○ ●
<10 10-50 50-1000 >1000
Figure 8. Distribution of lead in 155 samples from veins and fractures.



Zinc, in parts per million
+ □ ○ ●
<20 20-50 50-1000 >1000
Figure 9. Distribution of zinc in 155 samples from veins and fractures.



Molybdenum, in parts per million
+ □ ○ ●
<5 5-30 >30
Figure 10. Distribution of molybdenum in 155 samples from veins and fractures.



MAPS SHOWING GEOCHEMICAL DISTRIBUTION OF ELEMENTS IN VEINS AND FRACTURES IN THE DUN GLEN QUADRANGLE, PERSHING COUNTY, NEVADA

BY
DONALD H. WHITEBREAD, JOCELYN A. PETERSON, AND PAUL E. VENUIT

Interior—Geological Survey, Reston, Va., 1978
For sale by Branch of Distribution, U.S. Geological Survey
Box 2526B, Federal Center, Denver, CO 80225