



SR-4

GRADIENT HOLE NO.

8/7/80

DATE OF SURVEY

| DEPTH | °C   | °F    | DEPTH | °C   | °F     | DEPTH | °C   | °F     | DEPTH | °C   | °F     |
|-------|------|-------|-------|------|--------|-------|------|--------|-------|------|--------|
| 0     | 28.2 | 82.76 | 460   | 37.5 | 99.5   | 920   | 57.5 | 135.5  | 1380  | 55.7 | 132.26 |
| 10    | 18.2 | 64.76 | 70    | 38.0 | 100.4  | 30    | 58.0 | 136.4  | 90    | 55.7 |        |
| 20    | 16.8 | 62.24 | 80    | 38.5 | 101.3  | 40    | 58.4 | 137.12 | 1400  | 55.7 |        |
| 30    | 17.1 | 72.78 | 90    | 39.0 | 102.2  | 50    | 58.7 | 137.66 | 10    | 55.7 |        |
| 40    | 17.3 | 63.14 | 500   | 39.4 | 102.92 | 60    | 58.9 | 138.02 | 20    | 55.6 | 132.08 |
| 50    | 17.8 | 64.04 | 10    | 39.8 | 103.64 | 70    | 59.0 | 138.2  | 30    | 55.6 |        |
| 60    | 18.5 | 65.3  | 20    | 40.2 | 104.36 | 80    | 59.9 | 138.38 | 40    | 55.6 |        |
| 70    | 19.1 | 66.38 | 30    | 40.8 | 105.44 | 90    | 59.0 | 138.38 | 50    | 55.6 |        |
| 80    | 19.7 | 67.46 | 40    | 41.1 | 105.98 | 1000  | 59.0 | 138.38 | 60    | 55.6 |        |
| 90    | 20.3 | 68.54 | 50    | 41.5 | 106.7  | 10    | 58.9 | 138.02 | 70    | 55.6 |        |
| 100   | 20.7 | 69.26 | 60    | 41.9 | 107.42 | 20    | 58.8 | 137.84 | 80    | 55.6 |        |
| 10    | 21.0 | 69.8  | 70    | 42.3 | 108.14 | 30    | 58.5 | 137.3  | 90    | 55.5 | 136.9  |
| 20    | 21.3 | 70.34 | 80    | 42.6 | 108.68 | 40    | 58.2 | 136.76 | 1500  | 55.5 | 1      |
| 30    | 21.6 | 70.88 | 90    | 42.9 | 109.22 | 50    | 57.9 | 136.22 |       |      |        |
| 40    | 22.1 | 71.78 | 600   | 43.2 | 109.76 | 60    | 57.6 | 135.68 |       |      |        |
| 50    | 22.4 | 72.32 | 10    | 43.5 | 110.3  | 70    | 57.3 | 135.14 |       |      |        |
| 60    | 22.8 | 73.04 | 20    | 43.8 | 110.84 | 80    | 57.0 | 134.6  |       |      |        |
| 70    | 23.2 | 73.76 | 30    | 44.1 | 111.38 | 90    | 56.8 | 134.24 |       |      |        |
| 80    | 23.7 | 74.66 | 40    | 44.4 | 111.92 | 1100  | 56.6 | 133.88 |       |      |        |
| 90    | 24.1 | 75.38 | 50    | 45.0 | 113.0  | 10    | 56.4 | 133.52 |       |      |        |
| 200   | 24.6 | 76.28 | 60    | 45.8 | 114.44 | 20    | 56.2 | 133.16 |       |      |        |
| 10    | 25.1 | 77.18 | 70    | 46.1 | 114.98 | 30    | 56.2 | 133.16 |       |      |        |
| 20    | 25.6 | 78.08 | 80    | 46.4 | 115.52 | 40    | 56.3 | 133.34 |       |      |        |
| 30    | 26.1 | 78.98 | 90    | 46.9 | 116.42 | 50    | 56.3 | 133.34 |       |      |        |
| 40    | 26.6 | 79.88 | 700   | 47.4 | 117.32 | 60    | 56.3 | 133.34 |       |      |        |
| 50    | 27.1 | 80.78 | 10    | 47.9 | 118.22 | 70    | 56.2 | 133.16 |       |      |        |
| 60    | 27.6 | 81.68 | 20    | 48.4 | 119.12 | 80    | 56.1 | 132.98 |       |      |        |
| 70    | 28.1 | 82.58 | 30    | 48.9 | 120.02 | 90    | 56.1 | 132.98 |       |      |        |
| 80    | 28.6 | 83.48 | 40    | 49.4 | 120.92 | 1200  | 56.0 | 132.98 |       |      |        |
| 90    | 29.0 | 84.2  | 50    | 49.9 | 121.82 | 10    | 56.0 | 132.98 |       |      |        |
| 300   | 29.4 | 84.92 | 60    | 50.4 | 122.72 | 20    | 55.9 | 132.62 |       |      |        |
| 10    | 29.8 | 85.64 | 70    | 50.9 | 123.62 | 30    | 55.9 | 132.62 |       |      |        |
| 20    | 30.2 | 86.36 | 80    | 51.4 | 124.52 | 40    | 55.9 | 132.62 |       |      |        |
| 30    | 30.6 | 87.08 | 90    | 51.9 | 125.42 | 50    | 55.9 |        |       |      |        |
| 40    | 31.1 | 87.98 | 800   | 52.4 | 126.32 | 60    | 55.9 |        |       |      |        |
| 50    | 31.6 | 88.88 | 10    | 52.9 | 127.22 | 70    | 55.9 |        |       |      |        |
| 60    | 32.1 | 89.78 | 20    | 53.3 | 127.94 | 80    | 55.8 | 132.44 |       |      |        |
| 70    | 32.8 | 91.04 | 30    | 53.7 | 128.66 | 90    | 55.8 |        |       |      |        |
| 80    | 33.3 | 91.94 | 40    | 54.1 | 129.38 | 1300  | 55.8 |        |       |      |        |
| 90    | 33.9 | 93.02 | 50    | 54.5 | 130.1  | 10    | 55.8 |        |       |      |        |
| 400   | 34.5 | 94.1  | 60    | 54.9 | 130.82 | 20    | 55.8 |        |       |      |        |
| 10    | 35.0 | 95.0  | 70    | 55.3 | 131.54 | 30    | 55.8 |        |       |      |        |
| 20    | 35.5 | 95.9  | 80    | 55.7 | 132.26 | 40    | 55.8 |        |       |      |        |
| 30    | 36.0 | 96.8  | 90    | 56.1 | 132.98 | 50    | 55.7 | 132.26 |       |      |        |
| 40    | 36.5 | 97.7  | 900   | 56.5 | 133.7  | 60    | 55.7 |        |       |      |        |
| 50    | 37.0 | 98.6  | 10    | 57.0 | 134.6  | 70    | 55.7 |        |       |      |        |

Log No. ....  
 Permit No. ....  
 Basin: .....

**WELL DRILLERS REPORT**

Please complete this form in its entirety

1. OWNER Southland Royalty Company ADDRESS 1600 First National Bank Building  
Fort Worth, Texas 76102

2. LOCATION NW 1/4 SE 1/4 Sec 22 T. 23 N 8 R. 38 E Churchill County  
 PERMIT NO. Thermal Gradient Hole No. SR-4

|   |  |   |
|---|--|---|
| 3. TYPE OF WORK   | 4. PROPOSED USE  | 5. TYPE WELL  |
| New Well <input checked="" type="checkbox"/> Recondition <input type="checkbox"/> | Domestic <input type="checkbox"/> Irrigation <input type="checkbox"/> Test <input checked="" type="checkbox"/> | Cable <input type="checkbox"/> Rotary <input checked="" type="checkbox"/> |
| Deepen <input type="checkbox"/> Other <input type="checkbox"/>                    | Municipal <input type="checkbox"/> Industrial <input type="checkbox"/> Stock <input type="checkbox"/>          | Other <input type="checkbox"/>  |

| 6. LITHOLOGIC LOG |              |      |      |            |
|-------------------|--------------|------|------|------------|
| Material          | Water Strata | From | To   | Thick-ness |
| Top Soil          |              | 0    | 2    | 2          |
| Rhyolite          |              | 2    | 310  | 308        |
| Fault Gouge       |              | 310  | 330  | 20         |
| Granite           |              | 330  | 340  | 10         |
| Rhyolite          |              | 340  | 390  | 50         |
| Granite           |              | 390  | 420  | 30         |
| Rhyolite          |              | 420  | 530  | 110        |
| Granite           |              | 530  | 650  | 120        |
| Sandstone         |              | 650  | 660  | 10         |
| Fault Gouge       |              | 660  | 670  | 10         |
| Rhyolite          |              | 670  | 930  | 260        |
| Shale             |              | 930  | 940  | 10         |
| Rhyolite          |              | 940  | 1150 | 210        |
| Shale             |              | 1150 | 1160 | 10         |
| Rhyolite          |              | 1160 | 1180 | 20         |
| Fault Gouge       |              | 1180 | 1190 | 10         |
| Shale             |              | 1190 | 1210 | 20         |
| Rhyolite          |              | 1210 | 1240 | 30         |
| Shale             |              | 1240 | 1280 | 40         |
| Granite           |              | 1280 | 1290 | 10         |
| Shale             |              | 1290 | 1470 | 180        |
| Rhyolite          |              | 1470 | 1500 | 30         |

8. WELL CONSTRUCTION

Diameter hole 9-7/8 inches Total depth 1,500 feet  
 Casing record 7-5/8" 0 - 151 Ft.  
 Weight per foot 20 Lbs. Thickness.....  
 Tubing 1" 0 - 1,503 Ft.  
 Diameter From To  
6-1/4 inches 160 feet 1,260 feet  
6-1/8 inches 1,260 feet 1,500 feet  
 .....inches .....feet .....feet  
 .....inches .....feet .....feet  
 .....inches .....feet .....feet  
 .....inches .....feet .....feet

Surface seal: Yes  No  Type Cement  
 Depth of seal 1" Tubing Cemented 32 feet  
 Gravel packed: Yes  No   
 Gravel packed from 1,500 feet to 1,468 feet

Perforations: None  
 Type perforation.....  
 Size perforation.....  
 From.....feet to.....feet  
 From.....feet to.....feet  
 From.....feet to.....feet  
 From.....feet to.....feet  
 From.....feet to.....feet

Date started June 8, 19 80  
 Date completed June 27, 19 80

| 7. WELL TEST DATA |        |           |                  |
|-------------------|--------|-----------|------------------|
| Pump RPM          | G.P.M. | Draw Down | After Hours Pump |
|                   |        |           |                  |
|                   |        |           |                  |
|                   |        |           |                  |
|                   |        |           |                  |

| BAILER TEST |           |       |
|-------------|-----------|-------|
| G.P.M.      | Draw down | hours |
|             |           |       |
|             |           |       |
|             |           |       |


9. No Water WATER LEVEL

Static water level 0 Feet below land surface 0  
 Flow.....G.P.M.....  
 Water temperature.....° F. Quality.....

10. DRILLERS CERTIFICATION

This well was drilled under my supervision and the report is true to the best of my knowledge.

Name Jerrold D. Christiansen  
 Address 557 Ely Ave., Ely, Nevada 89301  
 Nevada contractor's license number 14790  
 Nevada driller's license number 641  
 Signed Jerrold D. Christiansen  
 Date July 5, 1980



Southland Royalty Company  
NATURAL RESOURCES DISTRICT  
FORT WORTH, TEXAS

DIXIE VALLEY NEVADA

LOCATION MAP FOR  
PROPOSED GRADIENT HOLE  
SR-4 SPRING CREEK

D.S. McMurdie      April 10, 1980  
Scale: 1" = 1 Mile

SUMMARY OF LITHOLOGIES AND ALTERATION IN SR-4  
DIXIE VALLEY, CHURCHILL COUNTY, NEVADA

M. J. Sweeney, September 1980

INTRODUCTION

Samples of washed, rotary cuttings taken every ten feet from drill-hole SR-4 were examined under a stereomicroscope. A thin-section of the cuttings was described from every 100 feet throughout the 1500 foot length of SR-4. Detailed descriptions of the cuttings and thin-sections are attached to this report. Photomicrographs of the thin-sections are also attached. X-ray diffraction patterns of two tuff samples are enclosed.

LITHOLOGIES

SR-4 intersects 900 feet of rhyolitic, vitric-crystal-lithic tuffs and 600 feet of carbonate rocks, limestones and dolomites, which are interbedded with narrow intervals of calcareous siltstones and sandstones. The tuffs overly the carbonate rocks.

Rhyolitic tuffs: The tuffs were originally composed largely of volcanic glass which was deposited as vitric shards, dust and pumice fragments. Shards and dust occurring in about subequal amounts usually comprise 75 to 85% of the tuffs and pumice fragments 5 to 15%. Devitrification of the volcanic glass is variably developed throughout this volcanic section.

Crystal clasts of sanidine, plagioclase and biotite occur throughout the tuff section. The abundances of the crystal clasts are variable; biotite is very rare in some intervals. Sanidine clasts usually comprise 5 to 7% of the tuff, plagioclases 1 to 5% and biotites 0 to 3%. Quartz crystal clasts were seen only in the samples at the bottom of the volcanic section. The feldspar crystal clasts usually range between 0.5 and 2 mm in length. They were originally subhedral to euhedral in shape, but most have been broken prior to deposition. The biotites, 0.2 to 1 mm across, exhibit very dark brown pleochroism, suggesting that they are very Fe-rich. Magnetite is locally exsolved from biotite.

Lithic clasts comprise 5 to 20% of the tuff; their abundance is variable throughout this section. Clasts of andesites, basalts, latites, quartzite and carbonate were seen; the first three types are the most ubiquitous. The lithic clasts observed are less than 10 mm in diameter.

The entire 900' section of this volcanic unit is very similar. Primary mineralogy and textures were virtually identical. One of the major variations seen in the tuff now is hardness. Most of the tuff is relatively soft and also gritty to the touch. However, between the depths of 520' and 640' the tuff is hard, exhibits a vitreous luster and fractures conchoidally. Examination of the thin-section at 590'-600' shows that the vitric components have been well compacted and welded. Devitrification to Kspar and quartz is complete. This well welded interval may be the base of a thick ignimbrite unit. Other intervals in the tuff have also been well devitrified, but no other intervals appear to have been as intensely compacted as the interval between 520 and 640 feet.

Dolomites, Limestones, Sandstones, Siltstones: Beneath the tuffs, SR-4 intersected dolomites, limestones (both of which are locally cherty), and calcareous sandstones and siltstones.

Cherty dolomites occur in the first 150 feet below the tuffs. Limestones occur in the next 60 feet, calcareous siltstones and sandstones in the next 110 feet, and then silty, locally cherty or argillaceous, limestones occur in the lowermost 280 feet of the drill hole.

The dolomites and limestones are dominantly fine-grained; they are micritic. Fossils were seen rarely.

The siltstones and sandstones are well sorted; the calcite in them is of detrital origin.

#### ALTERATION

Rhyolitic tuffs: Hydrothermal alteration in the tuffs is seen mainly in the glass and in the mafic minerals in the lithic clasts. Only rarely have crystal clasts suffered any alteration.

The volcanic glass has been replaced most commonly by clays and lesser calcite. Local occurrences of jarosite (90'-100'), gypsum (40'-50', 90'-100') and manganese-oxides (10'-20', 180'-240') were observed. Trace amounts of hematite and goethite occur throughout the tuff. These limonites replace disseminated and vein magnetite and sulfides; it was not often possible to determine which of these was the precursor of the limonites. Trace amounts of zeolites may occur in vugs in the upper portion of the tuff.

The extent and type of clay replacement is highly variable. Replacement may range between 5 and 100%. The most intensely clay-altered intervals are easily identified by observing the degree of rock disaggregation during washing; easily disaggregated rocks are intensely clay-replaced. The clay types present appear to be variable. X-ray diffraction analysis of clays in the sample at 590'-600' gives a strong 12.6A<sup>0</sup> peak possibly indicative of the presence of a mixed layer clay or possibly of hydrobasaluminite. X-ray analysis of sample 930'-940' indicates that the abundant white clay is montmorillonite; this rock disaggregates completely on wetting.

The present color of the tuffs is controlled by the type and distribution of limonites and/or clays present. Red, pink and orange colored tuffs are the result of limonite staining. Bright green tuffs contain a bright kelly-green clay mineral, probably celadonite.

In the interval between 860' and 900', the tuff has been brecciated. The spaces among the fragments have been filled with microcrystalline quartz, celadonite (?) and calcite.

Only rarely was fresh pyrite seen in the tuffs (at 520' to 530').

Slickensided chips derived from fault zones are sporadically present throughout the tuff; the gouge chips are usually white and weakly stained with hematite.

Dolomites, Limestones, Sandstones, Siltstones: Evidence of hydrothermal alteration in the carbonates and sandstones is largely confined to veins. Thin calcite veins, less than 1-2 mm wide, with and without pyrite occur throughout this interval. Pyrite also is disseminated in these rocks; it usually occurs as very fine-grained crystals forming less than 0.2% of the rock. This disseminated pyrite may be syngenetic in origin. Traces of magnetite also occur as disseminations. Total pyrite content throughout the carbonate-siltstone interval never exceeds 0.3 volume % and is usually less.

Evidence of faulting occurs throughout the carbonate-siltstone interval. Slickensided chips are present in nearly every sample of every rock type. Weak hematite stains are present on these slickensided surfaces.

THIN-SECTION DESCRIPTIONS

OF SR-4 SAMPLES



SR-4 90'-100' Devitrified Vitric-Crystal Tuff

This tuff contains sanidine crystal clasts, 0.3 to 1 mm long, most of which are angular, broken fragments; they form 5-7% of rock. These sanidines are totally unaltered. A few phenocrysts of albite-twinned plagioclase are present, 1-2%; they are unaltered. The groundmass is composed of devitrified glass shards; the shard outlines are preserved by axiolitic devitrification structures. Spherulitic devitrification structures occur occasionally. Traces of zeolite minerals may be present in some vugs.

Jarosite occurs in this tuff; it pseudomorphically replaces individual shards and occurs in irregular blebs disseminated throughout rock. Jarosite chips as large as 0.5 mm across are present. There is no evidence as to the origin of the jarosite; it does not occur in pyrite-like pseudomorphs. Jarosite also lines vugs.

Traces of hematite occur as disseminations (after magnetite). A few, free chips of gypsum occur in the section.

PRIMARY MINERALS:

|             |        |   |
|-------------|--------|---|
| Quartz      | 10-30% | Anhedra! to subhedra!; occurs only in groundmass. Grain size 0.01 mm or less. |
| Sanidine    | 5-7%   | Crystal clasts: 0.2 to 1 mm long; unaltered.                                  |
|             | 50-70% | Groundmass: anhedra! to subhedra!; grain size 0.01 mm or less.                |
| Plagioclase | 1-2%   | Crystal clasts: 0.2 to 1 mm long; unaltered; albite-twinned.                  |

SECONDARY MINERALS:

|            |      |   |
|------------|------|---|
| Jarosite   | 3-5% | Pseudomorphically replaces vitric shards and biotites?; also in irregular blebs disseminated throughout tuff. Very fine-grained crystals, 0.005 mm or less. |
| Gypsum     | tr   | Free crystals in sample.  |
| Zeolites?? | 1-5% | In vugs; minerals with birefringence up to first-order red occur in vugs. Colorless; lath shaped.   |
| Hematite   | tr   | Disseminated.   |

SR-4 190'-200' Devitrified Vitric-Crystal Tuff

This tuff originally was composed largely of vitric shards; it also contains crystal clasts of: sanidine, 0.3-0.6 mm, 5-7%; plagioclase, 0.3-1 mm long, albite-twinned (An<sub>30-40</sub>), 2-3%; and of biotite, 0.3-1 mm, very dark brown pleochroism, 1-3%. The phenocrysts (crystal clasts) are unaltered. The glass shards have devitrified to quartz, Kspar and clays. Irregular aggregates of Mn-oxides are distributed throughout rock; they form 2-3% of rock.

PRIMARY MINERALS:

|        |        |  |
|--------|--------|--|
| Quartz | 10-35% | Anhedra! to subhedra!; occurs in groundmass intergrown with Kspar. |
| Kspar  | 5-7%   | Sanidine crystal clasts: angular, crystal fragments.               |
|        | 50-70% | Groundmass: anhedra! to subhedra!; grain size less than 0.01 mm.   |

|             |      |  |
|-------------|------|--|
| Plagioclase | 2-3% | Crystal clasts: unaltered.                   |
| Biotite     | 1-3% | Crystal clasts: very dark brown pleochroism. |

SECONDARY MINERALS:

|           |       |  |
|-----------|-------|--|
| Clay      | 5-20% | Very fine-grained; occurs in matrix.   |
| Mn oxides | 2-3%  | Black reflecting; brown streak, disseminated throughout tuff in clots up to 0.1 mm across. |

SR-4 290'-300' Vitric-Crystal Tuff

This tuff is composed of: glass shards and even finer-grained, glass dust both of which form 80% of rock; glassy pumice fragments, 0.5 to 4 mm, 10% of rock; crystal clasts of sanidine, 0.1-0.5 mm, 5-7% of rock; crystal clasts of plagioclase, 0.1-0.5 mm, 1-2%; and crystal clasts of very dark brown biotite, 0.1-1.5 mm across, 2-4%. A very few, tiny lithic fragments of very fine-grained basalt are also present. Magnetite is disseminated throughout rock; magnetite replaces some of the biotite phenocrysts.

The glass particles have been less than 50% devitrified. Devitrification of pumice fragments is more advanced than that of vitric shards and ash. Spherulitic devitrification structures occur most commonly in pumice fragments. Devitrification products include Kspar, quartz and clay.

Calcite occurs in narrow veins (less than 0.1 mm wide); it also replaces feldspar phenocrysts, pumice fragments and vitric shards. Traces of celadonite (bright green) are disseminated throughout rock.

PRIMARY MINERALS:

|             |        |   |
|-------------|--------|---|
| Quartz      | 10-20% | Groundmass: devitrification product of glass. Very fine-grained.                            |
| Kspar       | 5-7%   | Crystal clasts.   |
|             | 10-20% | Groundmass: devitrification product of glass. Very fine-grained.                            |
| Plagioclase | 1-2%   | Crystal clasts: 0.2-1 mm long; 10-90% (average 15% replaced by calcite).                    |
| Biotite     | 2-4%   | Crystal clasts: very dark brown; 0.1-1 mm across; locally, partially replaced by magnetite. |
| Magnetite   | 0.5-1% | Disseminated; also common in biotite sites.   |
| Glass       | 30-40% | Vitric shards, ash, pumice fragments.   |

SECONDARY MINERALS:

|            |       |   |
|------------|-------|---|
| Celadonite | tr-1% | Bright green; disseminated throughout groundmass. |
| Calcite    | 3%    | Vein and disseminated.                            |
| Clay       | 5-20% | Alteration product of vitric clasts.              |

SR-4 390'-400' Vitric-Lithic-Crystal Tuff

This tuff is composed mostly of glass which includes shards, pumice fragments and very fine-grained vitric dust; glass forms 65 to 75% of the tuff. The fine dust has been altered to illite/sericite. Most of the shards and pumice fragments are unaltered. Crystal clasts of sanidine are present; they are

completely unaltered and form 7-10% of rock. Crystal clasts of unaltered plagioclase comprise 1-2% of rock. Crystal clasts of biotite constitute 2-4% of the tuff; the biotite exhibits dark brown pleochroism and is not altered.

Lithic fragments of andesite, basalt(?), quartzite(?), latite and carbonate occur; they are listed in order of abundance. The lithic fragments form about 10-20% of the tuff; most of these fragments are less than 5 mm across. The plagioclases, both phenocrysts and groundmass laths, are unaltered in these fragments. Original mafic minerals, excepting biotite phenocrysts, have been replaced by Fe-oxides and clays (montmorillonite?); biotite phenocrysts are unaltered. Primary magnetite is disseminated throughout the lithic fragments; it forms 0.5-1% of the fragments.

In one latite fragment, the feldspar phenocrysts have been replaced by epidote.

SECONDARY MINERALS:

|                                      |        |  |
|--------------------------------------|--------|--|
| Illite/sericite                      | 10-30% | Alteration product of glass dust.            |
| Montmorillonite?                     | 5-7%   | Alteration product of pyroxenes/hornblendes. |
| Fe-oxides<br>(goethite/<br>hematite) | 1%     | Alteration product of pyroxenes/hornblendes. |

SR-4 490'-500' Clay-altered Vitric-Crystal-Lithic Tuff

The tuff contains crystal clasts of sanidine which occur as either broken crystal fragments or euhedral, Carlsbad-twinned crystals up to 3 mm in length; they form 5 to 7% of rock. These sanidines are completely unaltered. Biotite crystals are rare, forming less than 0.5% of the rock. Plagioclase crystal clasts, up to 4 mm in length, form less than 1% of the tuff; they are completely unaltered. Lithic fragments of other tuffs and andesites are usually less than 2 mm across; they form about 3% of the tuff.

The groundmass or matrix of this tuff was originally composed of vitric shards and dust. The original glass is now composed mostly of fine-grained quartz and Kspar; locally devitrified shards exhibit axiolitic structures. Original vitric dust is locally replaced by fine-grained clays, celadonite?(green) and illite(colorless). Celadonite also occurs in aggregates filling gas bubbles and possibly replacing primary biotites. Minor amounts of carbonate are irregularly distributed throughout groundmass.

SECONDARY MINERALS:

|            |        |  |
|------------|--------|--|
| Illite(?)  | 10-30% | Replaces glass in matrix.  |
| Celadonite | 5%     | Occurs in groundmass; also fills vugs and may replace biotite phenocrysts. |
| Carbonate  | 1-3%   | Occurs in irregular patches throughout groundmass.                         |

SR-4 590'-600' Devitrified Welded Vitric-Crystal-Lithic Tuff

This tuff contains crystal clasts of: sanidine, 0.2-2mm long, 3 to 5% of rock; plagioclase, 0.2 to 1 mm long, 1-2% of rock; and of biotite, 0.1-0.5 mm across, 0.5-1% of rock, exhibits very dark brown pleochroism. None of these crystal clasts have suffered any alteration.

The groundmass was once composed of flattened pumic fragments, vitric shards

and dust. The glass has been devitrified to fine-grained, anhedral quartz, feldspar and very fine-grained disseminated magnetite. Clay also (celadonite? =green; montmorillonite?=orange) occurs in the groundmass; clay spottily replaces 10 to 40% (average 20%) of groundmass.

SECONDARY MINERALS:

|                  |       |   |
|------------------|-------|---|
| Celadonite       | 7-10% | Alteration product of glass; bright kelly green.                            |
| Montmorillonite? | 7-10% | Alteration product of glass; dirty orange.                                  |
| Pyrite           | 0.1%  | Vein and disseminated. Occurs in a vein in an intensely clay-replaced chip. |

SR-4 690'-700' Vitric-Lithic-Crystal Tuff

This tuff is composed dominantly of vitric shards and dust. The dust has been partially devitrified to clay (sericite) and to very fine-grained orthoclase and quartz. It also contains crystal clasts of: sanidine, up to 2 mm long, 3 to 5% of rock; plagioclase, up to 2 mm long--these plagioclase crystals are strongly zoned, they form 5 to 7% of rock; and of biotites which exhibit very dark brown pleochroism--they form 1-2% of rock. None of the crystal clasts has suffered any alteration.

Lithic clasts, fine-grained andesites/basalts?, are usually less than 2 mm in diameter. Such clasts form 5-10% of the tuff. The feldspars are unaltered; mafic minerals have been replaced by clays (montmorillonite?, illite?) and Fe-oxides (goethite, hematite).

SECONDARY MINERALS

|                   |        |   |
|-------------------|--------|---|
| Sericite/clay     | 5-20%  | Replaces glass dust; also replaces mafic minerals in lithic clasts.                 |
| Goethite/hematite | 0.5-1% | Alteration product of mafic minerals; some primary magnetite is partially oxidized. |
| Carbonate         | 2-5%   | Locally floods rock--replaces all components.                                       |

SR-4 790'-800' Devitrified Welded Vitric-Crystal Tuff

This rock was formed as a welded vitric-crystal tuff. The glass has been completely devitrified. Spherulites, usually about 0.5 mm across, are common devitrification structures. Axiolitic structures developed in shards are also common.

Crystal clasts of sanidine form 3-5% of rock; they are unaltered. Clasts of plagioclase(?) formed 1-3% of rock; they have been replaced by celadonite and illite. Biotites have also been replaced by celadonite. Primary magnetite is disseminated in trace amounts. A weak stain of hematite/goethite is disseminated throughout many chips.

Lithic clasts of latite/andesite form less than 50% of sample.

This tuff has been subjected to cataclasis or brecciation prior to devitrification. The breccia fragments are 0.1 to 2 mm across. The spaces among the fragments have been filled with microcrystalline quartz, celadonite and calcite.

SECONDARY MINERALS:

|                   |       |  |
|-------------------|-------|--|
| Sericite          | 2-5%  | Replaces plagioclase clasts.   |
| Celadonite        | 3-5%  | Replaces mafic minerals; fills vugs; occurs with chert in interbreccia spaces. |
| Quartz            | 1-10% | Fills spaces among breccia clasts; microcrystalline or cherty.                 |
| Calcite           | 1-2%  | Occurs in interbreccia spaces; occurs in post-brecciation fractures.           |
| Hematite/goethite | 1-2%  | Stains rock.   |

SR-4 890'-900' Devitrified Crystal-Vitric Tuff

This is the first tuff sample from this drill hole to contain quartz crystal clasts. Several of those present are rounded and embayed by resorption. The quartz clasts are usually about 1 mm in diameter; they comprise about 5% of the tuff. Crystal clasts of sanidine form about 10% of rock; those of plagioclase about 1%. Biotite phenocrysts occur in trace amounts. None of these phenocrysts have suffered any alteration.

The groundmass once composed of vitric shards and dust is now composed mostly of anhedral quartz and Kspar; spherulites occur occasionally. Clay alteration (sericite/illite) of groundmass is weak to moderate. Trace amounts of magnetite are disseminated throughout groundmass.

Calcite and rarely microcrystalline quartz occur as fracture fillings.

Lithic fragments of andesites/basalts form less than 5% of this rock.

SECONDARY MINERALS:

|               |        |   |
|---------------|--------|---|
| Sericite/clay | 10-25% | Very fine-grained material occurs in groundmass.            |
| Calcite       | 3-7%   | Vein and disseminated.                                      |
| Quartz        | 1-3%   | Microcrystalline; occurs in fracture fillings with calcite. |

SR-4 990'-1000' Carbonates (limestones/dolomites), Cherts and Devitrified Welded Tuffs

This thin-section contains chips of fine- to medium-grained, recrystallized carbonates; these chips form 50-60% of the section. A few of these chips contain disseminated chert. Pyrite fills fractures in a few of the carbonate chips.

The next most abundant rock type is chert; most of these chips are carbonate-free. Chert comprises 30-40% of rock chips in this thin-section.

Quartz "eye" tuff constitutes 5-10% of the chips. The matrix and plagioclase phenocrysts are weakly clay (illite)-altered.

SECONDARY MINERALS:

|        |      |  |
|--------|------|--|
| Pyrite | tr   | Fills or coats fractures in carbonate rocks. |
| Illite | 2-5% | Alteration product of tuff.                  |

SR-4 1090'-1100' Limestones, Sandy Limestones and Cherty Limestones

This sample contains pure limestones, sandy limestones and cherty limestones.

About 30% of the chips are composed of pure carbonate, mostly calcite judging by how reactive the rock is to cold, dilute HCL. The calcite is mostly very fine-grained (micritic). A few chips contain relict fossil structures (bryozoans?, shell fish). A small proportion of these chips have been recrystallized to medium-grained marbles.

About 50% of the limestone chips contain 10-15% silt and fine-sand sized quartz. Disseminated pyrite occurs most commonly in these chips; pyrite occurs in trace amounts up to 0.5 volume %.

About 15% of the limestone chips contain chert which occurs as microcrystalline quartz or fibrous chalcedony. Chert occurs in limestones with and without sand. Pyrite also occurs in cherty limestones.

Thin calcite veins occur in all of the above described chips.

SECONDARY MINERALS:

|         |      |                        |
|---------|------|------------------------|
| Pyrite  | 0.5% | Vein and disseminated. |
| Calcite | 2-3% | Vein.                  |

SR-4 1190'-1200' Calcareous Siltstone

This rock is composed largely of angular, detrital quartz grains which average 0.04 mm in size. Quartz comprises about 50% of the rock. Silt-sized orthoclase grains form about 5% of the rock. The remainder of the rock is composed of calcite. This calcite is also fine-grained; it is disseminated throughout the sandstone. The calcite is usually orange or brown in color, presumably from exsolved iron. Locally illite is common as a matrix for quartz grains; it may form 5-15% of rock.

Thin calcite and calcite-quartz veins crosscut the sandstone. Vein calcite is colorless. Euhedral cubes of hematite-replaced pyrite (or magnetite) up to 0.1 mm across are disseminated through the sandstone.

SECONDARY MINERALS:

|         |      |                                     |
|---------|------|-------------------------------------|
| Pyrite  | tr   |                                     |
| Calcite | 1-2% | In veins by itself and with quartz. |
| Quartz  | 1-2% | In veins with calcite.              |

SR-4 1290'-1300' Silty Limestones, Argillaceous Calcareous Siltstones and Limestones

Silty limestone is the most common rock type in this sample. Angular, silt-sized, quartz grains comprise 5 to 30% of these limestones. Most of the calcite in these chips is colorless in thin-section; however, orange or brown-stained calcite constitutes up to 30% of some rock chips.

The argillaceous rocks contain 10 to 40% clays (illite? and kaolinite?). Silt and fine-grained calcite, present in subequal amounts, are the other constituents of the clayey rocks; such chips comprise about 25% of this sample.

A few pure calcite chips are present. The calcite in these chips is mostly very fine-grained (micritic) and colorless.

Traces of pyrite, both fresh and oxidized, occur in silty limestones and argillaceous siltstones. Pyrite occurs in veins and as disseminations.

SECONDARY MINERALS:

Pyrite/hematite 0.1-0.2% Vein and disseminated.

SR-4 1390'-1400' Silty Limestones

These rocks are composed dominantly of very fine-grained calcite. Angular, detrital, silt-sized grains of quartz and feldspar form 5 to 20% of these limestones. Clay (illite, kaolinite) can comprise near 15% of the limestone; usually clay are present in small amounts.

Trace amounts of magnetite are disseminated throughout the limestones. Pyrite occurs in trace amounts in calcite veins.

SECONDARY MINERALS:

Calcite 1-2% Occurs in veins.  
Pyrite tr Occurs in veins with calcite.

SR-4 1490'-1500' Limestones

The rocks in this sample are micritic limestones which contain 2 to 20% silt. Clay comprises 5 to 25% (average 10% or less) of these limestones.

Thin calcite veins, some with goethite (after pyrite?), occur. Traces of very fine-grained pyrite and magnetite are disseminated throughout these limestones; these disseminated minerals are probably syngenetic.

SECONDARY MINERALS:

Calcite 2-3% Veins, sometimes with goethite (after pyrite?).  
Pyrite/goethite 0.2% Vein and disseminated; the disseminated pyrite may be of syngenetic origin.

WELL: SR-4

## GEOLOGICAL REPORT

PAGE: 1 of 10LOCATION: Dixie Valley, Churchill Co., Nevada WELL CUTTINGS SAMPLE DESCRIPTIONDATE: September 1930

ELEVATION: \_\_\_\_\_

DEPTH: 0 TO 150'

HOLE SIZE: \_\_\_\_\_

EXAMINED BY: M. J. Sweeney

| DEPTH | COLOR                                       | SAMPLE DESCRIPTION  |   |
|-------|---|---|---|
|       |   | PRIMARY   | SECONDARY   |
| feet  |   |   |   |
| 0     | Mottled pale green and pale orange-pink     | Tuff: Rock contains crystal clasts of sanidine and biotite; they range between 1 and 3mm in size. Feldspars form 5 to 10 volume % of rock; biotites 1-3%. The groundmass is soft and gritty to the touch. Dark green spots (0.2-1mm across) are possibly concentrations of celadonite which locally replaces phenocrysts. | Rare veins, 1-2mm wide, contain a soft, pink mineral -- a zeolite? Goethite occurs in 1/2mm wide veins; it also occurs as disseminations (after mgf/p)  |
| 10    | Pale pink, pale green                       | Tuff: Very similar to above sample. Less celadonite occurs, therefore less green.   | Manganese oxides coat some fractures -- this black material is very fine-grained. It has a brown streak. Traces of goethite/hematite as disseminations. |
| 20    | Very pale pinkish buff                      | Tuff: Very similar to above sample.   | Mn-oxides, traces. Traces of disseminated goethite/hematite.  |
| 30    | Very pale buff; mottled pale green and pink | Tuff: Very similar to above sample. Rock contains crystal clasts of sanidine and biotite in a soft matrix.  | Traces of goethite/hematite occur as disseminations and as coatings on fractures. Disseminated oxides may be in mafic mineral sites.                    |
| 40    | Pale green                                  | Tuff: Very similar to above sample.   | Traces of goethite in vugs. "Satin" spar gypsum crystals are present; they are rare. Disseminated celadonite (bright green clay) occurs.                |
| 50    | Pale green                                  | Tuff: same as above. As in all previous samples, the sanidine crystals are very clear; they have not suffered any alteration.   | Goethite and/or hematite occurs in vugs, 0.2-0.5%. A few gypsum crystals are present.   |
| 60    | Pale green                                  | Tuff: Same rock as above.   | Tiny aggregates of crystalline goethite and of earthy hematite are disseminated throughout rock, 0.5-0.75%.   |
| 70    | Pale green                                  | Tuff: Same rock as above. Crystal clear clasts of sanidine are present; 5-15 volume % of rock; clasts of very black biotite occur in trace amounts. Some original biotite may have been replaced by celadonite and limonites.   | Traces of disseminated goethite and hematite. Tiny clots of green celadonite also occur as disseminations.  |
| 80    | Very pale green                             | Tuff: Similar to above sample.  | Same as above.  |
| 90    | Very pale buff and green                    | Tuff: Contains phenocrysts of sanidine. No biotite present. See thin-section description. Original biotites probably replaced by jarosite; jarosite also replaces vitric shards.  | Contains almost no goethite or hematite; that present occurs as thin films on slickensided surfaces. Contains disseminated jarosite.                    |
| 100   | Very pale buff and green                    | Tuff: Same as above. No biotite present.  | Traces of goethite/hematite. Traces of celadonite.  |
| 110   | Very pale buff and green                    | Tuff: Same as above.  | Same as above.  |
| 120   | Pale buff                                   | Tuff: Same as above.  | Traces of hematite disseminated throughout rock.  |
| 130   | Pale buff with tinges of green.             | Tuff: Same as above. Contains crystal clasts of sanidine, 0.5 to 3mm in length, 5 to 10% of rock.   | Traces of hematite. Traces of calcite.  |
| 140   | Pale pink                                   | Tuff: Same as above. Rock rapidly disaggregates on wetting. Sanidines are unaltered.  | Contains traces of calcite, traces of hematite. Rock rapidly disaggregates on wetting - groundmass is clay-rich.  |
| 150   |   |   |   |



WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION



PAGE: 2 of 10  
 DATE: September 1980  
 DEPTH: 150' TO 300'  
 EXAMINED BY: M. J. Sweeney

| DEPTH | COLOR                                     | SAMPLE DESCRIPTION   |  |
|-------|---|--|--|
|       |   | PRIMARY  | SECONDARY  |
| feet  |   |  |  |
| 150   | Pale pink, green                          | Tuff: Pink tuff disaggregates on wetting; it is very clay-rich; green tuff remains intact. Traces of unoxidized magnetite remain.  | Traces of hematite. Clay replacement of vitric shards is locally intense.  |
| 160   |   |  |  |
| 160   | Pale pink, green.                         | Tuff: Same as above.   | Same as above.   |
| 170   |   |  |  |
| 170   | Pale brown-pink with green patches.       | Tuff: looks very different than previous sample. This rock remains intact on wetting. Groundmass is pale brown-pink with greenish patches. Contains crystal clasts of sanidine, 0.1-1mm long, 5-10% and of biotite, 5-7%, the same crystal clasts as present in above samples. | Traces of goethite irregularly disseminated throughout rock.   |
| 180   | Pale green with pink and white patches    | Tuff: similar to above sample.   | Traces of goethite. Mn oxides (black) intergrown with green clays disseminated throughout rock; Mn oxides = 2%               |
| 190   |   |  |  |
| 190   | Pale green                                | Tuff: same rock as above. See thin-section description.  | Tiny, Mn-oxide dendrites disseminated throughout rock; Mn-oxides also coat fractures. Traces of disseminated goethite.       |
| 200   |   |  |  |
| 200   | Nearly white to pale green                | Tuff: same rock as above.  | Mn-oxides disseminated throughout rock. A dark green mica/clay occurs in veins, sometimes it is associated with MnOX.        |
| 210   |   |  |  |
| 210   | Very pale brown-orange with green patches | Tuff: same rock as above.  | Same as above.   |
| 220   |   |  |  |
| 220   | Very pale orange-pink                     | Tuff: same rock as above. Contains trace amounts of disseminated magnetite.  | Contains only traces of Mn oxides.   |
| 230   |   |  |  |
| 230   | Very pale pink and white                  | Tuff: Very similar to previous sample. Contains crystal clasts of sanidine and biotite. Traces of disseminated magnetite present.  | Contains traces of disseminated Mn oxides and of calcite. Hematite (red) stains fractures.                                   |
| 240   |   |  |  |
| 240   | Pale grey to white                        | Tuff: same as above. Disseminated magnetite present in trace amounts.  | Traces of disseminated goethite/hematite.  |
| 250   |   |  |  |
| 250   | Pale grey to white                        | Tuff: same as above.   | Traces of disseminated goethite and calcite present.   |
| 260   |   |  |  |
| 260   | Light green                               | Tuff: very similar to above sample. Green color due to disseminated celadonite(?).   | Traces of disseminated Mn oxides, celadonite(?)  |
| 270   |   |  |  |
| 270   | Light green                               | Tuff: same as above.   | Same as above.   |
| 280   |   |  |  |
| 280   | Light green with patches of dark green    | Tuff: same as above.   | Celadonite (= bright green mica/clay) occurs both as disseminations and in veinlets, less than 0.1mm wide.                   |
| 290   |   |  |  |
| 290   | Pale green                                | Tuff: same as above. See thin-section description.   | Contains calcite both as disseminations and in rare veins (less than 0.1mm wide). Green color due to presence of celadonite. |
| 300   |   |  |  |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 3 of 10  
 DATE: September 1980  
 DEPTH: 300' TO 450'  
 EXAMINED BY: M. J. Sweeney



| DEPTH | COLOR                | SAMPLE DESCRIPTION   |  |
|-------|----------------------|--|--|
|       |                      | PRIMARY  | SECONDARY  |
| feet  |                      |  |  |
| 300   | Pale grey to white   | Tuff: contains crystal clasts of sanidine and biotite. Also disseminated, very fine-grained magnetite. Matrix is so clay-rich, it disappears during sample washing.  | Clay replacement of vitric matrix is intense.  |
| 310   |                      |  |  |
| 310   | Light grey to white. | Tuff: very similar to above sample, but not as strongly clay-altered. Does not disaggregate so completely during washing.  | Clay alteration of vitric groundmass of tuff.  |
| 320   | Dark grey.           | Dark grey rhyolite porphyry chips comprise 30% of this sample; rhyolite = dike?, flow?, xenoliths?, welded interval?   |  |
| 320   | Light grey to white. | Tuff: same as above.   | Same as above;   |
| 330   | Dark grey.           | Dark grey rhyolite porphyry forms 10-15% of this sample.   | also traces of disseminated celadonite.  |
| 330   | Very pale brown.     | Tuff: same rock as above.  | Traces of disseminated calcite.  |
| 340   |                      | Only a few chips, less than 1% of sample, are from the dark grey rhyolite porphyry present in previous two samples.  |  |
| 340   | Grey-green           | Tuff: Very similar to above samples. Contains crystal clasts of sanidine and biotite. Trace amounts of disseminated magnetite occur. The groundmass is colored like epidote or nontronite.   | Clay replaced groundmass; epidote present??  |
| 350   |                      |  | Traces of disseminated Mn oxides.  |
| 350   | Grey-green           | Tuff: same as above.   | Same as above.   |
| 360   |                      |  |  |
| 360   | Grey                 | Tuff: contains crystal clasts of sanidine and biotite. Trace amounts of disseminated magnetite present. Pumaceous lithic fragments form 5-10% of rock.   | Mn-oxides occur rarely. A few, loose pyrite crystals present in this sample.               |
| 370   |                      |  |  |
| 370   | Grey                 | Tuff: same as above.   | Minor amounts of a pale lime-green clay replaces matrix; occurs in irregular patches.      |
| 380   |                      |  |  |
| 380   | Grey to grey-green   | Tuff: same rock as above. Groundmass is more clay-altered than previous samples; rock disaggregates a lot during washing. Tuff contain pumice fragments and lithic fragments of andesite which form less than 10% of rock. Andesites more Mg-rich than tuff. | Pale lime-green clay more abundant than in previous sample.                                |
| 390   |                      |  |  |
| 390   | Grey to grey-green   | Tuff: Same as above. Groundmass disaggregates on wetting.  | Same as above.   |
| 400   |                      |  |  |
| 400   | Grey to grey-green   | Tuff: same as above. Disaggregates on wetting.   | Same as above.   |
| 410   |                      |  |  |
| 410   | Grey to grey-green.  | Tuff: same rock as above. This sample remained intact during washing. Lithic fragments of andesites, etc., are common; they are more magnetic than tuff.   | Patches of translucent lime-green clay.  |
| 420   |                      |  |  |
| 420   | Light lime-green     | Tuff: same rock as above. Rock disaggregates when wet.   | Groundmass strongly replaced by translucent lime-green clay.                               |
| 430   |                      |  |  |
| 430   | Grey and earthy red  | Tuff: same rock as above; disaggregates on wetting.  | Groundmass strongly replaced by grey clay. Irregular stains of hematite throughout rock.   |
| 440   |                      |  |  |
| 440   | White and pink.      | Tuff: same as above. Contains crystal clasts of sanidine and biotite. Lithic fragments of andesite form 5% of sample. Rock disaggregates on wetting.   | Specks of earthy hematite disseminated throughout rock. Groundmass strongly clay-replaced. |
| 450   |                      |  |  |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 4 of 10  
 DATE: September 1980  
 DEPTH: 450' TO 600'  
 EXAMINED BY: M. J. Sweeney



| DEPTH | COLOR                                 | SAMPLE DESCRIPTION  |   |
|-------|---------------------------------------|---|---|
|       |                                       | PRIMARY   | SECONDARY   |
| feet  |                                       |   |   |
| 450   | Deep maroon red                       | Tuff: contains crystal clasts of sanidine and biotite in abundances similar to previously described samples. Tuff also contains andesite lithic fragments.  | Groundmass (no vitric shards) has been clay-altered and flooded with earthy hematite.   |
| 460   | White and pale earthy red             | Tuff: similar to above sample.  | Weak hematite staining. Weak to moderate clay-alteration of groundmass.   |
| 470   | Light grey-green                      | Tuff: same as above. Lithic fragments form 3-5% of rock.  | Weak hematite staining on fractures. Groundmass replaced by green-grey clays; tiny spots of celadonite (bright green) also present. |
| 480   | Grey-green                            | Tuff: similar to above sample.  | Clay-alteration of groundmass.  |
| 490   | Light Kelly green                     | Tuff: Contains crystal clasts of sanidine and biotite. Groundmass disaggregates when rock is wetted. See thin-section description.  | Clay-alteration of vitric components; some celadonite (?) present (* bright green clay/mica).                                       |
| 500   | Light Kelly green                     | Tuff: same as above.  | Same as above.  |
| 510   | Light Kelly green                     | Tuff: same as above.  | Same as above. Traces of disseminated goethite present.   |
| 520   | Dark grey with patches of Kelly green | Tuff: grey portions of the tuff are quite hard -- these portions were probably well welded and are not now clay-altered. Rock contains crystal clasts of sanidine and of biotite. Trace amounts of magnetite occur as disseminations. | Celadonite/clay replaces groundmass. Pyrite (fresh) coats fractures - only trace amounts seen.                                      |
| 530   | Dark grey with patches of Kelly green | Tuff: same as above. These rock chips are hard; grey-colored portions exhibit a vitreous luster.  | Celadonite/clay alteration of groundmass. Traces of calcite present.  |
| 540   | Dark grey with patches of light green | Tuff: same as above. Hard.  | Celadonite/clay replacement of groundmass. Trace amounts of disseminated goethite (after pyrite).                                   |
| 550   | Green-grey                            | Tuff: same as above. Hard.  | Same as above.  |
| 560   | Green-grey                            | Tuff: same as above. Hard.  | Same as above.  |
| 570   | Green-grey                            | Tuff: same as above. Hard.  | Same as above.  |
| 580   | Green-grey                            | Tuff: same as above. Hard.  | Same as above.  |
| 590   | Greenish grey-brown                   | Tuff: same as above. Hard. See thin-section description and X-ray diffraction pattern - main clay peak at 12.6 Å.   | Clay replacement of vitric components averages ~20%. Trace of pyrite in a vein.   |
| 600   |                                       |   |   |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 5 of 10  
 DATE: September 1980  
 DEPTH: 600' TO 750'  
 EXAMINED BY: M. J. Sweeney



| DEPTH | COLOR                             | SAMPLE DESCRIPTION  |  |
|-------|-----------------------------------|---|--|
|       |                                   | PRIMARY   | SECONDARY  |
| feet  |                                   |   |  |
| 600   | Brownish grey-green               | Tuff: contains crystal clasts of sanidine, 3-5% of rock; biotites are very rare. Rock is hard; it was probably well welded and is now largely devitrified. Very similar to previously described sample. | Celadonite/clay alteration of groundmass. Traces of goethite and calcite.                          |
| 610   | Green-grey                        | Tuff: same as above. Hard.  | Same as above.   |
| 620   | Grey-brown                        | Tuff: similar to above sample. Hard.  | Contains very little celadonite. Traces of calcite, goethite present.                              |
| 630   | Dark grey with green patches      | Tuff: similar to above samples. Hard.   | Patchy celadonite. Thin veins, 0.1-0.5mm wide, filled with pale green and white minerals.          |
| 640   | White with grey and green patches | Tuff: contains crystal clasts of sanidine in a white, gritty groundmass with patches of grey and green.   | Patches of celadonite. Groundmass (a vitric component) partially replaced by clay.                 |
| 650   | Pale lime-green                   | Tuff: very clay-rich -- groundmass disaggregates on wetting. Crystal clasts same as in previous samples.  | Groundmass very clay-altered.  |
| 660   | pale red                          |   |  |
| 660   | Pale lime-green                   | Tuff: same as above. Contains crystal clasts of sanidine and biotite. Groundmass disaggregates readily on wetting.  | Same as above  |
| 670   | Pale lime-green                   | Tuff: same as above.  | Same as above.   |
| 680   | White with pale green patches     | Tuff: same as above.  | Same as above.   |
| 690   | Greyish white                     | Tuff: contains crystal clasts of sanidine and biotite in a greyish white, gritty groundmass. See thin-section description.  | Weakly to moderately clay-altered. Minor carbonate present.  |
| 700   | Greyish white                     | Tuff: same rock as above. Groundmass disaggregates on wetting.  | Same as above.   |
| 710   | Greyish white                     | Tuff: same as above.  | Same as above.   |
| 720   | Light grey-green, black           | Tuff: similar to above sample. Clay-altered vitric groundmass disaggregates on wetting. Andesite/basalt lithic fragments form ~30% of this tuff sample.   | Same as above.   |
| 730   | Light grey with green patches     | Tuff: similar to above samples. Contains irregular patches of celadonite (bright green clay/mica) alteration.   | Same as above. Also contains patches of celadonite. Moderate calcite alteration.                   |
| 740   | Light grey                        | Tuff: similar to above sample. Clay-altered vitric groundmass disaggregates on wetting. Lithic fragments comprise 10-15% of rock.   | Weak to moderate clay alteration including patchy celadonite. Calcite present in moderate amounts. |
| 750   |                                   |   |  |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co.,  
Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

**GEOLOGICAL REPORT**  
**WELL CUTTINGS SAMPLE DESCRIPTION**

PAGE: 6 of 10  
 DATE: September 1980  
 DEPTH: 750' TO 900'  
 EXAMINED BY: M. J. Sweeney



| DEPTH | COLOR                       | SAMPLE DESCRIPTION  |  |
|-------|-----------------------------|---|--|
|       |                             | PRIMARY   | SECONDARY  |
| feet  |                             |   |  |
| 750   | Light grey                  | Tuff: same as previous samples. Clay-altered matrix disaggregates on wetting.   | Moderate clay-alteration. Calcite moderately abundant.   |
| 760   |                             |   |  |
| 760   | Light grey                  | Tuff: similar to previous samples. Contains crystal clasts of feldspar and biotite. Lithic clasts of latite/andesite form 10 to 15% of tuff. Matrix disaggregates on wetting.   | Moderate clay alteration of vitric groundmass. Calcite moderately abundant. Hematite/goethite present in 1mm wide veins (after pyrite?)              |
| 770   |                             |   |  |
| 770   | Pale green and pale red     | Tuff: distinctly different from above sample. Does not disaggregate when wet. Contains crystal clasts of sanidine and biotite in a green- or a red-colored groundmass. Flattened pumice fragments form about 5% of rock. Lithic fragments form 5% of rock (= andesites, etc.).  | Green matrix contains minor, disseminated celadonite. Red matrix is weakly hematite stained. Traces of calcite present.                              |
| 780   | Pale green and red          | Tuff: same as above. Green and red patches may occur both in the same chip.   | Same as above.   |
| 790   |                             |   |  |
| 790   | Pale green and purple       | Tuff: about 30% of these chips are very similar to above sample. Remainder are mottled in color; mottling is on a small scale. These chips are green, creamy white or mottled purplish and creamy white -- this white has the appearance of opal or chert. Many chips exhibit features of small-scale brecciation -- result of faulting? SEE THIN-SECTION DESCRIPTION | Celadonite on fractures. Weak hematite staining present; calcite moderately abundant. Chert-celadonite-calcite fills spaces among brucite fragments. |
| 800   |                             |   |  |
| 800   | Light buff, green, red      | Tuffs: A pale buff, gritty, vitric-crystal tuff forms ~50% of this sample. Green and red vitric-lithic tuffs each form ~25% of sample.  | Clay alteration is moderately well developed. Calcite is abundant. Traces of goethite (after pyrite/magnetite?) occur.                               |
| 810   |                             |   |  |
| 810   | Grey-green, red             | Tuff: contains crystal clasts of feldspar and biotite in a grey-green (clay-rich) or red (hematite-stained) groundmass.   | Moderate clay-alteration. Calcite fairly abundant.   |
| 820   |                             |   |  |
| 820   | Pale buff with grey streaks | Tuff: contains crystal clasts of feldspars and biotite in a pale buff - almost white - groundmass with vitreous grey streaks (bands of devitrification?).   | Same as above.   |
| 830   |                             |   |  |
| 830   | Pale buff with grey streaks | Tuff: same as above. Examination in thin-section shows that the groundmass has been totally devitrified to fine-grained, anhedral quartz and Kspar.   | Weak to moderate clay-alteration of groundmass. Calcite disseminated and in veins.   |
| 840   |                             |   |  |
| 840   | Grey-white                  | Tuff: dominantly vitric -- contains few feldspar and biotite phenocrysts.   | Same as above.   |
| 850   |                             |   |  |
| 850   | Grey-white                  | Tuff: same as above.  | Same as above.   |
| 860   |                             |   |  |
| 860   | Grey-white                  | Tuff: same as above. Examination in thin-section reveals that this is a vitric-crystal tuff. Glass has been completely replaced by fine-grained, anhedral Kspar and quartz. Crystal clasts of sanidine, plagioclase and biotite are present.  | Same as above. Calcite occurs in veins and locally floods groundmass. Microcrystalline quartz occurs in veins.                                       |
| 870   |                             |   |  |
| 870   | Grey-white                  | Tuff: same as above.  | Same as above.   |
| 880   |                             |   |  |
| 880   | Grey-white                  | Tuff: similar to above sample. Ratio of feldspar clasts to groundmass is higher; groundmass may be relatively more prone to disaggregation on wetting.  | Same as above.   |
| 890   |                             |   |  |
| 890   | Grey-white                  | Tuff: contains quartz "eyes", otherwise similar to previous samples. SEE THIN-SECTION DESCRIPTION   | Clay-alteration of groundmass. Vein and disseminated calcite. Microcrystalline quartz occurs in fractures with calcite.                              |
| 900   |                             |   |  |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co.,  
Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 7 of 10  
 DATE: September 1980  
 DEPTH: 900' TO 1050'  
 EXAMINED BY: M. J. Sweeney



| DEPTH | COLOR                             | SAMPLE DESCRIPTION   |   |
|-------|-----------------------------------|--|---|
|       |                                   | PRIMARY  | SECONDARY   |
| feet  |                                   |  |   |
| 900   | Grey white                        | Tuff: same as previous sample. Groundmass (clay-altered vitric shards and dust) disappears during washing.   | Same as above.  |
| 910   |                                   |  |   |
| 910   | Mottled dark grey and grey-white  | Cherty Dolomite: fine-grained; even textured.  | Pyrite fills fractures; py forms ~0.5% of rock.   |
| 920   |                                   |  |   |
| 920   | Same as above                     | Cherty Dolomite  | Same as above.  |
| 930   |                                   |  |   |
| 930   | Dark grey, white                  | Dolomite is dark grey. White rock is a vitric-crystal tuff; it contains crystal clasts of sanidine and biotite in a white clay-rich matrix which disaggregates on wetting. X-ray diffraction analysis indicates clay is montmorillonite. | Pyrite occurs in fractures in dolomite. clay developed in tuff.                               |
| 940   |                                   |  |   |
| 940   | Same as above                     | Dolomite and Tuff: same as above.  | Same as above.  |
| 950   |                                   |  |   |
| 950   | Same as above                     | Dolomite and Tuff: same as above.  | Same as above.  |
| 960   |                                   |  |   |
| 960   | Same as above                     | Dolomite and Tuff: same as above.  | Same as above.  |
| 970   |                                   |  |   |
| 970   | Mottled grey and white            | Dolomite: fine-grained; cherty.  | Same as above.  |
| 980   |                                   | About 10 to 20% of the chips in this sample are from same white, clay-rich tuff described above.   |   |
| 980   | Same as above.                    | Dolomite: same as above.   | Traces of pyrite on fracture surfaces.  |
| 990   |                                   |  |   |
| 990   | Same as above                     | Dolomite: same as above.<br>SEE THIN-SECTION DESCRIPTION   | Same as above.  |
| 1000  |                                   |  |   |
| 1000  | Mottled dark grey and white       | Dolomite: same as above.   | Pyrite on fractures. forms 0.1% of rock.  |
| 1010  |                                   |  |   |
| 1010  | Same as above                     | Dolomite: same as above.   | Same as above.  |
| 1020  |                                   |  |   |
| 1020  | Same as above                     | Dolomite: same as above.   | Same as above.  |
| 1030  |                                   |  |   |
| 1030  | Mottled dark grey and white, buff | Dolomites are mottled grey and white; these chips form 40-50% of sample.<br>Fine-grained sandstones are buff; these chips form 40-50% of sample. A few chips have slickensided surfaces.   | Pyrite occurs on fractures in dolomite. Traces of goethite occurs on fractures in sandstones. |
| 1040  |                                   |  |   |
| 1040  | Same as above                     | Dolomites and sandstones: same as above.   | Same as above.  |
| 1050  |                                   |  |   |

WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co.,  
Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

**GEOLOGICAL REPORT**  
**WELL CUTTINGS SAMPLE DESCRIPTION**



PAGE: 8 of 10  
 DATE: September 1980  
 DEPTH: 1050' TO 1200'  
 EXAMINED BY: M.J. Sweeney

| DEPTH | COLOR                       | SAMPLE DESCRIPTION   |  |
|-------|-----------------------------|--|--|
|       |                             | PRIMARY  | SECONDARY  |
| feet  |                             |  |  |
| 1050  | Light grey                  | Limestone: fine-grained.   | Thin (less than 1mm wide) calcite veins occur.   |
| 1060  |                             |  |  |
| 1060  | Light grey                  | Limestone: fine-grained.   | Thin calcite veins. Traces of pyrite in fractures.   |
| 1070  |                             |  |  |
| 1070  | Mottled dark and light grey | Limestone: fine-grained.   | Same as above.   |
| 1080  |                             |  |  |
| 1080  | Same as above               | Limestone: same as above.  | Same as above.   |
| 1090  |                             |  |  |
| 1090  | Same as above               | Limestone: sandy and cherty; fine-grained. SEE THIN-SECTION DESCRIPTION  | Disseminated and vein pyrite. Thin calcite veins.  |
| 1100  |                             |  |  |
| 1100  | Light grey                  | Sandy Limestone: rock contains a small proportion of fine-sand. A few chips of slickensided fault gouge occur in this sample.                              | Traces of vein and disseminated pyrite, both fresh and oxidized. Some pyrite is in quartz veins.                           |
| 1110  |                             |  |  |
| 1110  | Light grey-buff             | Calcareous Sandstone: composed of fine sand and calcite. Yellow carbonate (Fe-rich?) coats some fracture surfaces - it also disseminated in parts of rock. | Traces of pyrite. Slickensided fault gouge chips present. A few, bleached, hematite-stained and veined siltstones present. |
| 1120  |                             |  |  |
| 1120  | Light grey and buff         | Calcareous Sandstone: same as above.   | Pyrite (mostly oxidized) occurs in trace amounts as disseminations (0.1%). Fault gouge chips present.                      |
| 1130  |                             |  |  |
| 1130  | Light grey                  | Calcareous Sandstone: same as above.   | Traces of fresh and oxidized pyrite, mostly in thin veins, less than 1mm wide. Less pyrite than in previous sample.        |
| 1140  |                             |  |  |
| 1140  | Light grey                  | Calcareous Sandstone: same as above.   | Traces of oxidized pyrite - less than in above sample.   |
| 1150  |                             |  |  |
| 1150  | Light brown-orange          | Calcareous Siltstone: finer-grained than above sample. Thinly laminated.   | Contains traces of oxidized and fresh disseminated pyrite.   |
| 1160  |                             |  |  |
| 1160  | Light brown-orange          | Calcareous Siltstone: same as above.   | Contains less pyrite than above sample. Fault gouge chips present.   |
| 1170  |                             |  |  |
| 1170  | Light brown-orange          | Calcareous Siltstone: same as above.   | Same as above.   |
| 1180  |                             |  |  |
| 1180  | Light brown                 | Calcareous Sandstone: coarser-grained than previous sample.  | Same as above.   |
| 1190  |                             |  |  |
| 1190  | Light orange-brown          | Calcareous Sandstone: similar to previous sample. SEE THIN-SECTION DESCRIPTION   | Traces of oxidized, disseminated magnetite. Thin calcite and calcite/gts veins.  |
| 1200  |                             |  | Fault-gouge chips present.   |

WELL: SR-4  
 LOCATION: Diriz Valley, Churchill Co.,  
Nevada.  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 9 of 10  
 DATE: September 1980  
 DEPTH: 1200' TO 1350'  
 EXAMINED BY: M.J. Sweeney



| DEPTH | COLOR                      | SAMPLE DESCRIPTION   |   |
|-------|----------------------------|--|---|
|       |                            | PRIMARY  | SECONDARY   |
| feet  |                            |  |   |
| 1200  | Pale orange, pale red      | Calcareous Siltstones: fine-grained; well cemented. Reacts vigorously to cold, dilute HCL. | Traces of oxidized mgt and/or pyrite -- gone to hematite/goethite. Some fresh pyrite as loose crystals. |
| 1210  |                            |  |   |
| 1210  | Pale brown-orange          | Calcareous Siltstones: same as above.  | Traces of oxidized mgt/py. Chips of limonite-stained, slickensided fault gouge.                         |
| 1220  |                            |  |   |
| 1220  | Medium grey                | Limestone: fine-grained; not laminated.  | Thin (1mm wide or less) calcite veins. Traces of fresh pyrite on fractures.                             |
| 1230  |                            |  |   |
| 1230  | Medium to dark grey        | Limestone: same as above.  | Same above. Py also seen in wider (1-5mm) calcite veins.  |
| 1240  |                            |  |   |
| 1240  | Same as above              | Limestone: same as above.  | Same as above.  |
| 1250  |                            |  |   |
| 1250  | Same as above              | Limestone: same as above.  | Same as above.  |
| 1260  |                            |  |   |
| 1260  | Medium grey                | Limestone: same as above.  | Thin calcite veins. Traces of disseminated pyrite (oxidized). Slickensided chips present.               |
| 1270  |                            |  |   |
| 1270  | Medium to dark grey        | Limestone: same as above.  | Thin calcite veins occur; a few contain pyrite. Pyrite also coats fractures.                            |
| 1280  |                            |  |   |
| 1280  | Medium grey and grey-brown | Limestones: same as above.   | Same as above. Pyrite often oxidized. Slickensided chips present.                                       |
| 1290  |                            |  |   |
| 1290  | Medium grey and grey-brown | Limestone: same as above. SEE THIN-SECTION DESCRIPTION                                     | Pyrite (partially oxidized to hematite) occurs in veins and as disseminations.                          |
| 1300  |                            |  |   |
| 1300  | Medium grey                | Limestone: same as above.  | Thin calcite veins present; some contain goethite/hematite (after pyrite?). Slickensided chips present. |
| 1310  |                            |  |   |
| 1310  | Medium grey.               | Limestone: same as above.  | Thin calcite veins present. Fresh pyrite seen on a few fracture surfaces. Slickensided chips present.   |
| 1320  |                            |  |   |
| 1320  | Medium grey                | Limestone: same as above.  | Thin calcite veins occur. Traces of disseminated goethite.  |
| 1330  |                            |  |   |
| 1330  | Medium grey                | Limestone: same as above.  | Thin calcite veins occur, some of which contain pyrite.   |
| 1340  |                            |  |   |
| 1340  | Medium grey                | Limestone: same as above.  | Same as above.  |
| 1350  |                            |  |   |



WELL: SR-4  
 LOCATION: Dixie Valley, Churchill Co.,  
Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

**GEOLOGICAL REPORT**  
 WELL CUTTINGS SAMPLE DESCRIPTION




PAGE: 10 of 10  
 DATE: September 1980  
 DEPTH: 1350' TO 1500'  
 EXAMINED BY: M.J. Sweeney

| DEPTH | COLOR                    | SAMPLE DESCRIPTION   |   |
|-------|--------------------------|--|---|
|       |                          | PRIMARY  | SECONDARY   |
| feet  |                          |  |   |
| 1350  | Medium grey              | Limestone: very fine-grained; even textured.   | Thin calcite veins present, some of which contain fresh pyrite. Slickensided chips occur.                       |
| 1360  |                          |  |   |
| 1360  | Medium grey              | Limestone: same as above.  | Same as above.  |
| 1370  |                          |  |   |
| 1370  | Light grey               | Limestone: same as above.  | Rare calcite veins, some of which contain hematite (after pyrite?). Pyrite in thin discontinuous seams (no. 1%) |
| 1380  |                          |  |   |
| 1380  | Light grey               | Limestone: same as above.  | Thin calcite veins present; some contain pyrite (no. 2%). Slickensided chips present.                           |
| 1390  |                          |  |   |
| 1390  | Grey                     | Limestone: same as above.<br>SEE THIN-SECTION DESCRIPTION  | Calcite veins contain pyrite some of which is oxidized to hematite.   |
| 1400  |                          |  |   |
| 1400  | Grey                     | Limestone: same as above.  | Same as above.  |
| 1410  |                          |  |   |
| 1410  | Grey                     | Limestone: same as above.  | Same as above.  |
| 1420  |                          |  |   |
| 1420  | Brown-grey and grey      | Limestone: same as above.  | Traces of fresh pyrite. Slickensided chips present.   |
| 1430  |                          |  |   |
| 1430  | Same as above.           | Limestone: same as above.  | Rare, thin, calcite veins.  |
| 1440  |                          |  |   |
| 1440  | Grey                     | Limestone: same as above.  | Same as above.  |
| 1450  |                          |  |   |
| 1450  | Grey                     | Limestone: same as above.  | Same as above. Slickensided chips present.  |
| 1460  |                          |  |   |
| 1460  | Grey                     | Limestone: same as above.  | Same as above.  |
| 1470  |                          |  |   |
| 1470  | Grey, red                | Limestone: same as above.  | Rare, thin, calcite veins present, some of which contain fresh and oxidized pyrite.                             |
| 1480  |                          | Fault gouge (?): hematite-red, clayey material which is often slickensided forms 1-3% of sample. |   |
| 1480  | Grey, red.               | Limestone: same as above.  | Rare, thin, calcite veins occur.  |
| 1490  |                          | Fault gouge (!): red; forms 3-5% of sample.  |   |
| 1490  | Grey and light grey, red | Limestone: same as above.<br>SEE THIN-SECTION DESCRIPTIONS                                       | Pyrite (partially oxidized) in thin veins, less than 1mm wide. Thin calcite veins also present.                 |
| 1500  |                          | Fault gouge (?): red; forms 2-5% of sample.  |   |

# SHALLOW TEMPERATURE GRADIENT HOLE RECORD



|                   |                                   |                                    |                           |
|-------------------|-----------------------------------|------------------------------------|---------------------------|
| <u>SR-3</u>       | <u>Dixie Valley</u>               | <b>Southland Royalty Company</b>   |                           |
| GRADIENT HOLE NO. | PROSPECT                          |                                    |                           |
| <u>NW NE</u>      | <u>S 32 T 24N R 37E</u>           | <u>Churchill</u>                   | <u>Nevada</u>             |
| LOCATION          | SEC. TOWNSHIP RANGE               | COUNTY                             | STATE                     |
| <u>1/27/80</u>    | <u>2/14/80</u>                    | <u>1472</u>                        | <u>4.83</u>               |
| SPUD DATE         | COMPLETION DATE                   | BOTTOM HOLE TEMP.                  | AT _____ FT. _____ F/100' |
| <u>1,500'</u>     | <u>R. Jodry</u>                   | GRADIENT <u>570</u> TO <u>1470</u> |                           |
| TOTAL DEPTH       | GEOLOGIST OR PERSON READING TEMP. | <u>Christiansen Bros.</u>          |                           |
|                   |                                   | COMPANY DRILLING HOLES             |                           |



| FOR HOLES 0-500'      |        |       |         |    |
|-----------------------|--------|-------|---------|----|
| TEMP                  | °C     | °F    | °F      | °F |
| DATE OF SURVEY        | 3-5-80 | 35-80 | 5-29-80 |    |
| TIME SINCE COMPLETION | 20     | 20    | 105     |    |
| 0                     | 10.4   | 50.7  | 75.2    |    |
| 100                   | 48.0   | 118.4 | 121.5   |    |
| 200                   | 55.2   | 131.3 | 134.6   |    |
| 300                   | 59.2   | 139.1 | 142.2   |    |
| 400                   | 62.8   | 145.0 | 147.4   |    |
| 500                   | 65.5   | 149.9 | 152.4   |    |
| 600                   | 68.7   | 155.6 | 157.1   |    |
| 700                   | 71.6   | 160.8 | 162.1   |    |
| 800                   | 74.8   | 166.6 | 167.4   |    |
| 900                   | 77.8   | 172.0 | 172.0   |    |
| 1000                  | 80.6   | 177.0 | 176.5   |    |
| 1100                  | 83.4   | 182.1 | 181.0   |    |
| 1200                  | 85.8   | 186.4 | 185.4   |    |
| 1300                  | 88.2   | 190.7 | 189.5   |    |
| 1400                  | 90.6   | 195.0 | 193.6   |    |
| 1474                  | 92.4   | 198.3 | 196.2   |    |

LOST CIRCULATION ZONES:

---

WATER ENTRIES: 150'-160'

Static Water Level-140'

USE OTHER SIDE FOR GRADIENT HOLES OVER 500'  
DISCUSS DRILLING PROBLEMS ON ANOTHER SHEET

SR-3

GRADIENT HOLE NO.

3/5/80

DATE OF SURVEY

| DEPTH | °C   | °F     | DEPTH | °C   | °F     | DEPTH | °C   | °F     | DEPTH | °C   | °F     |
|-------|------|--------|-------|------|--------|-------|------|--------|-------|------|--------|
| 0     | 10.4 | 50.72  | 460   | 64.7 | 148.46 | 920   | 78.4 | 173.12 | 1380  | 90.2 | 194.36 |
| 10    | 19.6 | 67.28  | 470   | 65.0 | 149.0  | 930   | 78.7 | 173.66 | 1390  | 90.4 | 194.72 |
| 20    | 29.1 | 84.38  | 480   | 64.8 | 148.64 | 940   | 79.0 | 174.2  | 1400  | 90.6 | 195.08 |
| 30    | 33.5 | 92.3   | 490   | 65.2 | 149.36 | 950   | 79.2 | 174.56 | 1410  | 90.8 | 195.44 |
| 40    | 36.8 | 98.24  | 500   | 65.5 | 149.9  | 960   | 79.5 | 175.1  | 1420  | 91.1 | 195.98 |
| 50    | 39.6 | 103.28 | 510   | 66.0 | 150.8  | 970   | 79.8 | 175.64 | 1430  | 91.3 | 196.34 |
| 60    | 43.6 | 110.48 | 520   | 66.4 | 151.52 | 980   | 80.0 | 176.0  | 1440  | 91.6 | 196.7  |
| 70    | 44.7 | 112.46 | 530   | 66.7 | 152.06 | 990   | 80.3 | 176.54 | 1450  | 91.9 | 197.06 |
| 80    | 45.8 | 114.44 | 540   | 67.1 | 152.78 | 1000  | 80.6 | 177.08 | 1460  | 92.1 | 197.42 |
| 90    | 46.9 | 116.42 | 550   | 67.4 | 153.32 | 1010  | 80.9 | 177.62 | 1470  | 92.3 | 197.78 |
| 100   | 48.0 | 118.4  | 560   | 67.7 | 153.86 | 1020  | 81.2 | 178.16 | 1472  | 92.4 | 198.32 |
| 110   | 48.9 | 120.02 | 570   | 67.9 | 154.22 | 1030  | 81.5 | 178.7  |       |      |        |
| 120   | 49.8 | 121.64 | 580   | 68.2 | 154.76 | 1040  | 81.8 | 179.24 |       |      |        |
| 130   | 50.6 | 123.08 | 590   | 68.5 | 155.3  | 1050  | 82.0 | 179.6  |       |      |        |
| 140   | 51.3 | 124.34 | 600   | 68.7 | 155.66 | 1060  | 82.3 | 180.14 |       |      |        |
| 150   | 52.0 | 125.6  | 610   | 69.0 | 156.2  | 1070  | 82.5 | 180.5  |       |      |        |
| 160   | 52.6 | 126.68 | 620   | 69.2 | 156.56 | 1080  | 82.8 | 181.04 |       |      |        |
| 170   | 53.2 | 127.76 | 630   | 69.5 | 157.1  | 1090  | 83.1 | 181.58 |       |      |        |
| 180   | 53.9 | 129.02 | 640   | 69.8 | 157.64 | 1100  | 83.4 | 182.12 |       |      |        |
| 190   | 54.6 | 130.28 | 650   | 70.1 | 158.18 | 1110  | 83.7 | 182.66 |       |      |        |
| 200   | 55.2 | 131.36 | 660   | 70.4 | 158.72 | 1120  | 83.9 | 183.02 |       |      |        |
| 210   | 55.7 | 132.36 | 670   | 70.7 | 159.26 | 1130  | 84.1 | 183.38 |       |      |        |
| 220   | 56.2 | 133.16 | 680   | 71.0 | 159.8  | 1140  | 84.4 | 183.92 |       |      |        |
| 230   | 56.7 | 134.06 | 690   | 71.3 | 160.34 | 1150  | 84.7 | 184.46 |       |      |        |
| 240   | 57.2 | 134.96 | 700   | 71.6 | 160.88 | 1160  | 84.9 | 184.82 |       |      |        |
| 250   | 57.5 | 135.5  | 710   | 72.1 | 161.78 | 1170  | 85.1 | 185.18 |       |      |        |
| 260   | 58.0 | 136.4  | 720   | 72.5 | 162.5  | 1180  | 85.4 | 185.72 |       |      |        |
| 270   | 58.3 | 136.94 | 730   | 72.8 | 163.04 | 1190  | 85.6 | 186.08 |       |      |        |
| 280   | 58.8 | 137.84 | 740   | 73.0 | 163.4  | 1200  | 85.8 | 186.44 |       |      |        |
| 290   | 59.2 | 138.56 | 750   | 73.3 | 163.94 | 1210  | 86.1 | 186.98 |       |      |        |
| 300   | 59.5 | 139.1  | 760   | 73.7 | 164.66 | 1220  | 86.3 | 187.34 |       |      |        |
| 310   | 59.8 | 139.64 | 770   | 74.0 | 165.2  | 1230  | 86.5 | 187.7  |       |      |        |
| 320   | 60.0 | 140.0  | 780   | 74.3 | 165.74 | 1240  | 86.8 | 188.24 |       |      |        |
| 330   | 60.3 | 140.54 | 790   | 74.6 | 166.28 | 1250  | 87.0 | 188.6  |       |      |        |
| 340   | 60.7 | 141.26 | 800   | 74.8 | 166.64 | 1260  | 87.2 | 188.96 |       |      |        |
| 350   | 61.1 | 141.98 | 810   | 75.2 | 167.36 | 1270  | 87.5 | 189.5  |       |      |        |
| 360   | 61.4 | 142.52 | 820   | 75.4 | 167.72 | 1280  | 87.7 | 189.86 |       |      |        |
| 370   | 61.7 | 143.06 | 830   | 75.7 | 168.26 | 1290  | 88.0 | 190.4  |       |      |        |
| 380   | 62.0 | 143.60 | 840   | 76.0 | 168.8  | 1300  | 88.2 | 190.76 |       |      |        |
| 390   | 62.4 | 144.32 | 850   | 76.3 | 169.34 | 1310  | 88.5 | 191.3  |       |      |        |
| 400   | 62.8 | 145.04 | 860   | 76.6 | 169.88 | 1320  | 88.7 | 191.66 |       |      |        |
| 410   | 63.1 | 145.58 | 870   | 76.9 | 170.42 | 1330  | 88.9 | 192.02 |       |      |        |
| 420   | 63.5 | 146.3  | 880   | 77.2 | 170.96 | 1340  | 89.1 | 192.38 |       |      |        |
| 430   | 63.8 | 146.84 | 890   | 77.5 | 171.5  | 1350  | 89.4 | 192.92 |       |      |        |
| 440   | 64.1 | 147.38 | 900   | 77.8 | 172.04 | 1360  | 89.7 | 193.46 |       |      |        |
| 450   | 64.4 | 147.92 | 910   | 78.1 | 172.58 | 1370  | 90.0 | 194.0  |       |      |        |

# SHALLOW TEMPERATURE GRADIENT HOLE RECORD



SR-3  
GRADIENT HOLE NO.

NW NE  
LOCATION

01/27/80  
SPUD DATE

1500  
TOTAL DEPTH

Dixie Valley  
PROSPECT

S 32 T 25N R 37E  
SEC. TOWNSHIP RANGE

02/14/80  
COMPLETION DATE

Richard Jodry  
GEOLOGIST OR PERSON READING TEMP.

196.2°F  
BOTTOM HOLE TEMP.


Reading for 05/29/80

Southland Royalty Company

Churchill Nevada  
COUNTY STATE

1474 4.38°  
AT FT. F/100

GRADIENT 500 TO 147'  
J. D. Christiansen Drilling  
COMPANY DRILLING HOLES Ely



FOR HOLES 0-500'

| TEMP                  | °C       | °F       | °F | °F |
|-----------------------|----------|----------|----|----|
| DATE OF SURVEY        | 5-29-80  | 5-29-80  |    |    |
| TIME SINCE COMPLETION | 105 DAYS | 105 DAYS |    |    |
| 0                     | 24.0     | 75.2     |    |    |
| 100                   | 49.7     | 123.2    |    |    |
| 200                   | 57.0     | 134.6    |    |    |
| 300                   | 61.2     | 149.6    |    |    |
| 400                   | 64.1     | 147.4    |    |    |
| 500                   | 66.9     | 152.4    |    |    |
| 600                   | 69.5     | 157.1    |    |    |
| 700                   | 72.3     | 162.1    |    |    |
| 800                   | 75.2     | 167.4    |    |    |
| 900                   | 77.8     | 172.0    |    |    |
| 1000                  | 80.3     | 176.5    |    |    |
| 1100                  | 82.8     | 181.0    |    |    |
| 1200                  | 85.2     | 185.4    |    |    |
| 1300                  | 87.5     | 189.4    |    |    |
| 1400                  | 89.8     | 193.6    |    |    |
| 1474                  | 91.2     | 196.2    |    |    |

LOST CIRCULATION ZONES:

WATER ENTRIES: 150'-160'  
Gravel bed.

Prep. by DCM





R - 37 - E

T  
25  
N

T  
25  
N

R - 37 - E



**Southland Royalty Company**  
*NATURAL RESOURCES DISTRICT*  
 FORT WORTH, TEXAS

FIGURE 1  
**DIXIE VALLEY, NEVADA**  
 LOCATION AND INDEX MAP  
**SRC-3**

Scale: 1" 2000'



SUMMARY OF LITHOLOGY AND ALTERATION IN SR-3,  
DIXIE VALLEY, CHURCHILL COUNTY, NEVADA

M. J. Sweeney, July 1980

INTRODUCTION

Samples of washed, rotary cuttings taken every ten feet from drill-hole SR-3 were examined under a stereomicroscope. A thin-section of the cuttings was described from every 100 feet throughout the 1500 foot length of SR-3. Detailed descriptions of the cuttings and thin-sections are attached to this report. Photomicrographs of the thin-sections are also attached.

LITHOLOGIES

SR-3 is in gravels for its entire 1500 foot depth. All the gravel was derived from a common source. It is of nearly uniform composition for 1500 feet, i.e., the same types of clasts occur in about the same proportions throughout the entire hole. Metasandstones and phyllites are the most common lithic clasts. Locally the gravels are cemented by a mixture of clay<sup>±</sup>calcite containing sand-sized lithic and crystal clasts; this clay-sand material may be derived from volcanic ash.

The rock types present in the gravel are listed below.

- 1) Limestone: Black, foliated, very fine-grained, carbonaceous, locally phlogopitic limestone.
- 2) Phyllites: Black, foliated, carbonaceous phyllites composed of varying proportions of biotite, sericite and silt-sized quartz and feldspar. Occasionally porphyroblasts of biotite or of clinozoisite occur in some of the phyllite particles. Often disseminated anhedral magnetite is present. Pyrite was observed rarely.
- 3) Diorite: Dark grey to dark grey green, fine-grained diorite or diabase. The original mafic minerals in this rock type were pyroxenes; in some particles, unaltered pyroxene remains, but most often it has been replaced by chlorite and/or montmorillonite or sericite<sup>±</sup>carbonate. Black Fe-oxides, most probably mixtures of ilmenite/magnetite, are present in the diorite; locally ilmenite/magnetite is oxidized to earthy red hematite. Pyrite was not observed in any diorite fragments.
- 4) Metasandstones: White, buff, pale pink, orange and greenish. These particles are well indurated and never friable. A small proportion of the metasandstones in most every sample contain goethite pseudomorphing a disseminated cubic mineral (pyrite??). Only rarely was unoxidized pyrite observed in metasandstone particles.



It is composed 1) of clay which appears in thin-section to be a montmorillonite, 2) of angular sand grains of quartz, feldspar, mafic minerals, and carbonate, 3) of very fine-grained calcite cement, and 4) of sand-, pebble- and cobble-sized lithic clasts. The lithic clasts include all of the rock types described above in this report.

The clay-sand-lithic material is locally abundant in the hole; its abundance seems to correlate with how clayey the unwashed samples are. The letters, n, s, m and st in the depth column of the stratigraphic log indicate how clayey the unwashed samples are. The symbol "n" indicates that the sample was fairly free of clay, "s" slightly clayey, "m" moderately clayey and "st" strongly clay-cemented. The clay-sand-lithic material is compositionally and behaviorly similar to the bentonite-gel drilling mud used in SR-3; there is no definitive way of deciding which type of material, drilling mud or clay-sand-lithic material, is dominant in the clayey horizons. Some of the clay-rich samples, particularly those at 600', contain pyritic mudstone; the mudstone may have been deposited originally as a lacustrine bed.

The clay-sand material may have originated as volcanic ash. This is suggested by the heterogeneity of the crystal clasts and by the possibility that the clay could be a product of altered glass shards. The clay-sand material and the lithic clasts are thoroughly mixed together; the clasts and ash were probably transported together to the site of SR-3.

Thin (less than 1 mm wide) calcite veins were seen in consolidated fragments of clay-sand-lithic material which survived the washing process. Only once was a pyrite crystal seen in the calcite veins; the pyrite was attached to the surface of the vein. No disseminated pyrite was seen in the clay-sand material.

14) Pyrite crystals<sup>±</sup>clay: Loose pyrite crystals, which occasionally are associated with a white clay (kaolinite?), very fine-grained sericite? and calcite occur in all samples below 590 feet. Frequently these crystals are oxidized. Possibly, these loose crystals were derived from pyrite deposited in open fractures in the consolidated gravels.

#### Alteration

Unraveling the alteration history of gravels, especially those sampled by rotary drilling, is not straightforward. The lithic clasts may contain alteration assemblages developed originally at their sources and not at the present site. Soft or friable assemblages and veins are easily disaggregated by the process of drilling rotary holes and these same assemblages are removed from the sample by washing.

Pyritization. Pyrite occurs 1) as loose particles of subhedral crystal aggregates, sometimes associated with a white clay and calcite, 2) as loose euhedral crystals 1 mm or less across, and 3) as disseminations in siltstone, metasandstones, phyllites and mudstones. Nearly all of the pyrite? in the metasandstones has been replaced by goethite. A smaller proportion of the pyrite in siltstones has been oxidized and even less of that in mudstones has been oxidized. The significance of the disseminated pyrite cannot be evaluated without knowing if pyrite is present at the source area of these clasts.

The loose pyrite crystals are possibly from open veins or from a mudstone disaggregated by drilling/sample washing. The total volume of pyrite in the samples from SR-3 ranges between 0.1 and 0.5 volume % through the hole, including both disseminated and loose pyrite crystals. There is a fair probability that little of this pyrite was deposited as the result of hydrothermal processes occurring at the site of this drill hole, SR-3. The possibility that little of the pyrite was locally deposited is supported by the following observations: 1) only a trace of pyrite was observed in a vein in the clay-sand-lithic matrix material, and 2) no pyrite was seen in the diorite/diabase fragments. Both of these lithologies are usually more susceptible to pyritization than quartz-rich rocks.

Calcite Veining. Thin calcite veins occur in particles of clay-sand-lithic matrix material. Only once was a small pyrite crystal observed on the surface of one of these calcite veins. Very fine-grained calcite was observed occasionally on the surface of lithic clasts; this material is possibly caliche.

Quartz Veining. Drusy quartz veins occur in metasandstones and phyllites. Most probably this veining is related to events at the source area of these particles.

Opal(?) Veining. Opal-quartz occurs in thin veins in clay-sand matrix material. They were observed only in upper 600' of the drill hole.

Summary. Drill hole SR-3 has sampled 1500 feet of gravels locally cemented by sand-clay-carbonate material that is possibly derived from volcanic ash. Many of the clasts in the gravel are weakly pyritized; the pyrite possibly was deposited initially at the source area of these clasts.

The only alteration definitely related to the site of the drill hole is weak calcite as well as opal-quartz veining developed in the sand-clay matrix material.

Oxidation of pyrite is fairly complete down to 1500', the total depth of the hole.

## Clay Analysis

An unwashed sample of strongly clay-cemented drill cuttings was prepared for clay analysis. The sample selected for analysis is from the 1130'-1140' interval. The sample of chips was washed in de-ionized water in a blender. Sodium tripolyphosphate was added to peptize suspended clays. The suspended material was then centrifuged at 1000 r.p.m. to remove larger than clay-sized material. Suspended clay was dropped by centrifuging at 4000 r.p.m. A portion of the dropped clay fraction was smeared on a glass slide and air dried. This sample was analyzed by x-ray diffraction. The clay fraction from the 1130'-1140' interval contains calcite, montmorillonite, illite and kaolinite in order of decreasing abundance.

Clay analysis was also made of a composite of clay-sand fragments which had survived washing. The sample consisted of fragments collected from throughout the hole during chip logging. X-ray diffraction analysis shows that this sample also contains calcite, montmorillonite, illite and kaolinite. Relative peak intensities on this XRD pattern and the previously discussed pattern are very similar. Drilling mud does not appear to have strongly affected peak intensities obtained on the 1130'-1140' sample.

The two patterns are attached to this report.

DESCRIPTIONS OF THIN-SECTIONS FROM SR-3,  
DIXIE VALLEY, CHURCHILL CO., NEVADA

100'-110'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u> : Mostly very fine-grained; grain size 0.1 mm or less, usually 0.01 mm. Rock exhibits fine-scale foliation. Contains porphyroblasts? or detrital flakes of phlogopite; these mica flakes oriented parallel to rock foliation. Very fine-grained carbonaceous? particles (=black dust) occur throughout rock, distributed both inter- and intragranularly. One of the limestone chips contains a vein filled with elongate quartz crystals oriented perpendicularly to vein walls. Ghosts of fossils are still recognizable in most chips. | 7                         | 33                         |
| <u>Phyllite</u> : Very fine-grained. Composed of foliated biotite, muscovite and silt-sized quartz and feldspar grains.  | 2                         | 10                         |
| <u>Diorite/Diabase</u> : One unaltered chip is composed of unoriented plagioclase laths, and lesser pyroxene and possibly olivine. In another chip, mafic minerals have been replaced by chlorite. In another, montmorillonite has replaced mafic minerals, and kaolinite has replaced plagioclase.  | 3                         | 14                         |
| <u>Metasandstone</u> : One grain composed of sand-sized quartz grains and muscovite-chlorite porphyroblasts. One composed of subangular to subrounded quartz grains about 0.1 mm across; sorting good. Silica, clay and calcite occur as cement.   | 2                         | 10                         |
| <u>Marble</u> : Grain size about 1 mm; granoblastic. One chip contains patches of chert. Another contains patches of more coarsely crystalline quartz (recrystallized chert?). The latter also contains patches of montmorillonite after an unidentified mineral (low birefringence=first order grey; low 2V, 10-30°; opt +; good cleavage).   | 3                         | 14                         |
| <u>Chert</u> : Recrystallized; quartz occurs as fine-grained feathery crystals.  | 3                         | 14                         |
| <u>Silicified Cataclasite</u> : Original rock type not known; now composed of subhedral quartz of variable grain size. Patches of brown clay (montmorillonite?) form 5-10% of chip.  | 1                         | 5                          |
| Total Number of Chips Examined   | 21                        | 100%                       |

190'-200'

| Rock Type   | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|---|---------------------------|----------------------------|
| <p><u>Phyllite:</u> This category also includes a few clasts of hornfelsed muddy siltstones; the only significant difference between hornfelses and phyllites is the degree or complete lack of foliation of mica in the phyllites.</p> <p>Very fine-grained, metamorphosed clayey siltstones and silty mudstones; usually composed of subequal amounts of silt-sized quartz and feldspar and of very fine-grained white micas, lesser biotite and of variable amounts of chlorite (0-50%). Micas are usually well foliated. In some chips, biotite occurs in sparse, small clots of unoriented crystals; these are spotted hornfelses. Porphyroblasts of clinozoisite also occur in some biotite-spotted hornfelses.</p> <p>Variable amounts of very fine-grained, black (= carbonaceous material?, magnetite?) are disseminated throughout these rocks. Trace amounts of goethite occur (after disseminated magnetite?, pyrite?).</p> | 27                        | 73                         |
| <p><u>Diorite:</u> Composed of 0.3 to 1.5 mm long laths of plagioclase, interstitial chlorite-clay altered mafic crystals which comprise 7 to 15% of rock. Interstitial quartz forms 2 to 4% of rock. Alteration of plagioclase varies from chip to chip; sericite-clay<sup>±</sup>carbonate replacement of feldspar ranges between 20 and 70%. In the least altered chip, disseminated magnetite/ilmenite forms 1 to 2% of rock.</p>   | 4                         | 11                         |
| <p><u>Metasandstones:</u> Fine-grained; sand grains of quartz; one contains 5-10% carbonate as cement; one is moderately foliated.</p>  | 3                         | 8                          |
| <p><u>Marble:</u> Fine- to medium-grained.</p>  | 2                         | 5                          |
| <p><u>Clay-Sand-Lithic Matrix Material:</u> Clayey matrix containing sand-sized, angular crystal fragments of quartz, feldspar and pyroxene. This material is attached to one of the phyllite chips; presumably, the clay-sand material is the matrix for the alluvial clasts.</p>  | 1                         | 3                          |
| Total Number of Particles Examined  | 37                        | 100%                       |

290'-300'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u> : Very fine-grained; foliated.  | 11                        | 10                         |
| <u>Phyllite</u> : Usually composed of foliated, fine-grained sericite and biotite in variable proportions and of silt-sized quartz and feldspar. Also contains very fine-grained, disseminated, black material (=graphite?, magnetite?). | 20                        | 18                         |
| <u>Diorite</u> : Mafic minerals altered to chlorite, montmorillonite and calcite.  | 14                        | 12                         |
| <u>Metasandstone</u> : Composed of fine-grained, well sorted quartz; calcite cement present in some particles.   | 25                        | 22                         |
| <u>Siltstone</u>   | 1                         | 1                          |
| <u>Clayey Siltstone</u>  | 2                         | 2                          |
| <u>Mudstone</u> : Composed of very fine-grained sericite/illite, sometimes with biotite porphyroblasts. Traces of goethite (after pyrite?) in a few chips.   | 14                        | 12                         |
| <u>Volcanics</u> : Rhyolite?, composed of feathery feldspar crystals and quartz.   | 3                         | 3                          |
| <u>Marble</u>  | 5                         | 4                          |
| <u>Chert</u> : Recrystallized. Composed of very fine-grained, feathery crystals of quartz; contains variable proportions of silt-sized quartz grains.  | 16                        | 14                         |
| <u>Vein Quartz</u>   | 2                         | 2                          |
| Total Number of Particles Examined   | 113                       | 100%                       |

390'-400'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u> : Very fine-grained; foliated; $\pm$ mica; contains abundant, fine-grained, carbonaceous? material. Black or dark grey in hand-specimen.  | 2                         | 2                          |
| <u>Phyllite</u> : Fine-grained; foliated; contains variable proportions of sericite and biotite, as well as, silt-sized grains of quartz and feldspar. Very fine-grained, black material (graphite?, Fe-oxides?) is disseminated throughout most clasts.   | 43                        | 33                         |
| <u>Diorite</u> : Sericite-chlorite-clay altered.   | 2                         | 2                          |
| <u>Metasandstones</u> : Such particles range in composition from nearly pure quartzites to containing 30% sericite $\pm$ chlorite and/or calcite. Goethite/hematite (after pyrite?) porphyroblasts(?) poikilitically encloses quartz grains in one chip; no other chips contained disseminated goethite and/or pyrite. | 65                        | 50                         |
| <u>Clayey Siltstone</u> : Contains abundant sericite/illite as well as silt.   | 2                         | 2                          |
| <u>Mudstone</u> : Composed of very fine-grained unfoliated sericite and chlorite in varying proportions; also contains 0 to 30% silt.  | 3                         | 2                          |
| <u>Granite</u>   | 1                         | 1                          |
| <u>Marble</u> : Fine- to medium-grained.   | 4                         | 3                          |
| <u>Chert</u> : Recrystallized to feathery quartz.  | 6                         | 5                          |
| <u>Pyrite</u> : Unoxidized; a free, 0.5 mm grain.  | 1                         | 1                          |
| <u>Chlorite Schist</u>   | 2                         | 2                          |
| Total Number of Particles Examined   | 131                       | 100%                       |

490'-500'

| Rock Type  | Number of Particles | Percentage of Sample |
|--|---------------------|----------------------|
| <u>Limestone</u> : Very fine-grained; usually foliated; carbonaceous; rarely contains phlogopite flakes.   | 15                  | 19                   |
| <u>Phyllite</u> : Very fine-grained; composed of variable proportions of sericite, biotite and silt-sized quartz and feldspar. Foliated. Contains very fine-grained, disseminated black material, graphite?, Fe-oxides?. Coarse-grained Fe-oxides, magnetite $\pm$ hematite; occur in trace amounts as disseminations. | 23                  | 29                   |
| <u>Metasandstone</u>   | 14                  | 18                   |
| <u>Siltstone</u> : Very fine-grained; contains silt-sized quartz and feldspar as well as variable amounts of sericite/illite $\pm$ chlorite.   | 3                   | 4                    |
| <u>Marble</u>  | 12                  | 15                   |
| <u>Chert</u> : Composed of fine-grained feathery quartz with variable amounts of carbonate and silt- and/or sand-sized quartz grains.  | 10                  | 13                   |
| <u>Clay-Sand-Lithic Matrix Material</u> : Sand- and pebble-sized clasts in a clay-rich matrix; this material is probably the matrix of the alluvium.   | 2                   | 3                    |
| Total Number of Particles Examined   | 79                  | 100%                 |



590'-600'

| Rock Type  | Number of Particles | Percentage of Sample |
|--|---------------------|----------------------|
| <u>Limestone</u>   | 5                   | 6                    |
| <u>Phyllite</u>  | 28                  | 36                   |
| <u>Diorite</u>   | 3                   | 4                    |
| <u>Metasandstones</u>  | 26                  | 33                   |
| <u>Siltstone</u>   | 4                   | 5                    |
| <u>Mudstone</u> : Pyritic; very soft; pale grey-green in hand-specimen; disaggregates on wetting; forms about 30% of washed sample, most chips were destroyed in thin-section making process so they are rare in thin-section. | 1                   | 1                    |
| <u>Volcanics</u> : Latite?   | 1                   | 1                    |
| <u>Marble</u>  | 2                   | 3                    |
| <u>Chert</u>   | 6                   | 8                    |
| <u>Vein Quartz</u>   | 1                   | 1                    |
| <u>Clay-Sand-Lithic Matrix Material</u>  | 1                   | 1                    |
| Total Number of Particles Examined   | 78                  | 100%                 |

790'-800'

| Rock Type   | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|---|---------------------------|----------------------------|
| <u>Limestone</u>  | 10                        | 6                          |
| <u>Phyllite</u> : Composed of very fine-grained foliated sericite, biotite (10-70%) and silt-sized grains of quartz and feldspar. Contains 0 to 1% disseminated pyrite. Rarely contains porphyroblasts of feldspar and/or biotite. Almost always contains finely disseminated black material, graphite? | 46                        | 25                         |
| <u>Diorite</u> : Mafic minerals have gone to montmorillonite.   | 3                         | 2                          |
| <u>Metasandstone</u> : Goethite (after pyrite?) in a few particles.   | 49                        | 27                         |
| <u>Siltstone</u> : Usually sericitic.   | 12                        | 7                          |
| <u>Mudstone</u> : Composed of very fine-grained, clay-sized material. Contains 0.1-0.3% pyrite (usually oxidized to goethite); contains 0 to 15% silt-sized quartz and feldspar.  | 26                        | 14                         |
| <u>Volcanics</u> : Andesite, illite-replaced.   | 2                         | 1                          |
| <u>Marble</u>   | 14                        | 8                          |
| <u>Chert</u>  | 11                        | 6                          |
| <u>Vein Quartz</u>  | 1                         | 1                          |
| <u>Vein Calcite</u>   | 1                         | 1                          |
| <u>Quartz and/or Feldspar Schist</u>  | 4                         | 2                          |
| <u>Chlorite Schist</u>  | 1                         | 1                          |
| <u>Sand-Clay Matrix Material</u>  | 1                         | 1                          |
| Total Number of Particles Examined  | 181                       | 100%                       |

890'-900'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u>   | 14                        | 12                         |
| <u>Phyllite</u>  | 17                        | 14                         |
| <u>Diorite</u>   | 1                         | 1                          |
| <u>Metasandstones</u>  | 40                        | 33                         |
| <u>Siltstones</u>  | 14                        | 12                         |
| <u>Mudstone</u>  | 6                         | 5                          |
| <u>Volcanics</u> : Andesite?, altered to clay and hematite.  | 1                         | 1                          |
| <u>Marble</u>  | 10                        | 8                          |
| <u>Chert</u> : Recrystallized.   | 5                         | 4                          |
| <u>Vein Quartz</u> : One piece coated with limonite and quartz; one vein cuts marble; one contains vugs filled with clay-pyrite. | 5                         | 4                          |
| <u>Vein Calcite</u> : Associated with hematite and quartz.   | 1                         | 1                          |
| <u>Schist</u>  | 2                         | 2                          |
| <u>Clay-Sand-Calcite Matrix Material</u>   | 4                         | 3                          |
| Total Number of Particles Examined   | 120                       | 100%                       |

990'-1000'

| Rock Type   | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|---|---------------------------|----------------------------|
| <u>Limestones</u> : Black; foliated; very fine-grained; sometimes contains phlogopite flakes; contains black, carbonaceous? material. | 27                        | 13                         |
| <u>Phyllites</u>  | 25                        | 12                         |
| <u>Diorite</u>  | 1                         | 0.5                        |
| <u>Metasandstones</u>   | 91                        | 45                         |
| <u>Siltstone</u>  | 8                         | 4                          |
| <u>Mudstone</u> : Occasionally contains large flakes of phlogopite.   | 8                         | 4                          |
| <u>Marble</u>   | 27                        | 13                         |
| <u>Chert</u>  | 12                        | 6                          |
| <u>Vein Quartz</u>  | 2                         | 1                          |
| <u>Schist</u>   | 2                         | 1                          |
| <u>Clay-Sand-Lithic Matrix Material</u>   | 1                         | 0.5                        |
| Total Number of Particles Examined  | 204                       | 100%                       |

1090'-1100'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u>   | 13                        | 8                          |
| <u>Phyllite</u>  | 20                        | 12                         |
| <u>Diorite</u>   | 1                         | 1                          |
| <u>Metasandstone</u>   | 87                        | 51                         |
| <u>Siltstone</u> : Some are chloritic.                               | 8                         | 5                          |
| <u>Mudstone</u> : Composed of very fine-grained clay-sized material. | 9                         | 5                          |
| <u>Volcanics</u> : Chlorite-carbonate replaced.                      | 1                         | 1                          |
| <u>Marble</u>  | 13                        | 8                          |
| <u>Chert</u>   | 12                        | 7                          |
| <u>Vein Quartz</u>   | 5                         | 3                          |
| <u>Clay-Sand-Lithic Matrix Material</u>                              | 3                         | 2                          |
| Total Number of Particles Examined                                   | 172                       | 100%                       |

1190'-1200'

| Rock Type   | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|---|---------------------------|----------------------------|
| <u>Limestone</u>  | 25                        | 15                         |
| <u>Phyllite</u>   | 17                        | 10                         |
| <u>Diorite</u>  | 2                         | 1                          |
| <u>Metasandstone</u>  | 81                        | 50                         |
| <u>Siltstone</u>  | 7                         | 4                          |
| <u>Mudstone</u>   | 11                        | 7                          |
| <u>Volcanics</u> : Andesite; plagioclase replaced by montmorillonite and calcite. | 1                         | 0.5                        |
| <u>Marble</u>   | 8                         | 5                          |
| <u>Chert</u> : Recrystallized.  | 9                         | 6                          |
| <u>Vein Quartz</u>  | 1                         | 0.5                        |
| <u>Quartz Schist</u>  | 1                         | 0.5                        |
| Total Number of Particles Examined  | 163                       | 100%                       |

1290' - 1300'

| <u>Rock Type</u>  | <u>Number<br/>of<br/>Particles</u> | <u>Percentage<br/>of<br/>Sample</u> |
|---|------------------------------------|-------------------------------------|
| <u>Limestones</u>   | 13                                 | 8                                   |
| <u>Phyllites</u>  | 32                                 | 20                                  |
| <u>Metasandstones</u>   | 65                                 | 41                                  |
| <u>Siltstones</u>   | 9                                  | 6                                   |
| <u>Mudstones</u>  | 7                                  | 4                                   |
| <u>Volcanics:</u> Andesites, clay-chlorite and clay-hematite<br>replaced. | 4                                  | 3                                   |
| <u>Marble</u>   | 12                                 | 8                                   |
| <u>Chert</u>  | 9                                  | 6                                   |
| <u>Vein Quartz</u>  | 7                                  | 4                                   |
| <u>Quartz-Feldspar Schist</u>   | 1                                  | 0.6                                 |
| <u>Sericite Schist</u>  | 1                                  | 0.6                                 |
| Total Number of Particles Examined  | 160                                | 100%                                |

1390' - 1400'

| Rock Type   | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|---|---------------------------|----------------------------|
| <u>Limestones</u>   | 13                        | 7                          |
| <u>Phyllites</u>  | 38                        | 19                         |
| <u>Diorite</u> : Primary pyroxene preserved in one chip;<br>in most, montmorillonite replaced mafic minerals. | 5                         | 3                          |
| <u>Metasandstones</u>   | 96                        | 49                         |
| <u>Siltstone</u>  | 17                        | 9                          |
| <u>Mudstone</u>   | 14                        | 7                          |
| <u>Volcanics</u> : Andesites; one replaced by hematite-clay;<br>another by montmorillonite-illite.            | 2                         | 1                          |
| <u>Marble</u>   | 7                         | 4                          |
| <u>Chert</u>  | 4                         | 2                          |
| Total Number of Particles Examined  | 196                       | 100%                       |



1490'-1500'

| Rock Type  | Number<br>of<br>Particles | Percentage<br>of<br>Sample |
|--|---------------------------|----------------------------|
| <u>Limestone</u>   | 14                        | 6                          |
| <u>Phyllite</u>  | 35                        | 16                         |
| <u>Diorite</u> : Mafic minerals replaced by chlorite <sup>±</sup><br>clays <sup>±</sup> carbonate. | 8                         | 4                          |
| <u>Metasandstone</u>   | 88                        | 41                         |
| <u>Siltstone</u>   | 18                        | 8                          |
| <u>Mudstone</u>  | 11                        | 5                          |
| <u>Volcanics</u> : Some are partially sericitized.   | 2                         | 1                          |
| <u>Marble</u>  | 27                        | 13                         |
| <u>Chert</u>   | 11                        | 5                          |
| <u>Vein Quartz</u> : In phyllite   | 1                         | 0.5                        |
| <u>Chlorite Schist</u>   | 1                         | 0.5                        |
| Total Number of Particles Examined   | 216                       | 100%                       |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 1 of 10 pages  
 DATE: July 1920  
 DEPTH: 0 TO 150'  
 EXAMINED BY: M. J. Swaneey



| DEPTH         | COLOR                           | SAMPLE DESCRIPTION   |  |
|---------------|---------------------------------|--|--|
|               |                                 | PRIMARY  | SECONDARY  |
| fact          |                                 |  |  |
| 0 to 10 s*    | Black, white, pink, pale orange | Estimated proportions: 25% black, fine-grained, foliated, micaceous limestone; 30% black to green diorite/diabase; 40% white, pink, orange metasandstones; 5% calcite vein fragments; 3% drusy quartz vein fragments                     | Diorite, weakly to strongly clay-chlorite altered. Drusy quartz vein in metasandstone.                             |
| 10 to 20 n*   | Same as above.                  | Estimated proportions: 20% dark grey to black, fine-grained, foliated limestone; 40% black to green, fine to medium-grained diorite/diabase; 10% white to light grey marble; 25% metasandstones; 1-2% drusy quartz.                      | Drusy quartz veins in chlorite-clay altered basalts. Traces of goethite in diorite and metasandstones.             |
| 20 to 30 m*   | Same as above.                  | Based on examination of 64 chips: 10% black foliated limestone; 35% diorite/diabase; 11% black phyllite; 2% sericite phyllite; 27% metasandstone; 5% marble; 2% chert; 5% quartz veins; 5% calcite veins.                                | 2% rhyolite. Traces of goethite in diorite. Drusy quartz veining.  |
| 30 to 40 n*   | Same as above.                  | Based on 69 chips: 15% black limestone; 23% black phyllite; 21% diorite/diabase; 2% sericite phyllite; 2% marble; 25% metasandstone; 2% latite porphyry; 2% calcite veins; 2% quartz veins.  | Traces of goethite (after pyrite?) in metasandstones and diorite. Quartz veins in black phyllite.                  |
| 40 to 50 n    | Same as above.                  | Based on 123 particles: 29% black, foliated, micaceous limestone; 20% diorite/diabase; 14% black phyllite; 6% marble; 2% sericite phyllite; 17% metasandstone; 2% meta siltstone; 2% latite porphyry; 7% clay-sand.                      | Particles; 2% vein calcite; 1% vein quartz. Opal(?) veins occur in clay-sand matrix; goethite in metasandstone.    |
| 50 to 60 n*   | Black, dark grey, white, pink   | Based on 121 particles: 17% black, foliated limestone; 15% black phyllite; 20% diorite/diabase; 7% white and grey marble; 12% white, pink, buff metasandstone; 3% vein quartz; 1% vein calcite; 24% clay-sand matrix.                    | Traces of goethite in metasandstone. Quartz vein fragments.  |
| 60 to 70 n*   | Same as above.                  | Based on 90 particles: 20% black, foliated limestone; 6% black phyllite; 3% sericite phyllite; 25% diorite/diabase; 2% latite porphyry; 8% marble; 1% meta-arkose; 2% chert; 22% metasandstone; 2% vein calcite; 3% clay-sand matrix.    | Clay-sand coated with opal(?) / quartz veins; also occur on limestone and phyllite particles.                      |
| 70 to 80 s*   | Same as above.                  | Based on 75 particles: 25% black, foliated limestone; 18% diorite/diabase; 8% black phyllite; 3% sericite phyllite; 25% marble; 15% metasandstone; 1% vein quartz; 1% vein calcite; 3% clay-sand matrix.                                 | Caliche coats a few diorite particles. Goethite in diorite. Opal/quartz veins on limestones.                       |
| 80 to 90 n*   | Same as above.                  | Based on 87 particles: 24% black, foliated limestone; 21% diorite/diabase; 6% black phyllite; 30% marble; 10% metasandstone; 1% vein quartz; 3% vein calcite; 6% clay-sand matrix.   | Quartz vein in quartzite; weak chlorite-epidote in same diorite chips.   |
| 90 to 100 n*  | Same as above.                  | Based on 77 particles: 30% black, foliated limestone; 27% diorite/diabase; 6% black phyllite; 13% grey/black and white flow-banded marble; 8% metasandstone; 5% siltstone and clayey siltstone; 1% arkose; 3% vein quartz; 8% clay-sand. | Quartz/opal(?) veins in metasandstone; phyllite, clay-sand material, chlorite-clay alteration of diorite.          |
| 100 to 110 n* | Black, white, pink.             | Based on 21 particles examined in thin-section: 35% black, foliated, micaceous limestone; 5% black phyllite; 15% diorite/diabase; 15% white to grey cherty marbles; 10% white, pink metasandstone; 20% chert.                            | Quartz-calcite veins in limestones contain goethite (after pyrite?).   |
| 110 to 120 n* | Same as above.                  | Alluvium. Particle distribution similar to previously described samples.   | Traces of goethite (after pyrite?) in metasandstone and in diorite, clay-chlorite                                  |
| 120 to 130 n* | Same as above.                  | Alluvium. Same as above.   | Diorite weakly to moderately clay-chlorite altered; also andesite. Rare ribbon quartz veins in phyllites.          |
| 130 to 140 s* | Same.                           | Alluvium. Same.  | Traces of goethite (after pyrite?) in diorite, which is chlorite-clay or sericite-clay altered.                    |
| 140 to 150 s* | Same.                           | Based on 69 particles: 17% black, foliated limestone; 6% basalt and andesite; 16% black phyllite; 15% diorite/diabase; 1% meta-arkose; 16% foliated grey and white marble; 10% sericite phyllite; 25% metasandstone; 1% vein quartz;     | 3% vein calcite. Traces of goethite in metasandstone; also rare quartz veins. Chlorite-clay alteration of diorite. |

\* These letters are explained in text of report.

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 2 of 10  
 DATE: July 1980  
 DEPTH: 150' TO 300'  
 EXAMINED BY: M.J. Sweeney



| DEPTH  | COLOR | SAMPLE DESCRIPTION  |   |
|--------|-------|---|---|
|        |       | PRIMARY   | SECONDARY   |
| 150'to | Same  | Based on 63 particles: 10% black, micaceous limestone; 16% black phyllite; 27% diorite; 16% grey and white, foliated marble; 8% clayey metasiltstone; 18% grey, white, buff metasandstone; 3% metaconglomerate; 3% clay-sand. | Diorite, weakly to moderately chlorite-clay altered. Traces of goethite (after py?) in metasandstone.                     |
| 160'n  |       |   |   |
| 160'to | Same  | Alluvium. Particle distribution similar to previously described samples.  | Diorite, chlorite-clay altered. Very minor quartz/opal veining in metasandstone.  |
| 170'n  |       |   |   |
| 170'to | Same  | Alluvium. Same.   | Diorite, chlorite-clay altered.   |
| 180'n  |       |   |   |
| 180'to | Same  | Alluvium. Same  | Traces of goethite (after py?) in some metasandstone chips. Diorite, chl-clay altered.                                    |
| 190'm  |       |   |   |
| 190'to | Same  | Based on 80 particles (37 in thin-section): 1% black limestone; 60% black phyllites and sericite and chlorite phyllites; 11% diorite; 8% marble; 16% metasandstones; 1% vein quartz.  | Diorite, sericite-clay-chlorite-carbonate altered.  |
| 200'n  |       |   |   |
| 200'to | Same  | Alluvium. Same.   | Diorite, strongly chl-calcite altered. Traces of goethite (after py?) in meta sandstone which also contains quartz veins. |
| 210'n  |       |   |   |
| 210'to | Same  | Alluvium. Same.   | Same.   |
| 220'n  |       |   |   |
| 220'to | Same  | Alluvium. Same  | Same.   |
| 230'n  |       |   |   |
| 230'to | Same  | Alluvium. Same  | Same.   |
| 240's  |       |   |   |
| 240'to | Same  | Alluvium. Same.   | Same.   |
| 250'm  |       |   |   |
| 250'to | Same  | Based on 73 particles; 1% black, foliated limestone; 30% black phyllite; 6% chlorite phyllite; 7% diorite; 4% marble; 43% metasandstone; 1% vein quartz; 8% clay-sand.  | Same.   |
| 260'm  |       |   |   |
| 260'to | Same  | Alluvium. Same.   | Same.   |
| 270'm  |       |   |   |
| 270'to | Same  | Alluvium. Same.   | Same.   |
| 280'm  |       |   |   |
| 280'to | Same  | Alluvium. Same.   | Same.   |
| 290's  |       |   |   |
| 290'to | Same  | 113 particles examined in thin-section: 10% black, foliated limestone; 18% phyllite; 12% diorite; 4% marble; 2% clayey siltstone; 1% siltstone; 12% mudstone; 22% metasandstone; 14% chert; 2% vein quartz; 3% latite.        | Diorite, chl-mont-calcite altered. Traces of goethite (after py?) in mudstone.  |
| 300's  |       |   |   |

WELL: SR-3

## GEOLOGICAL REPORT

PAGE: 3 of 10LOCATION: Churchill Co., Nevada

## WELL CUTTINGS SAMPLE DESCRIPTION

DATE: July 1980

ELEVATION: \_\_\_\_\_

DEPTH: 300' TO 450'

HOLE SIZE: \_\_\_\_\_

EXAMINED BY: M. J. Sweeney

| DEPTH              | COLOR               | SAMPLE DESCRIPTION   |  |
|--------------------|---------------------|--|--|
|                    |                     | PRIMARY  | SECONDARY  |
| Feet               |                     |  |  |
| 300' to<br>310' s  | Same.               | Very similar to previous sample. Clay-sand material is attached to surface of some clasts.   | Same.  |
| 310' to<br>320' st | Same.               | Based on 75 particles: 50% black, foliated phyllite; 10% andesite; 5% diorite; 15% gray and white marble; 5% chloritic meta sandstone; 18% meta sandstone; 1% chert; 3-5% clay-sand matrix.                    | Same.  |
| 320' to<br>330' st | Same.               | Alluvium. Same as above.   | Same.  |
| 330' to<br>340' m  | Same.               | Alluvium. Same as above.   | Same.  |
| 340' to<br>350' m  | Black, buff, white. | Based on 50 particles: 26% black phyllite; 20% chlorite phyllite; 2% sericite phyllite; 66% metasandstone; 2% vein quartz; 2% chert.   | Traces of goethite (after py?) in metasandstone; goethite in qtz vein in phyllite.                     |
| 350' to<br>360' m  | Same.               | Alluvium. Same as above.   | Same.  |
| 360' to<br>370' m  | Same.               | Alluvium. Same as above.   | Same.  |
| 370' to<br>380' s  | Same.               | Alluvium. Same as above.   | Same.  |
| 380' to<br>390' s  | Same.               | Alluvium. Same as above.   | Same.  |
| 390' to<br>400' s  | Same.               | 131 particles examined in thin-section: 2% black, foliated limestone; 33% black phyllite; 2% diorite; 3% marble; 2% clayey siltstone; 2% mudstone; 50% metasandstone; 5% chert; 2% chlorite schist; 1% pyrite. | One fragment of fresh pyrite seen. Thin quartz veins in most rock types. Tr goethite in metasandstone. |
| 400' to<br>410' s  | Same.               | Alluvium. Same as above.   | Same.  |
| 410' to<br>420' st | Same.               | Alluvium. Same as above.   | Same.  |
| 420' to<br>430' st | Same.               | Alluvium. Same as above.   | Same.  |
| 430' to<br>440' s  | Same.               | Alluvium. Same as above.   | Same.  |
| 440' to<br>450' s  | Same.               | Alluvium. Same as above.   | Same.  |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 4 of 10  
 DATE: July 1980  
 DEPTH: 450' TO 600'  
 EXAMINED BY: M.J. Sweeney



| DEPTH            | COLOR  | SAMPLE DESCRIPTION   |  |
|------------------|--|--|--|
|                  |  | PRIMARY  | SECONDARY  |
| feet             |  |  |  |
| 450'to<br>460's  | Buff,<br>black,<br>pink                      | Based on 63 particles: 30% black phyllite; 3% fluorite; 3% marble; 62% buff, grey or white metasandstone; 2% clay-sand matrix.   | Traces of goethite (after py?) in metasandstone; goethite also in stevein in phyllite. |
| 460'to<br>470'st | Same.  | Alluvium. Same as above.   | Same.  |
| 470'to<br>480'm  | Same.  | Alluvium. Same as above.   | Same.  |
| 480'to<br>490'm  | Same.  | Alluvium. Same as above.   | Same.  |
| 490'to<br>500'm  | Same.  | Based on 79 particles examined in thin-sections: 19% black, foliated limestone; 29% phyllite; 15% marble; 4% siltstone; 18% metasandstone; 13% chert; 3% clay-sand matrix.             | Same.  |
| 500'to<br>510'st | Same.  | Alluvium. Same as above.   | Traces of goethite (after pyrite?) in meta sandstones.                                 |
| 510'to<br>520'st | Same.  | Alluvium. Same as above.   | Same. Quartz vein in phyllite.   |
| 520'to<br>530's  | Same.  | Alluvium. Same as above.   | Same.  |
| 530'to<br>540's  | Same.  | Alluvium. Same as above.   | Same.  |
| 540'to<br>550'st | Same.  | Alluvium. Same as above.   | Same.  |
| 550'to<br>560'st | Same.  | Alluvium. Same as above.   | Same.  |
| 560'to<br>570'st | Same.  | Alluvium. Same as above.   | Same.  |
| 570'to<br>580'st | Same.  | Alluvium. Same as above.   | Same.  |
| 580'to<br>590'st | Same.  | Alluvium. Same as above.   | Same.  |
| 590'to<br>600'st | Black,<br>green-<br>grey,<br>buff,<br>white. | Alluvium. About 30% of sample is composed of dark green-grey mudstone which contains 0.1 to 0.5 volume % pyrite. Mudstone disaggregates on wetting. Remainder of sample same as above. | Fresh pyrite disseminated in mudstone.   |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 5 of 10  
 DATE: July 1980  
 DEPTH: 600' TO 750'  
 EXAMINED BY: M.J. Sweeney



| DEPTH            | COLOR                     | SAMPLE DESCRIPTION  |  |
|------------------|---------------------------|---|--|
|                  |                           | PRIMARY   | SECONDARY  |
| feet             |                           |   |  |
| 600'to<br>610'st | Same                      | Alluvium. Same as above.  | Traces of goethite (after py?) in metasandstones.                                      |
| 610'to<br>620'st | Same.                     | Alluvium. Same clast types as described above. Sand-clay material attached to a few particles.  | Same.  |
| 620'to<br>630'st | Same.                     | Alluvium. Same as above.  | Same. A few particles from drusy quartz veins. A few, free, partially oxidized py.     |
| 630'to<br>640'st | Same.                     | Alluvium. Same as above. Contains a few, free or loose pyrite crystals.   | An unoxidized pyrite-quartz fragment; py also in chert(?); goethite in metasandstones. |
| 640'to<br>650'st | Same.                     | Alluvium. Same as above.  | Same.  |
| 650'to<br>660'st | Same.                     | Alluvium. Same as above.  | Pyrite unoxidized in some metasandstone particles.                                     |
| 660'to<br>670'm  | Black, green-grey, white. | Based on 68 particles: 1% black limestone; 13% black phyllite; 3% dark red mudstone; 31% diorite/andesite; 6% marble; 4% siltstone; 4% clayey siltstone; 22% metasandstone; 3% chloritic sandstone. | 4% chert; 3% clay-sand matrix. Tr py in metasandstone.                                 |
| 670'to<br>680'n  | Same.                     | Alluvium. Same as above.  | Loose pyrite crystals in sample; calcite veins in clay-sand material.                  |
| 680'to<br>690'n  | Same.                     | Alluvium. Same as above.  | Same.  |
| 690'to<br>700's  | Same.                     | Alluvium. Essentially same as above. See thin-section description.  | Calcite veining in clay-sand material; trace of py in siltstone.                       |
| 700'to<br>710'n  | Same.                     | Alluvium. Same as above.  | Narrow drusy quartz veins in diorite and metasandstone.                                |
| 710'to<br>720'n  | Same.                     | Alluvium. Same as above.  | Goethite (after py?) in metasandstone; loose py grains.                                |
| 720'to<br>730'st | Same.                     | Alluvium. Same as above.  | Same as above; also py in siliceous particles, chert?                                  |
| 730'to<br>740'n  | Same.                     | Alluvium. Same as above.  | Same.  |
| 740'to<br>750'm  | Same.                     | Alluvium. Same as above.  | More loose py crystals in this sample than previous ones.                              |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION



PAGE: 6 of 10  
 DATE: July 1980  
 DEPTH: 750' TO 900'  
 EXAMINED BY: M. J. Sweeney

| DEPTH  | COLOR        | SAMPLE DESCRIPTION  |   |
|--------|--------------|---|---|
|        |              | PRIMARY   | SECONDARY   |
| feet   |              |   |   |
| 750'to | Black,       | Alluvium. Same as above.  | Loose, euhedral pyrites. Py also attached to siliceous particles. Fragments of botryoidal goethite w/ pyrite. |
| 760'st | buff, white  |   |   |
| 760'to | Same         | Alluvium. Same  | Same.   |
| 770'n  |              |   |   |
| 770'to | Same.        | Alluvium. Same  | Same. Py forms about 0.5% of washed sample.   |
| 780'st |              |   |   |
| 780'to | Same         | Alluvium. Same  | Same.   |
| 790'n  |              |   |   |
| 790'to | Black, grey, | Based on 181 particles examined in thin-section: 6% limestone; 25% phyllite; 1% clay-altered volcanics; 2% diorite; 2% marble; 7% siltstone; 27% metasandstone; 1% vein quartz; 1% vein calcite; 2% quartz-feldspar schist.                     | 6% chert; 14% mudstone; 1% sand-clay matrix material. Py in siltstone.  |
| 800'n  | buff, white  |   |   |
| 800'to | Same.        | Alluvium. Same.   | Same.   |
| 810'n  |              |   |   |
| 810'to | Same.        | Alluvium. Same.   | Same as above. Also particles of py associated with calcite and clay - from veins? Transported limonite also. |
| 820'n  |              |   |   |
| 820'to | Same.        | Alluvium. Same.   | Same.   |
| 830'n  |              |   |   |
| 830'to | Same.        | Alluvium. Same.   | Same.   |
| 840's  |              |   |   |
| 840'to | Same.        | Alluvium. Same.   | Same.   |
| 850'n  |              |   |   |
| 850'to | Same         | Alluvium. Same.   | Fragments of drusy qtz veins more abundant than usual.  |
| 860'm  |              |   |   |
| 860'to | Same.        | Alluvium. Same.   | Same.   |
| 870's  |              |   |   |
| 870'to | Same         | Alluvium. Same.   | Same.   |
| 880'st |              |   |   |
| 880'to | Same.        | Alluvium. Same.   | Same.   |
| 890'm  |              |   |   |
| 890'to | Same.        | 120 particles examined in thin-section: 12% limestone; 14% phyllite; 1% andesite; 1% diorite; 8% marble; 12% siltstone; 33% meta sandstone; 4% chert; 4% vein quartz; 1% vein calcite; 5% mudstone; 2% schist; 3% clay-carbonate-sand material. | Py content less than 1%; fragments of transported limonites in veins.   |
| 900'm  |              |   |   |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 7 of 10  
 DATE: July 1980  
 DEPTH: 900' TO 1050'  
 EXAMINED BY: M. J. Sweeney



| DEPTH               | COLOR                     | SAMPLE DESCRIPTION  |   |
|---------------------|---------------------------|---|---|
|                     |                           | PRIMARY   | SECONDARY   |
| 900' to<br>910' m   | Black,<br>buff,<br>white  | Alluvium. Same as previously described sample.  | Most py is oxidized to goethite. Py comprises less than 0.5 volume % of sample. |
| 910' to<br>920' st  | Same.                     | Alluvium. Same.   | Same.   |
| 920' to<br>930' m   | Same.                     | Alluvium. Same.   | Same.   |
| 930' to<br>940' m   | Same.                     | Alluvium. Same.   | Py is about half replaced by goethite.  |
| 940' to<br>950' s   | Same.                     | Alluvium. Same.   | Same.   |
| 950' to<br>960' m   | Same.                     | Alluvium. Same.   | Same.   |
| 960' to<br>970' s   | Same.                     | Alluvium. Same.   | Same.   |
| 970' to<br>980' s   | Same.                     | Alluvium. Same.   | Same.   |
| 980' to<br>990' s   | Same.                     | Alluvium. Same.   | Same.   |
| 990' to<br>1000' s  | Buff,<br>white,<br>black. | 204 particles examined in thin-section: 13% black limestone; 12% black phyllite; 0.5% diorite; 13% white and grey marble; 4% siltstone; 4% mudstone; 45% buff and white metasandstone; 6% grey chert; 1% vein Qtz; 1% schist; 0.5% clay-sand. | Disseminated py is oxidized; pieces of transported limonite crust present.      |
| 1000' to<br>1010' s | Same.                     | Alluvium. Same.   | Loose py crystals are oxidized; otherwise same as above.                        |
| 1010' to<br>1020' s | Same.                     | Alluvium. Same.   | Same.   |
| 1020' to<br>1030' s | Same.                     | Alluvium. Same.   | Same.   |
| 1030' to<br>1040' s | Same.                     | Alluvium. Same.   | Same.   |
| 1040' to<br>1050' s | Same.                     | Alluvium. Same.   | Same.   |



WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 8 of 10  
 DATE: July 1980  
 DEPTH: 1050' TO 1200'  
 EXAMINED BY: M.J. Sweeney



| DEPTH              | COLOR                    | SAMPLE DESCRIPTION  |   |
|--------------------|--------------------------|---|---|
|                    |                          | PRIMARY   | SECONDARY   |
| feet               |                          |   |   |
| 1050'to<br>1060's  | Black,<br>buff,<br>white | Alluvium. Clast lithology similar to description of sample at 990'-1000'  | Goethite (after py?) disseminated through siltstone, metasandstone. Loose py crystals oxidized.       |
| 1060'to<br>1070's  | Same                     | Alluvium. Same.   | Same.   |
| 1070'to<br>1080's  | Same                     | Alluvium. Same.   | Same.   |
| 1080'to<br>1090's  | Same                     | Alluvium. Same.   | Same.   |
| 1090'to<br>1100's  | Same                     | 172 particles examined in thin-section: 8% black, foliated limestone; 12% black phyllite; 10% andesite (chlorite-carbonate altered); 1% diorite; 8% marble; 5% siltstone; 5% mudstone; 51% metasandstone; 7% chert; 3% vein quartz; 2% clay-sand. | Traces of pyrite in siltstone and mudstone.   |
| 1100'to<br>1110'm  | Same                     | Alluvium. Same.   | Same. Loose py grains and all disseminated py oxidized.   |
| 1110'to<br>1120'n  | Same                     | Alluvium. Same.   | Same. Narrow calcite veins in clay-sand lithic material. Some fresh py; most oxidized.                |
| 1120'to<br>1130'm  | Same                     | Alluvium. Same.   | Py all oxidized.  |
| 1130'to<br>1140'st | Same                     | Alluvium. Same.   | Py mostly oxidized.   |
| 1140'to<br>1150'n  | Same                     | Alluvium. Same.   | Py mostly oxidized. Colloidal, transported limonite fragments present as in most of previous samples. |
| 1150'to<br>1160'st | Same                     | Alluvium. Same.   | Same.   |
| 1160'to<br>1170'm  | Same                     | Alluvium. Same.   | Same.   |
| 1170'to<br>1180's  | Same                     | Alluvium. Same.   | Same.   |
| 1180'to<br>1190's  | Same                     | Alluvium. Same.   | Trace of py in thin calcite vein on metasandstone. Most py oxidized.                                  |
| 1190'to<br>1200's  | Same                     | 162 particles examined in thin-section: 10% phyllite; 1% diorite; 1% andesite; 5% marble; 4% siltstone; 50% metasandstone; 6% chert; 0.5% vein quartz; 0.5% qtz-feldspar schist.  | Loose py crystals and disseminated py mostly oxidized.  |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL CUTTINGS SAMPLE DESCRIPTION

PAGE: 9 of 10  
 DATE: July 1980  
 DEPTH: 1200' TO 1350'  
 EXAMINED BY: M. J. Sweeney



| DEPTH               | COLOR           | SAMPLE DESCRIPTION  |   |
|---------------------|-----------------|---|---|
|                     |                 | PRIMARY   | SECONDARY   |
| feet                |                 |   |   |
| 1200' to<br>1210' n | Buff,<br>black  | Alluvium. Clast lithology similar to previous sample.   | Same as above.  |
| 1210' to<br>1220' n | Same.           | Alluvium. Same.   | As above; most py is oxidized.  |
| 1220' to<br>1230' m | Same.           | Alluvium. Same.   | Proportion of fresh py slightly higher. Py-grt vein attached to metasandstone.                    |
| 1230' to<br>1240' n | Same.           | Alluvium. Same.   | Same. Py content less than 0.5 volume %.  |
| 1240' to<br>1250' n | Same.           | Alluvium. Same.   | Same.   |
| 1250' to<br>1260' n | Same.           | Alluvium. Same.   | Same.   |
| 1260' to<br>1270' m | Same.           | Alluvium. Same.<br>Clay-sand matrix material is darker red than in previous samples.  | Same.   |
| 1270' to<br>1280' n | Same.           | Alluvium. Same.   | Same.   |
| 1280' to<br>1290' n | Same.           | Alluvium. Same.   | Same.   |
| 1290' to<br>1300' n | Buff,<br>black. | 160 particles examined in thin-section: 8% black, foliated limestone; 20% black phyllite; 3% clay-altered andesites; 8% marble; 6% siltstone; 4% mudstone; 41% meta sandstone; 6% chert; 4% vein quartz; 1% schist. | Most py oxidized; py occurs as loose crystals and as disseminations in metasandstones, phyllites. |
| 1300' to<br>1310' n | Same            | Alluvium. Same.   | Same.   |
| 1310' to<br>1320' n | Same            | Alluvium. Same.   | Same.   |
| 1320' to<br>1330' n | Same.           | Alluvium. Same.   | Same.   |
| 1330' to<br>1340' n | Same.           | Alluvium. Same.   | Same.   |
| 1340' to<br>1350' n | Same            | Alluvium. Same.   | Same.   |

WELL: SR-3  
 LOCATION: Churchill Co., Nevada  
 ELEVATION: \_\_\_\_\_  
 HOLE SIZE: \_\_\_\_\_

GEOLOGICAL REPORT  
 WELL-CUTTINGS SAMPLE DESCRIPTION

PAGE: 10 of 10  
 DATE: July 1980  
 DEPTH: 1350' TO 1500'  
 EXAMINED BY: M.J. Sweeney



| DEPTH             | COLOR | SAMPLE DESCRIPTION  |  |
|-------------------|-------|---|--|
|                   |       | PRIMARY   | SECONDARY  |
| feet              |       |   |  |
| 1350'to<br>1360'n | Same. | Alluvium.   | Same.  |
| 1360'to<br>1370'n | Same. | Alluvium.   | Same.  |
| 1370'to<br>1380'n | Same. | Alluvium.   | Same.  |
| 1380'to<br>1390'n | Same. | Alluvium.   | Py fresh in mudstones; oxidized in metasandstones. |
| 1390'to<br>1400'n | Same. | 196 particles examined in thin-section: 7% black, foliated limestone; 19% phyllite; 1% clay-altered andesites; 3% diorite (w/ fresh px); 4% marble; 9% siltstone; 7% mudstone; 49% meta sandstone; 2% chert.  |  |
| 1400'to<br>1410'n | Same. | Alluvium.   | Same. Some of loose py grains are fresh.           |
| 1410'to<br>1420'n | Same. | Alluvium.   | Same.  |
| 1420'to<br>1430'n | Same. | Alluvium.   | Same.  |
| 1430'to<br>1440'n | Same. | Alluvium.   | Most py oxidized.                                  |
| 1440'to<br>1450'n | Same. | Alluvium.   | Same.  |
| 1450'to<br>1460'n | Same. | Alluvium.   | Same.  |
| 1460'to<br>1470'n | Same. | Alluvium.   | Same.  |
| 1470'to<br>1480'n | Same. | Alluvium.   | Same.  |
| 1480'to<br>1490'n | Same. | Alluvium.   | Same.  |
| 1490'to<br>1500'n | Same. | 216 particles examined in thin-section: 6% black, foliated limestone; 16% phyllite; 1% clay-altered volcanics; 4% diorite; 13% marble; 8% siltstone; 5% mudstone; 41% meta sandstone; 5% chert; 1% vein quartz; 1% chlorite schist. Some of loose py crystals are unoxidized. |  |