

that slight environmental differences greatly influenced the morphology of the stromatolites. Near Rio Blanco, Colorado, large domed stromatolites can be traced into thin algal laminites over a distance of a few hundred meters. The large domed stromatolites are interpreted as growing on a promontory where wave activity was significant. In protected areas where wave activity was less, the domed structures were replaced by thin algal laminites. Many of the domed structures are elongated perpendicular to the shore line.

Modern stromatolites from Shark Bay are found only where high salinity has eliminated grazing of algae by other organisms. The absence of gastropods and pelecypods associated with the stromatolite units in the Green River Formation probably implies that high salinity was a necessary condition for stromatolite development in Eocene Lakes Gosiute and Uinta also.

**GEOPHYSICAL STUDIES OF A GEOTHERMAL AREA IN THE SOUTHERN RAFT RIVER VALLEY, IDAHO**

Mabey, Don R., Ackermann, Hans, Zohdy, Adel A.R., Hoover, Donald B., Jackson, Dallas B., and O'Donnell, James E., U.S. Geological Survey, Denver Federal Center, Denver, Colorado 80225

The U.S. Geological Survey in cooperation with the Atomic Energy Commission has made gravity, magnetic, refraction seismic, resistivity, audio-magnetotelluric, self-potential, and telluric current surveys in a study of the geothermal resources in the southern Raft River Valley. The geophysical data indicate that the maximum thickness of Cenozoic sedimentary and volcanic rock underlying the valley is about 2 km and that the valley is bounded by normal faults on the east and south and by a complex system of faults on the west. Large gravity, magnetic, and total field resistivity highs within the valley east of Jim Sage Mountains reflect a mass at a relatively shallow depth, which is probably igneous rock but too old to relate directly to a geothermal system. The seismic interpretation divides the valley into four areas where the Cenozoic rocks have distinctive seismic velocities. These areas appear to relate to known or inferred structures and to a suspected zone of shallow warm water. Resistivity anomalies reflect compositional variations in the Cenozoic rocks and variation in degree of induration and alteration. Although no large reservoir of hot water in the Cenozoic rocks has been identified yet, the resistivity soundings show a 2-5 ohmmeter resistivity unit with a thickness of 1 km underlying a large area of the valley, which may in part be indicative of hot water. Observed self-potential anomalies are believed to indicate zones where warm water is ascending to near the surface. The warm water occurring in springs and wells in the area northeast of The Narrows may relate to deep circulation control by the intersection of north-trending faults east of the Jim Sage Mountains with a northeast-trending structure passing through The Narrows.

**REGIONAL GRAVITY AND MAGNETIC STUDIES OF THE SNAKE RIVER PLAIN**

Mabey, Don R., Peterson, Donald L., and Wilson, Carol W., U.S. Geological Survey, Denver Federal Center, Denver, Colorado 80225

The western and eastern parts of the Snake River Plain are distinctly different, as shown by the gravity and magnetic anomalies. Over the western part gravity and magnetic highs are approximately coextensive

**UNIVERSITY OF UTAH  
RESEARCH INSTITUTE  
EARTH SCIENCE LAB.**

with the plain and and by a thick lay depth of about 2 k Group. Over the e magnetic values ar the anomalies are lying geology. Mo appear to be produ Superimposed on th of the crust under appear to reflect which is buried un The Snake River Pl the regions to the faults. The infer plain and the appa the eastern plain Miocene time and t Basin-Range struct that regional tens in the west and th subsidence of the

**INTERNAL STRUCTURE ALLUVIAL FAN**

Macke, David L. University, F Department of Fort Collins,

A small alluvial f source area compos form constant prec alluvial fans but logic record. Hea tations were anlay stratigraphy was p The most signif size and heavy min development of cha of the fan was int sed reworking of t and shifted heavy these erosional ph Heavy mineral c deposits and show fan-head channel d degree of longitud of the fan, and la Anomalous grain num median-grain-s From the apex due Both the heavy distribution in th alluvial deposits economic significa