

SAMPLE LOCATION	OWNER OR WATER USER	DATE OF COLLECTIO (Mo/Yr)	TEMPERATURE °C	pH	SPECIFIC CONDUCTANCE (Dmhos/cm at 25°C)	BICARBONATE (HCO <sub>3</sub> )	CHLORIDE (Cl <sup>-</sup> )	FLUORIDE (F <sup>-</sup> )	MIRRATE (as Na)	SILICA (SiO <sub>2</sub> )	TRITIUM (PCI/litter)	REFERENCES	REMARKS
25N/66E-31a1	—	6-50	32	—	309	141	0	—	24	7.4	34	—	—
15N/67E-13a2	—	6-20	12	7.5	460	200	0	288	39	17	36	2.9	57
15N/66E-25a1	—	6-50	12	—	112	63	0	—	10	3.6	12*	5	—
18N/67E-1c1	—	7-64	12	8.1	975	284	0	—	47	26	122*	—	—
17N/65E-3ab	—	6-80	9	7.2	28	12	0	—	2.2	1.7	2.4	2.4	2.8
17N/66E-15ac	—	6-80	6	7.5	23	8	0	3	2.0	0.7	1.0	0.4	0.04
16N/66E-13a1	—	7-64	13	7.8	287	172	0	—	38	7.8	15*	—	—
16N/66E-34ba	Clev Rch	6-80	12	7.6	84	44	0	35	13	3.2	1.8	0.4	6.4
16N/67E-3aa	Rosers Rch	6-80	16	7.3	580	360	0	285	56	27	20	1.5	0
16N/67E-21d	—	7-64	16	8.0	911	521	0	—	58	30	105*	—	12
15N/66E-21ac	Bastian Rch	6-80	11	8.2	315	204	0	147	53	7.0	3.7	0.5	—
15N/68E-8b	—	7-64	12	8.0	626	346	0	—	65	33	21	—	26
14N/66E-24a1	—	7-64	12	7.8	499	220	0	—	48	26	22*	—	63
14N/67-16dd	—	6-80	13	7.4	445	176	0	236	27	11	43	2.7	46
13N/67E-15d1	—	6-50	18	—	161	84	0	—	17	3.3	14*	—	7.0
13N/67E-18d	—	7-64	12	8.2	395	204	0	—	39	22	12*	—	34
13N/67E-33d	—	7-64	14	8.5	750	239	16	—	61	14	82*	—	52
13N/67E-35d	—	7-64	23	—	158	88	0	—	18	1.0	16*	—	5.8
13N/66E-17cb	—	6-80	10	7.5	48	28	0	—	4.6	2.6	2.5	0.4	4.4
13N/66E-32db	—	6-80	6.5	7.2	37	20	0	9	3.2	1.7	1.8	0.5	2.4
12N/67E-2a	—	6-80	23	7.9	114	86	0	—	20	2.7	9.2	1.1	4.4
11N/65E-55db	—	6-80	12	7.7	335	220	0	160	30	21	7.9	1.3	12
11N/67E-1c1	—	—	12	—	374	220	0	—	58	12	8.2*	—	14
11N/67E-1bc	Shoeshone Rch	6-80	11	7.5	305	196	0	144	47	10	3.8	0.7	6.2
11N/67E-4c	—	6-80	9	7.6	305	200	0	—	48	8.8	1.4	0.4	4.2
11N/66E-31cd	—	6-80	14	7.6	440	196	0	—	42	27	6.8	2.0	11
9N/67E-27a1	—	7-69	21	7.9	236	122	0	—	24	6.8	18*	—	11

\*Sodium plus potassium.

## References:

1. Rush and Kazmi, 1965

NOTE: SAMPLES FOR WATER QUALITY ANALYSIS COLLECTED BY FUGRO NATIONAL EXCEPT WHERE NOTED. ALL ANALYSIS USED IN mg/l EXCEPT WHERE NOTED. FUGRO NATIONAL ANALYSIS FOR DISSOLVED SOLIDS CALCULATED USING THE RESIDUE - ON - EVAPORATION AT 180°C METHOD. OTHER AUTHORS MAY USE DIFFERENT METHODS. NEVADA LOCATIONS BASED ON MT. DIABLO BASELINE AND UTAH LOCATIONS BASED ON SALT LAKE BASELINE AND MERIDIAN.

WATER QUALITY ANALYSES  
SPRING VALLEY, NEVADA
 MX SITING INVESTIGATION  
DEPARTMENT OF THE AIR FORCE - BMO

 TABLE C-1-9  
C-1-9

SPRING VALLEY

Table 3. Inorganic Water Chemistry.

<u>Time of Collection</u>	2000	2000
<u>Date of Collection</u>	9/14/80	9/18/80
<u>Constituent</u>	<u>Concentration, in mg/l</u>	
Calcium	24.5	24.5
Magnesium	12.2	12.2
Sodium	9.5	9.6
Potassium	3.3	3.3
Alkalinity as $\text{HCO}_3$	132.4	133.4
Chloride	11.7	12.1
Sulfate	8.8	8.7
Nitrate as N	1.0	1.1
Fluoride	0.29	0.25
Silica	57	57
Total Dissolved Solids, (Residue at 180° C)	188	193
Hardness as $\text{CaCO}_3$	113	112
Arsenic	0.004	0.006
Copper	< 0.01	< 0.01
Iron	0.03	0.01
Manganese	< 0.01	< 0.01
Mercury	< 0.0002	< 0.0002
Zinc	0.01	0.01

Analyst: Cranmer Engineering, Inc., Grass Valley California.

Tritium Analysis

$\text{H}^3$ (pCi/l $\pm 2\sigma$ )	0±100	0±100
-------------------------------------	-------	-------

Analyst: James M. Montgomery, Inc., Pasadena, California.