

PPh / Θv HAVALLAH / VALMY FMS. (AVERY)

COMPARISON - CONTRAST : WHY ?

IN THE MCCOY PROJECT AREA, THE POSSIBILITY OF DRILLING INTO BOTH THE PPH AND THE Θv IS PRETTY GOOD FOR ANY HOLE DEEPER THAN 2000'. THIS IS DUE TO THE MAJOR THRUST FAULTS THAT HAVE OCCURRED SINCE THE DEPOSITION OF THE VALMY. WHEN ONE LOOKS AT THE LITERATURE, DESCRIPTIONS OF THE TWO FORMATIONS REAR RATHER FRUSTRATING SIMILARITIES ! HOW WILL THE (AMAX) GEOLOGIST LOGGING CUTTINGS KNOW IF HE HAS PASSED FROM THE BEDDED CHERTS, SILTSTONES, SHALES, AND QUARTZITES OF THE PPH, INTO THE BEDDED CHERTS, SILTSTONES, ARGILLITES, AND QUARTZITES OF THE Θv ?

AFTER A RECONNAISANCE HIKE WITH DEAN PICKINGTON TO THE THRUST FAULT CONTACT OF THE PPH, Pp (After cong.) AND Θv (T22N R41E ON LANDER CTY GEOLOGIC MAP), WE DECIDED THAT THE QUARTZITES HOLD ONE GOOD MEANS OF RECOGNITION, AND SEPARATION. THE QUARTZITES OF THE PPH ARE F.M.G. SANDSTONES WHICH RETAIN MOST OF THEIR ORIGINAL GRAIN-SIZE, SHAPE, AND TEXTURE. THEY, LIKE THE Θv QUARTZITES, ARE OFTEN IRON-STAINED, AND THOUGH USUALLY THIN-BEDDED, DO HAVE SOME MASSIVE SECTIONS. THE VALMY QUARTZITES, ON THE OTHER HAND, DO NOT RETAIN ANY ORIGINAL GRAIN SIZE, SHAPE, OR TEXTURE. THESE MASSIVELY BEDDED, OFTEN IRON-STAINED QUARTZITES HAVE A GLASSY TEXTURE; INTERLOCKING, CRYSTALLINE QUARTZITE APPEARANCE TO THEIR GRAINS. (IT IS MORE OF A META-QUARTZITE). THE SAMPLES WE BROUGHT BACK SHOULD BE USED FOR A REFERENCE WHEN LOGGING QUARTZITES IN DEEPER PARTS OF OUR DRILL HOLES. MORE INFORMATION ON OTHER UNITS OF THE VALMY MAY PROVE HELPFUL (CHERT, ARGILLITE, ETC.), HOWEVER ONLY THE QUARTZITES WERE EXPOSED IN THE OUTCROP WE INVESTIGATED. DEAN IS GOING TO EXPLORE OTHER VALMY OUTCROPS LATER THIS WEEK, AND MAY HAVE SOME VALUABLE INSIGHTS THEN.

Mark Avery

NOTE: TRIASSIC QUARTZITES COMPARE WITH THE PPh AS WELL.