

A-38

STATE OF CALIFORNIA
RONALD REAGAN, *Governor*
THE RESOURCES AGENCY
NORMAN B. LIVERMORE, JR., *Administrator*
DEPARTMENT OF CONSERVATION
RAY B. HUNTER, *Director*

BOUGUER GRAVITY MAP OF CALIFORNIA


WEED SHEET

*filed by
folio*

Scale 1:250,000

1973

O'BRIEN RESOURCES



DIVISION OF MINES AND GEOLOGY
WESLEY G. BRUER, *State Geologist*
1416 Ninth Street, Sacramento
95814

Price \$2.00

WEED GRAVITY


EXPLANATORY DATA
WEED SHEET
GEOLOGIC MAP OF CALIFORNIA

OLAF P. JENKINS EDITION

Compiled by Rudolph G. Strand, 1963

INDEX TO GEOLOGIC MAPPING
USED IN THE COMPILATION OF
THE WEED SHEET

THIS DATA SHEET IS A REPRINT OF THE DATA SHEET ACCOMPANYING THE WEED SHEET, GEOLOGIC MAP OF CALIFORNIA, OLAF P. JENKINS EDITION, FIRST PUBLISHED IN 1963. IT HAS NOT BEEN ALTERED. THE GEOLOGY SHOWN ON THE WEED SHEET OF THE BOUGUER GRAVITY MAP OF CALIFORNIA IS ALSO REPRINTED FROM THE GEOLOGIC MAP OF CALIFORNIA, 1963. THE GRAVITY DATA PRESENTED WAS COMPILED IN 1972 AND PUBLISHED IN 1973.



- Aune, Q. A., Reconnaissance geologic map of a part of the Shoehorse Mtn. quadrangle and vicinity, scale 1:62,500, unpublished, 1963.
Danehy, E. A., Unpublished notes on the geology of the Shoehorse Mtn. quadrangle, 1960-61.
- Averill, C. V., 1931, Preliminary report on economic geology of the Shasta quadrangle, California Div. Mines Mining in California v. 27, no. 1, pp. 3-65, Map: Economic geology and location of mines, Shasta quadrangle by Southern Pacific Company Geological Department, scale 1:250,000. (Modified in part by S. J. Rice, California Div. Mines and Geology, unpublished, 1959.)
- Back, William, 1957, Geology and ground-water features of the Smith River plain, Del Norte County, California: U.S. Geol. Survey Water-Supply Paper 1254, 76 pp., Pl. 5, scale 1:62,500. (Additional information by S. J. Rice, ref. 18a.)
- Cater, F. W., Jr. and Wells, F. G., 1953, Geology and mineral resources of the Gasquet quadrangle, California-Oregon: U.S. Geol. Survey Bull. 995-C, pp. 79-133, Pl. 11, scale 1:50,000.
- Churkin, Michael, Jr. and Langenheim, R. L., Jr., 1960, Silurian strata of the Klamath Mountains, California: Am. Jour. Sci., v. 258, pp. 258-273, Fig. 1, scale 1:79,200.
- Davis, Gregory A., Metamorphic and igneous geology of pre-Cretaceous rocks, Coffee Creek area, northeastern Trinity Alps, Klamath Mountains, California, scale 1:21,120, University of California, Berkeley, unpublished Ph.D. thesis, 1961.
Davis, Gregory A., 1963, Structure and mode of emplacement of Caribou Mountain Pluton, Klamath Mountains, California: Geol. Soc. Am. Bull. v. 74, pp. 331-348, Pl. 1, scale approx. 1:21,500, (this map covers only the southern part of this area).
- Hail, W. Roger, Regional geologic map, Shasta Valley area, California Dept. Water Res., scale 1:62,500, unpublished departmental report, 1959.
- Harbaugh, J. W., Reconnaissance geological map of Shasta Lake area, California, scale 1:62,500, Stanford University, unpublished, 1956.
- Holdaway, M. J., Petrology and structure of metamorphic and igneous rocks of parts of northern Coffee Creek and Cecilville quadrangles, Klamath Mountains, California, plate 1, scale 1:42,240, and Fig. 1, scale approx. 1:425,000, University of California, Berkeley, unpublished Ph.D. thesis, 1962. (Glacial deposits in southern part of this area by R. P. Sharp, 1960, Pleistocene glaciation in the Trinity Alps of northern California: Am. Jour. Sci., v. 258, pp. 305-340, Fig. 3, scale approx. 1:150,000.)
- Hotz, P. E., Geology of the Condrey Mountain quadrangle, scale 1:48,000, U.S. Geol. Survey, work in progress 1963.
- Irwin, W. P., 1960, Geologic reconnaissance of the Northern Coast Ranges and Klamath Mountains, with a summary of the mineral resources: California Div. Mines Bull. 179, 80 pp., Pl. 1 (reconnaissance mapping by William P. Irwin and Donald B. Tatlock), scale 1:500,000. (Additional information regarding the geology at Slate Creek, Orleans quadrangle, by the California Dept. Water Resources, W. R. Hansen, B. G. Hicks, and R. S. Ford, Scale 1:2400, unpublished, 1954.)
- Jones, David L., Geologic Map of the Upper Cretaceous strata in the vicinity of Hornbrook, California, scale 1:24,000, U.S. Geol. Survey, unpublished, 1958.
- Mack, Seymour, 1958, Geology and ground-water features of Scott Valley, Siskiyou County, California: U.S. Geol. Survey Water-Supply Paper 1462, 98 pp., Pl. 1, scale 1:62,500. (Modified in part by mapping of Southern Pacific Company geologists, see ref. 22.)
- Mack, Seymour, 1960, Geology and ground-water features of Shasta Valley, Siskiyou County, California: U.S. Geol. Survey Water-Supply Paper 1484, 115 pp., Pl. 1, scale 1:62,500.
- Masson, P. H., Geology of the Gunsight Peak district, Siskiyou County, California, scale 1:33,000, University of California, Berkeley, unpublished M.A. thesis, 1949. (Stratigraphy reinterpreted in part by Southern Pacific Company geologists, see ref. 22.)
- Masson, J. H., 1933, Economic geology of portions of Del Norte and Siskiyou Counties, northwesternmost California: California Div. Mines Rept., v. 29, nos. 1 and 2, pp. 123-160, Pl. 4, scale 1:187,500. (Modified in part by Wm. P. Irwin, see ref. 11.)
- Pratt, Walden P., Geology of a part of the Scott Bar quadrangle, California, scale 1:62,500, Stanford University, unpublished Ph.D. thesis in preparation, 1963. (Marble deposits mapped by James R. Evans, California Div. Mines and Geology, unpublished limestone investigations, 1962. Additions to mapping by Carl Seyfert, Jr., 1963, see ref. no. 21.)
- Rice, S. J., Geology of the Crescent City, Klamath, Orick and Trinidad quadrangles, California, scale 1:62,500, California Div. Mines and Geology, unpublished, 1951-52. (Information regarding the upper Pliocene along the coast, Trinidad quadrangle, by J. Wyatt Durham, University of California, Berkeley, oral communication, 1962. Franciscan volcanic rocks in the Trinidad quadrangle by E. Alexander, J. DeLapp, E. Gladish and R. Skolmen of the Pacific Southwest Forest & Range Experiment Station—U.S. Forest Service, scale 1:31,680, 1959-60.)
- Rice, S. J., Data on the geology of the area adjacent to the Klamath River (Hornbrook quadrangle), California Div. Mines and Geology, unpublished, 1962.
- Romey, W. D., Geology of a part of the Etna quadrangle, Siskiyou County, California, scale 1:31,680, University of California, Berkeley, Ph.D. thesis, 1962.
- Rynearson, G. A. and Smith, C. T., 1940, Chromite deposits in the Seiad Valley, Siskiyou County, California: U.S. Geol. Survey Bull. 922-J, Pl. 40, scale 1:125,000. Map also published in U.S. Geol. Survey Bull. 948-B.
- Seyfert, Carl, Jr., Geology of the north one-half of the Sawyers Bar quadrangle and vicinity, California, scale 1:62,500, Stanford University, unpublished Ph.D. thesis in preparation, 1963.
- Southern Pacific Company, Land Dep't., Regional geologic mapping program, Geologic maps of T48N R3-10W, T43-47N R7-10W, T42N R2-10W, T40-41N R2-9W, T39N R3,4,6-9W, T38N, R4-9W, T36,37N R2-10W MDB&M by R. J. Antil, H. F. Bonham, Jr., J. T. Collier, J. W. Cooksley, Jr., W. L. Coonrad, A. B. Cunningham, E. A. Danehy, R. R. Dorsey, A. R. Glockzin, E. Kojan, R. T. Laird, W. A. Oesterling, G. W. Olcott, D. E. Pruss, J. V. A. Sharp, W. H. Spurck, F. A. Stejer, J. S. Vanderpool, R. E. Willson, scale 1:24,000, unpublished, 1957-1961. (Glacial deposits in the southwestern part of this area by Sharp, R. P., 1960, Pleistocene glaciation in the Trinity Alps of northern California: Am. Jour. Sci., v. 258, pp. 305-340, Figs. 3,5,6, and 7.)
- Stanford Geological Survey (W. R. Dickinson in charge), Geologic map of Seiad Valley area, Siskiyou Co., California, scale 1:62,500, Stanford University, unpublished, 1959.
- Wells, F. G., 1940, Preliminary geologic map of the Grants Pass quadrangle, Oregon: Oregon Dep't. Geol. and Min. Industries, Map 5, by F. G. Wells, G. O. Gates, R. M. Grantham, P. E. Holtz, H. L. James, W. E. Kennett, J. V. Neuman, Jr., G. A. Rynearson, C. T. Smith, E. C. Tabor, and E. J. Tate, scale 1:96,000.
- Williams, Howel, 1934, Mount Shasta, California: Zeitschrift für Vulkanologie, vol. 15, pp. 225-253, Pl. 16, scale 1:240,000. Additions by Southern Pacific Company geological mapping, see ref. 22.)
- Williams, Howel, 1949, Geology of the Macdoel quadrangle: California Div. Mines Bull. 151, pp. 7-60, Pl. 1, scale 1:125,000. Modifications in the Butte Valley area from: Wood, P. R., 1960, Geology and ground-water features of the Butte Valley region, Siskiyou County, California: U.S. Geol. Survey Water-Supply Paper 1491, 150 pp., Pl. 1, scale 1:62,500, and modifications in the northern part of quadrangle by Southern Pacific Company, see ref. 22.)
- Williams, Howel, 1949, Geology of the Macdoel quadrangle: California Div. Mines Bull. 151, pp. 7-60, Pl. 4, Sketch map of geology along Highway 99, Weed to Hilt, Siskiyou County, California, scale 1:125,000.

For a complete list of published geologic maps of this area see Division of Mines and Geology Special Reports 52 and 52-A.

STRATIGRAPHIC NOMENCLATURE— WEED SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>
CENOZOIC	Recent	Qs	RECENT DUNE SAND Dune sand and olderolian sand along the coast.
		Qal	RECENT ALLUVIUM Recent stream channel, flood plain and alluvial fan deposits. Fluvio-glacial deposits (may be Pleistocene in part), and mud flow deposits, notably along Mud Creek on southeast flank of Mt. Shasta.
		Qrv	RECENT VOLCANIC ROCKS: UNDIFFERENTIATED "Shastina" pyroxene andesite—dark gray or black andesitic basalt rich in bright green granules of pyroxene (these lavas are "identical" with the early flows of Shasta and hence it is not certain that they originated from the Shastina vent, Williams, 1934, p. 245). Hypersthene-rich vesicular andesite and basaltic andesite of Gonneseo volcano. Dark vesicular, glass-rich pyroxene andesite of Deer Mountain. Pale gray porphyritic pyroxene andesite lava, the latest flow from Shasta. Pale gray andesite lavas (containing long slender prisms of brownish hornblende) of Black Butte dome. Hornblende andesite talus breccia in Diller Canyon on west side of Mt. Shasta.
	Qrv ^a	ANDESITIC	
	Qrv ^b	BASALTIC	
	Qrv ^p	PYROCLASTIC	
	QUATERNARY	Ql	QUATERNARY LAKE DEPOSITS Recent diatomite at Copco Lake. Semi-consolidated clay, volcanic ash, diatomite, and sand in Butte Valley. Glacial lake sediments near the town of Mt. Shasta.
		Qg	QUATERNARY GLACIAL DEPOSITS Late (Morris Meadow), middle (Rush Creek), and early (Alpine Lake) Wisconsin (age) moraines in the Trinity Alps area. Undifferentiated glacial deposits on Mt. Shasta and elsewhere. Glacial deposits shown on this map are generalized; in some places the areas are exaggerated in size and may include less than 10% glacial debris.
		Qt	QUATERNARY NONMARINE TERRACE DEPOSITS Unconsolidated sand, clay, and gravel terrace deposits.
		Qm	PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS Battery Formation—buff and blue sand units interbedded with white, bluish-gray, brown, and blue clay units.
		Qc	PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS "Older" alluvium, fan and terrace deposits. "Older" surficial agate-bearing deposits northwest of Copco Dam. "Older" dune sand in the Trinidad quadrangle.
		Qpv	PLEISTOCENE VOLCANIC ROCKS: UNDIFFERENTIATED Andesitic basalt and dense gray pyroxene andesite (early flows forming the base of Mt. Shasta).
		Qpv ^r	RHYOLITIC Pale gray rhyolitic dacite characterized by large phenocrysts of plagioclase. White pumiceous lava (probably a glassy dacite) forming the dome south of Gray Butte on Mt. Shasta.
		Qpv ^a	ANDESITIC Porphyritic pale-gray and brown, pyroxene andesites having distinctive platy or shaly habit and abundant basic inclusions, present on Mt. Shasta. Fine-grained andesite or andesitic basalt of the Bear Butte and Spring Hill lava cones. Pale gray andesite and dacite of Gray Butte plug dome. Colorless glass-rich andesite of the twin domes on the north side of Mt. Shasta.
		Qpv ^b	BASALTIC Butte Valley Basalt—smooth craters, black to gray, vesicular olivine basalt (in part Recent). Pale gray to black, dense, massive to vesicular basalt that emanated from the Everett Hill volcano and flowed down the Sacramento River Valley.
Qpv ^p		PYROCLASTIC Well consolidated lapilli tuff and tuff-breccia in the Macdoel 30' quadrangle, (Pleistocene to Recent). Volcanic mud flow deposits and tuff-breccias exposed in walls of Mud Creek Canyon (perhaps oldest exposed rocks of Mt. Shasta). Black tuff, lapilli, and volcanic bombs on Mt. Shasta.	
TERTIARY	QP	PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS Probable Plio-Pleistocene deposits in the Orick quadrangle. Pleistocene or Late Tertiary terrace deposits that have been slightly deformed and contain deeply weathered boulders. (Hoops quadrangle).	
	Q*	QUATERNARY AND/OR PLIOCENE CINDER CONES Pliocene (?) to Recent cinder cones.	
	Pu	UPPER PLIOCENE MARINE SEDIMENTARY ROCKS Small areas of highly fossiliferous marine, near-shore sands and gravels deposited on very irregular surface along the present day coast.	
	Pliocene	Pv ^a	PLEIOCENE VOLCANIC ROCKS: ANDESITIC Pyroxene andesite and hornblende andesite in the Macdoel 30' quadrangle (possibly Plio-Pleistocene).
		Pv ^b	BASALTIC Olivine basalt and basaltic andesite in the Macdoel 30' quadrangle (possibly Plio-Pleistocene to late Pleistocene).
		Pv ^p	PYROCLASTIC Low mounds of red cinders, remnants of cinder cones in Macdoel quadrangle.
	MIOCENE	Tm	TERTIARY MARINE SEDIMENTARY ROCKS St. George Formation—faint gray-blue, massive fossiliferous siltstone and shale containing irregular lenses of sand (Pliocene).
		Mu	UPPER MIOCENE MARINE SEDIMENTARY ROCKS Wimer Formation—friable yellow shale and siltstone that weathers brown.
		Tc	TERTIARY NONMARINE SEDIMENTARY ROCKS Late Tertiary autiferous gravel deposits in the Gasquet and Klamath quadrangles.
		Undifferentiated	Ti
Ti ^r			RHYOLITIC Massive gray and olive-green porphyritic pyroxene dacite at Cedar Lake. Albite rhyolite porphyry dikes in the Gasquet quadrangle.
Ti ^a			ANDESITIC Andesitic volcanic necks.
Ti ^b			BASALTIC Basaltic volcanic neck in vicinity of Copco Dam.
Tertiary volcanic rocks:		Tv	UNDIFFERENTIATED Wason Formation, Raxy Formation, Coleraine Formation—hypersthene augite andesite, basalt, some dacite flows, volcanic conglomerate and sandstone (considered to be upper Eocene to upper Miocene by H. Williams).
		Tv ^r	RHYOLITIC Basaltic flows in the Gazelle area and in the southeast corner of the map area (age uncertain; may prove to be Quaternary).
		Tv ^b	BASALTIC White to pale cream very fine-grained rhyolite plug domes in the Macdoel 30' quadrangle.
	Tv ^p	PYROCLASTIC White to cream, and pale blue-green rhyolite tuff, glassy pumiceous dacite tuff, and andesitic tuff-breccia.	
	Ku	UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS Hornbrook Formation—greenish-gray massive arkosic sandstone, conglomerate, and bluish-gray shale, buff and white massive sandstone (nonmarine in part), cross-bedded sandstone, and coal. Coarse- to medium-grained sandstone, angular boulder conglomerate, and light-gray shale in the Gunsight Peak area.	
FRANCISCAN FORMATION	Kjf	Franciscan Formation—graywacke, interbedded shale, minor conglomerate, thin-bedded chert, some undifferentiated basaltic or siltitic rocks that have been altered to greenschist, and small masses of glauconitic schist. KJF = Dothan Formation—dark gray hard metagreywacke, metaconglomerate, thin-bedded metabasalt, and interbedded green metaconglomerate rocks (probably Late Jurassic, but believed by F. Wells and G. Walker (USGS Map GQ21, 1953) to be older than the Galice Formation shown as Ju on this map; the Franciscan Formation and the lithologically similar Dothan Formation have been extended from their type areas from the south and from the north to the Oregon-California line where the rocks are known by either name).	
	Kjfv	FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS Metamorphosed basic igneous rocks within the Franciscan Formation.	


STRATIGRAPHIC NOMENCLATURE— Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>	
MESOZOIC	JURASSIC	MESOZOIC GRANITIC ROCKS	gr	UNDIFFERENTIATED Quartz diorite, diorite, gabbro, granite and granodiorite that have not yet been differentiated in the mapping. Locally includes some ultramafic rocks.
			gr ^a	GRANITE AND ADAMELLITE (QUARTZ MONZONITE) Adamellite at Yellow Butte. Granite and adamellite in the Seiad Valley quadrangle.
			gr ^d	GRANODIORITE Hornblende-biotite granodiorite, and leucocratic granodiorite in the Etas and Coffee Creek quadrangles. Granodiorite in the Seiad Valley quadrangle.
			gr ^t	TONALITE (QUARTZ DIORITE) AND DIORITE Tondalite in Caribou Mtn. area (Coffee Creek quadrangle), tonalite, hornblende diorite, biotite diorite, and pyroxene diorite in the Etas and Coffee Creek quadrangles. Tonalite in the Seiad Valley quadrangle.
			bi	MESOZOIC BASIC INTRUSIVE ROCKS Hornblende gabbro and related dark diorite, augite-hornblende gabbro, hornblende gabbro and hornfels complex in the Coffee Creek quadrangle. Hornblende gabbro, and undifferentiated gabbro and related dark diorite in the Etas quadrangle. Gabbro, undifferentiated gabbro and diorite, and minor granodiorite elsewhere.
	TRIASIC	MESOZOIC ULTRABASIC INTRUSIVE ROCKS	ub	Predominantly serpenitized peridotite. Includes some unaltered peridotite, dunite, and pyroxenite, also lamplike deposits of ultramafic rocks in the Coffee Creek quadrangle. Most of the ultramafic rock in the eastern part of the map area may be part of a once-continuous sheet that intruded a low-angle fault along which Paleozoic strata to the east were thrust westward over the Abrams and Salmon Formations (Irwin and Lipman USGS Prof. Paper 410-C Article 67, 1962).
			Ju	UPPER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Galice Formation—dark gray to black slate, phyllite, and interbedded thin to massive, light gray, tauffaceous sandstone; includes some interbedded andesitic metaconglomerate.
			Jml	MIDDLE AND/OR LOWER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Potem Formation—argillite, tauffaceous sandstone and shale, and limestone; Arvision Formation—(pyroclastic rocks deposited in a marine environment) volcanic breccia and agglomerate containing interbedded tuff, tauffaceous sandstone and conglomerate, some limestone.
			Jrv	JURASSIC AND/OR TRIASSIC METAVOLCANIC ROCKS Metavolcanic rocks in the northwestern part of the Weed sheet (may be in part correlative with the Rogue Formation in Oregon and are in part interbedded within the Galice Formation). Bagley Andesite—fine grained, volcanic breccia and pyroclastic deposits (Middle Jurassic). Pit Formation, (volcanic part of)—Keratophyre, light to dark colored flow rocks, mafic flows and some pyroclastic rocks (Triassic).
			R	TRIASIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Medin Formation—shale, sandstone, tuff, agglomerate, andesite flow rocks, limestone, and conglomerate. Buck Shale(?)—thin dark shale, in part calcareous; Houselius Limestone—gray, lenticular, fossiliferous limestone; Pit Formation—slate, siltstone and sandstone and some undifferentiated volcanic rocks.
			m	PRE-CRETACEOUS METAMORPHIC ROCKS, UNDIFFERENTIATED Chiefly undivided phyllite, chert, and metavolcanic rocks of the Western Paleozoic and Triassic Belt of Irwin (1960), (in part correlative with the Applegate Group of southwestern Oregon); includes rocks of the so-called Southwestern Devonian and Southwestern Carboniferous Belts of Diller, the Blue Chert and part of the Lower State series of Herberich, and the Grayback Formation of Maxson (1933). In Scott Bar and Conroy Mtn. quadrangles, includes amphibolite, quartzite, and marble. Stuart Fork Formation—phyllite, quartzite, argillite, quartz-mica phyllite, gneiss and related tuff, minor marble (formerly included within the Abrams Mica Schist, Davis and Lipman, G.S.A. Bull., Dec., 1962. This formation is present in the Coffee Creek quadrangle and is probably correlative with similar rocks elsewhere in the central metamorphic belt.)
			ls	ls = LIMESTONE AND/OR DOLOMITE White, coarsely crystalline limestone (marble) in the Seiad quadrangle and the Marble Mountains.
			ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS Quartzite, metachert, and mica schist on Yellow Butte (Macdoel 30' quadrangle). Phyllite, blue-gray thin-bedded chert, some volcanic rocks, and minor limestone have been considered as correlative with the Triassic Applegate Group. Rocks considered to be more highly metamorphosed equivalents of the Galice Formation, includes Wetschpe Schist of Oscar Herberich (1904).
			mv	PRE-CRETACEOUS METAVOLCANIC ROCKS Greenstone and greenstone schist having metamorphic interbeds of chert, argillite and limestone (these rocks may be correlative with the Applegate Group). Metabasite intrusive rock south of Gunsight Peak northward to Paradise Crater in the Hornbrook quadrangle. The "mv" unit north of Seiad Valley depicts coarse-grained, foliate and/or laminate black amphibolite of metamorphic grade similar to that of the adjacent "ps" unit. However, F. Wells (personal communication to W. Dickinson) believes that the black amphibolite resembles metamorphic phases of undivided Applegate Group, whereas the hornblende schist ("ps" unit) is unlike anything known to be Applegate Group or its equivalent.
			PALEOZOIC	PERMIAN
R	PERMIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS McCloud Limestone—light gray, coarsely crystalline limestone, some dark gray massive or thin-bedded limestone.			
MISSISSIPPIAN	C	UNDIVIDED CARBONIFEROUS MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Baird Formation—Metamorphosed maroon and green mudstone, multicolored conglomerate, sandstone, tuff, and dark gray limestone (probably Mississippian).		
	CM	MISSISSIPPIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Bradon Formation—Metamorphosed shale, siltstone, mudstone, sandstone and coarse conglomerate.		
	D	DEVONIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Kennett Formation—Metamorphosed black shale and mudstone, and dark gray dense coral reef metamorphosed limestone.		
	Dv?	DEVONIAN AND PRE-DEVONIAN ? METAVOLCANIC ROCKS Copley Greenstone—Green meta-andesite that weathers buff, irregularly interbedded flows, tuff, and tuff-breccia.		
	S	SILURIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Gazelle Formation—Essentially unmetamorphosed volcanic graywacke, dark gray to black siltstone and mudstone, siliceous and feldspathic grit, chert conglomerate, limestone and limestone conglomerate.		
	O	ORDOVICIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS Dazel Formation—Pale gray-green schistose graywacke, phyllite, limestone, chert, and greenstone.		
	ps	PRE-SILURIAN ? METAMORPHIC ROCKS, UNDIFFERENTIATED Limestoned hornblende schist and medium to dark green plagioclase-chlorite schist in the Seiad Valley quadrangle. Chlorite-quartz-mica schist in the Conroy Mtn. and Seiad Valley quadrangles. Hornblende schist in the Preston Fork quadrangle. Hornblende and chlorite schists in the Scott Bar quadrangle. These units have been considered as equivalents of the Salmon Hornblende Schist and Abrams Mica Schist, however they may be nothing more than more highly metamorphosed equivalents of Triassic or Paleozoic rocks to the south. Dickinson (1945 written communication) believes that the amphibolites shown as "mv" north of Seiad Valley are of approximately equivalent metamorphic grade to the adjacent "ps" schists.		
	psS	PRE-SILURIAN ? METASEDIMENTARY ROCKS Quartz-mica schists that have been considered as Abrams Mica Schist. Grouse Ridge Formation (Davis and Lipman, G.S.A. Bull., Dec., 1962)—micaceous and feldspathic quartz schists, almandine-hornblende rocks, hornblende schists, hornblende gneiss, and calc-schists. Rocks shown as ps ^b (igneous amphibolites) are considered by Lipman (1962) as part of the Salmon Hornblende Schist.		
psv	PRE-SILURIAN ? METAVOLCANIC ROCKS Salmon Hornblende Schist—limestoned hornblende schist probably formed by the metamorphism of basaltic rocks. Included in psv is a "transitional" Stuart Fork-Salmon unit of Lipman and Davis, the mapped northward extension of this unit is however, a greenstone that has a similar metamorphic history to that of the Stuart Fork Formation. Davis and Lipman (G.S.A. Bull., Dec., 1962) have postulated that the Stuart Fork-Salmon contact is a major low angle thrust fault.			

NOTES

¹ These units are equivalent to a part of the Western Cascade Volcanic Series of H. Williams.
² Miller, J. S., 1903, Klamath Mountains section, California: Am. Jour. Sci., 4th ser., vol. 15, pp. 342-362.
³ Herberich, O. H., 1901, Metamorphic formations of northwestern California: Am. Geologist, vol. 37, pp. 225-245.
⁴ Herberich, O. H., 1906, Some western Klamath stratigraphy: Am. Jour. Sci., 4th ser., vol. 21, pp. 1-44.
⁵ The Dazel Formation was first described by F. Wells, G. Walker, and C. Merriam (G.S.A. Bull., May, 1959) as Upper Ordovician (?) based upon a fossil fauna at Horseshoe Gulch. This age was determined by Helen Duncan (of the U.S.G.S.) who recognizes the less likely possibility that this fauna may have been Silurian. However, she recognizes the possibility that both Ordovician and Silurian strata may be present.
⁶ The Dazel Formation was first described by F. Wells, G. Walker, and C. Merriam (G.S.A. Bull., May, 1959) as Upper Ordovician (?) based upon a fossil fauna at Horseshoe Gulch. This age was determined by Helen Duncan (of the U.S.G.S.) who recognizes the less likely possibility that this fauna may have been Silurian; however, she recognizes the possibility that both Ordovician and Silurian strata may be present.

TOPOGRAPHIC QUADRANGLES
 WITHIN THE WEED SHEET
 AVAILABLE FROM THE U.S. GEOLOGICAL SURVEY
 FEDERAL CENTER, DENVER, COLORADO 80225
 1963



Mt. Shasta as seen from the west, extends upward to 14,162 feet and forms a classic example of a Quaternary composite volcanic cone. Shastina is the large parasitic composite cone that appears on the near flank of the mountain. The wedge-shaped gash on Shastina is Diller Canyon, a feature which Howel Williams suggests may be due to violent downward explosions.

Photo by Fairchild Aerial Surveys, Inc.
 Oct. 5, 1930

INDEX TO CALIFORNIA GRAVITY MAP SHEETS

The date beneath each map sheet name indicates year of publication. If no date appears, the gravity map sheet had not been published at the time this cover was issued.

