


A-38




STATE OF CALIFORNIA
 THE RESOURCES AGENCY
 DEPARTMENT OF CONSERVATION

BOUGUER GRAVITY MAP OF CALIFORNIA
 NEEDLES SHEET

Scale 1:250,000
 1978

*filed in
 folio*



CALIFORNIA DIVISION OF MINES AND GEOLOGY
 1416 Ninth Street, Room 1341
 SACRAMENTO, CALIFORNIA 95814

NEEDLES GRAVITY


EXPLANATORY DATA
NEEDLES SHEET
GEOLOGIC MAP OF CALIFORNIA

OLAF P. JENKINS EDITION

Compiled by Charles C. Bishop, 1963

THIS DATA SHEET IS A REPRINT OF THE DATA SHEET ACCOMPANYING THE NEEDLES SHEET, GEOLOGIC MAP OF CALIFORNIA, OLAF P. JENKINS EDITION, FIRST PUBLISHED IN 1964. IT HAS NOT BEEN ALTERED. THE GEOLOGY SHOWN ON THE NEEDLES SHEET OF THE BOUGUER GRAVITY MAP OF CALIFORNIA IS ALSO REPRINTED FROM THE GEOLOGIC MAP OF CALIFORNIA, 1964. THE GRAVITY DATA PRESENTED WERE COMPILED IN 1976 AND PUBLISHED IN 1978.

INDEX TO GEOLOGIC MAPPING
USED IN THE COMPILATION OF THE
NEEDLES SHEET



1. Anonymous, Reconnaissance geology of the Granite Mountains area, scale 1:62,500, unpublished, 1958.
2. Anonymous, Reconnaissance geology of the Whipple Mountains area, scale 1:62,500, unpublished, 1961-62.
- 3a. Bishop, Charles C., Photogeologic interpretation of part of the Stepladder Mountains quadrangle, California, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1963.
- 3b. Bishop, Charles C., Photogeologic interpretation of the northern part of the Colton Well quadrangle, California, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1963.
- 4a. Bishop, Charles C. and Jennings, Charles W., Reconnaissance geologic map and photogeologic interpretation of parts of the Milligan, Essex, Cadiz Lake and Danby quadrangles, California, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1963.
- 4b. Bishop, Charles C. and Jennings, Charles W., Reconnaissance geologic map and photogeologic interpretation of alluvial and adjacent bedrock areas, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1963.
5. Evans, James R., Xenotime mineralization in the southern Music Valley area, Riverside County, California: California Div. Mines and Geology Special Rept. 79, 24 pp., in press, 1963, Map 1: Geologic map of the southern Music Valley, scale 4 inches = 1 mile.
6. Gale, H. S., 1951, Geology of the saline deposits of Bristol Dry Lake, San Bernardino County, California: California Div. Mines and Geology, Special Rept. 13, 21 pp., pl. 1, scale 1:62,500.
7. Giessner, F. W., Map of the Rice and Vidal Valley areas, California (showing reconnaissance geology and

- location of wells and springs) scale 1:62,500, U. S. Geol. Survey, Ground Water Branch, work in progress, 1962; to be published by Calif. Dept. Water Resources in Bull. 91-8, 1963.
8. Hamilton, Warren B., Geologic map of the Riverside and West Riverside Mountains, scale 1:24,000, U. S. Geol. Survey work in progress, 1963.
9. Hazzard, John C., 1933, Notes on the Cambrian rocks of the eastern Mojave Desert, California: California Univ. Dept. Geol. Sci. Bull., v. 23, no. 2, Map 1, scale 1:187,500. Southern Pacific Company, Land Dept., Regional geologic mapping program, geologic map of part of T5N R14E SBBM, scale 1:24,000, unpublished, 1959-60. Lanphere, M. A., Geochronologic studies in the Death Valley-Mojave Desert region, California, Pl. 8; Geologic map of part of the Marble Mountains, scale 1:24,000, Calif. Inst. of Tech., unpublished Ph.D. thesis, 1962. Lanphere, M. A., Geochronologic studies in the eastern Mojave Desert, California: Jour. Geol., in press, 1964.
10. Hazzard, John C., 1954, Rocks and structure of the northern Providence Mountains, San Bernardino County, California, in California Div. Mines and Geology, Bull. 170, Geology of Southern California, Chpt. IV, contr. 4, Pl. 2, scale 1:31,680.
11. Kennitzer, Luis E., Structural studies in the Whipple Mountains, southeastern California, scale 1:62,500, Calif. Inst. of Tech., unpublished Ph.D. thesis, 1937.
12. Kupfer, Donald H. and Bassett, Allen M., 1962, Geologic reconnaissance map of part of the southeastern Mojave Desert, California: U. S. Geol. Survey Mineral Investigations, Field Studies Map MF-205, scale 1:125,000.
13. Moyle, W. R., Jr., 1961, Data on water wells in the Dale Valley area, San Bernardino and Riverside Counties: California Dept. Water Resources, Bull. 91-5, 55 pp., fig. 2, scale 1:62,500, prepared by U. S. Geol. Survey, Ground Water Branch.

14. Noble, J. A., Reconnaissance geologic map and photogeologic interpretation of a part of the Whipple Mountains along the Colorado River, scale 1 in. = 880 feet and 1 in. = 1080 feet, unpublished 1963.
15. Noble, L. F., 1931, Nitrate deposits in southeastern California: U. S. Geol. Survey Bull. 820 (information concerning Quaternary lake bed deposits on p. 39, no map).
16. Rogers, Thomas H. and Jennings, Charles W., Reconnaissance geologic map and photogeologic interpretation of the southern part of the Valley Mountain and Dale Lake quadrangles, California, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1964. (In part after anonymous mining company reconnaissance geologic map of the Pinto-Sheephole Mountains area, scale 1:62,500, unpublished, 1957).
17. Saul, Richard B., Reconnaissance geology and photogeologic interpretation of the Turtle Mountain quadrangle and parts of the Vidal, Savahia Park, and Rice quadrangles, California, scale 1:62,500, California Div. Mines and Geology reconnaissance mapping for the State Geologic Map, 1963.
18. Southern Pacific Company, Land Dept., Regional geologic mapping program, geologic maps of all or parts of T3-11N, R9-18E; T2N, R11-16E; T5-11N, R19-24E SBBM by R. Antcliff; H. F. Bonham, Jr.; J. T. Collier; J. W. Cooksley; W. L. Coonrad; A. B. Cunningham; E. A. Danehy; James Gamble; R. T. Laird; Max Schafer; W. H. Spurck; M. S. Tischler, scale 1:24,000, unpublished, 1958-1960. Field checked and modified in part by California Div. Mines and Geology, 1963.

For a complete list of published geologic maps of this area see Division of Mines and Geology Special Reports 52 and 52-A.



View west from Coffin Spring in the northern part of the Turtle Mountains showing typical exposures of Tertiary volcanic rocks. Light colored pyroclastic material is interbedded with darker andesitic flow rocks. Photo by Richard B. Saul, 1963

STRATIGRAPHIC NOMENCLATURE—NEEDLES SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>
QUATERNARY	Recent	Qs	RECENT DUNE SAND
		*	QUATERNARY CINDER CONES
		Qrvb	RECENT VOLCANIC ROCKS: BASALTIC
		Qal	RECENT ALLUVIUM
		Qst	QUATERNARY SALT DEPOSITS
	Pleistocene	Ql	QUATERNARY LAKE DEPOSITS
		Qc	PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS
		Qpvb	PLEISTOCENE VOLCANIC ROCKS: BASALTIC
		Qvp	PYROCLASTIC
		QP	PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS
CENOZOIC	Miocene	Mv	MIOCENE VOLCANIC ROCKS: UNDIFFERENTIATED
		Mva	ANDESITIC
		Mvb	BASALTIC
		Mvp	PYROCLASTIC
		Tc	TERTIARY NONMARINE SEDIMENTARY ROCKS
	Undivided	Tjr	TERTIARY INTRUSIVE (HYPABYSSAL) ROCKS: RHYOLITIC
		Tja	ANDESITIC
		Tv	TERTIARY VOLCANIC ROCKS: UNDIFFERENTIATED
		Tva	ANDESITIC
		Tvb	BASALTIC
Cretaceous	gr	MESOZOIC GRANITIC ROCKS: UNDIFFERENTIATED	
	gra	ADAMELITE (QUARTZ MONZONITE)	
	grf	TONALITE (QUARTZ DIORITE)	
	bi	MESOZOIC BASIC INTRUSIVE ROCKS	
	Jrv	JURASSIC AND/OR TRIASSIC METAVOLCANIC ROCKS	
MESOZOIC	Triassic	R	TRIASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS


STRATIGRAPHIC NOMENCLATURE—Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit are listed in stratigraphic sequence from youngest to oldest.)</small>		
MESOZOIC	UNDIVIDED	PRE-CRETACEOUS METAMORPHIC ROCKS: UNDIFFERENTIATED	Metasedimentary to meta-igneous rock complex including hornfels, granophyre, phyllite, metadiorite, gneiss, and quartzite, with some Mesozoic intrusive rocks. Greenschist, greenstone, metamorphosed clastic sediments and subordinate metacarbonate rocks (Riverside Mountains).		
		Is	Is = LIMESTONE AND/OR DOLOMITE	Metacarbonate rocks of the Bristol Mountains, locally include tactite, splite, and other intrusive rocks. Dolomite marble and tactite in the Pinto Mountains.	
		ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS	White, buff, and gray quartzite in the Pinto Mountains.	
		mv	PRE-CRETACEOUS METAVOLCANIC ROCKS	Undifferentiated metavolcanic rocks in Valley Mountain northeast of Twentynine Palms.	
		gr-m	PRE-CENOZOIC GRANITIC AND METAMORPHIC ROCKS	Migmatites of Precambrian complex rocks and Mesozoic plutonic rocks (Calumet Mountains, Fenner Hills, and Piute Mountains). Undifferentiated plutonic and metamorphic rocks.	
	PALEOZOIC	UNDIVIDED	IP	PALEOZOIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Metasedimentary rocks of probable Paleozoic age, including metadolomite, white marble, varicolored marble and quartzite (Riverside Mountains and Arica Mountains).
			IPs	Is = LIMESTONE	Marble, locally cherty and fossiliferous (crinoid stems and brachiopods) in the Marble Mountains.
		PERMIAN	R	PERMIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Bird Spring Formation—fossiliferous limestone with minor amounts of calcareous shale and chert (Providence Mountains); lower part of Permian age. Limestone with minor calcareous shale (Ship Mountains).
			CP	PENNSYLVANIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Limestone and minor black chert (northwestern part of the Ship Mountains).
			CM	MISSISSIPPIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Monte Cristo Limestone—includes: Yellow Pine, Anchor, Bullion, and Dawn Limestone members and a basal sandstone member (Providence Mountains).
CAMBRIAN	DEVONIAN	D	DEVONIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Sultan Limestone—limestone and dolomite with local thin sandstone beds (Providence and southern Marble Mountains).	
		ε	CAMBRIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS	Cambrian sedimentary section in the Providence and southern Marble Mountains including: Upper Cambrian "Cornfield Springs" Formation—dolomite and shale; Middle Cambrian Bonanza King Formation—dolomite and partially dolomitized limestone, locally cherty and sandy; Cadiz Formation—muddy limestone, shale and quartzite; Lower Cambrian Chambliss Limestone—limestone with algal nodules; Latham Shale—platy shale with thin sandy limestone layers. Also includes small outcrop area of Prospect Mountain Quartzite of probable Cambrian age.	
	PRECAMBRIAN	pε	UNDIVIDED PRECAMBRIAN METAMORPHIC ROCKS: UNDIFFERENTIATED	Essex Series (in part)—quartzite, marble, dolomite, and interbedded schists. pCh = marble in the Old Woman Mountains.	
		pεs	SCHIST	Chlorite schists of the Essex Series ¹ (Old Woman Mountains).	
		pεg	GNEISS	Augen gneiss, granite to dioritic gneiss ² , Fenner Gneiss ² , Kilbeck Gneiss ² , Essex Series ² , and Pinto Gneiss ² ; includes meta-igneous intrusive rocks.	
pεgr	UNDIVIDED PRECAMBRIAN GRANITIC ROCKS	Plutonic igneous rocks ³ varying in composition from granite to diorite and gabbro ³ . Generally slightly to highly foliated or gneissic.			
pεc	PRECAMBRIAN IGNEOUS AND METAMORPHIC ROCK COMPLEX	Intimate mixture of Precambrian plutonic rocks (diorites to granites) and Precambrian metamorphic rocks, including migmatites. ⁴			
epε	EARLIER PRECAMBRIAN METAMORPHIC ROCKS	Augen gneiss (western Providence Mountains and southern Homer Mountains).			

NOTES

- Fossil vertebrates found in the Sacramento Mountains are of a fairly primitive species of *Merychippus* and are probably middle Miocene according to John F. Lance, personal communication 12/18/1963.
- Radiometric dates of granite rocks from the following areas indicate:
 Marble Mountains (central part)
 165 million years (minimum), Pb²⁰⁶/U²³⁸, L. T. Silver, personal communication, 2/5/1964.
 Ship Mountains
 150 million years (minimum), Pb²⁰⁶/U²³⁸, L. T. Silver, personal communication, 2/5/1964.
 Piute Mountains
 78.3 million years (± 3.0 m.y.), K/A, Geochron Laboratories, for Calif. Div. Mines and Geology, Sept. 1963.
 West Riverside Mountains
 98.5 million years (± 4.0 m.y.), K/A, Geochron Laboratories, for Calif. Div. Mines and Geology, Sept. 1963.
 Pinto Mountains
 163 million years (± 7 m.y.), K/A, Geochron Laboratories, for Calif. Div. Mines and Geology, Oct. 1963.
- The age of some of these rocks is uncertain and actually may be younger than Precambrian.
- Radiometric dates of granitic rocks from the following area indicate:
 Marble Mountains (southern part)
 1450 million years (± 20 m.y.), Pb²⁰⁶/U²³⁸, Silver, L. T. and McKinney, C. R., 1963, U/Pb isotopic age studies of a Precambrian granite, Marble Mountains: Geol. Soc. Amer. Spec. Papers, No. 73 (1962 meeting), p. 65 (abstract).
 1250 million years (± 20 m.y.), K/A, and 1300 million years (± 30 m.y.), Rb-Sr, Lanphere, M. A., 1964, Geochronologic studies in the eastern Mojave Desert, California: Jour. Geol., in press.
 1190 million years (± 20 m.y.), K/A, and 1215 million years (± 30 m.y.), Rb-Sr, Lanphere, M. A., 1964, Geochronologic studies in the eastern Mojave Desert, California: Jour. Geol., in press.
 1150 million years (± 20 m.y.), K/A, and 1410 million years (± 30 m.y.), Rb-Sr, Lanphere, M. A., 1964, Geochronologic studies in the eastern Mojave Desert, California: Jour. Geol., in press.

TOPOGRAPHIC QUADRANGLES
 WITHIN THE NEEDLES SHEET
 AVAILABLE FROM THE U.S. GEOLOGICAL SURVEY
 FEDERAL CENTER, DENVER, COLORADO 80225
 1963



View northwest of Amboy Crater, a very recent cone composed of volcanic ejecta, surrounded by dark basaltic flow rocks. A thin veneer of wind blown sand gives the basalt a light appearance; thicker sand deposits are white. The Bristol Mountains, composed of dark Precambrian complex rocks and lighter granitic rocks are seen in the upper right, with the Granite Mountains on the skyline behind. The Lava Hills, in the upper left, are composed of Tertiary volcanic rocks underlain by Mesozoic granitic rocks and pre-Cenozoic granitic and metamorphic rocks. Photo by R. C. Frampton and J. S. Shelton, Claremont, California.

INDEX TO CALIFORNIA GRAVITY MAP SHEETS

The date beneath each map sheet name indicates year of publication. If no date appears, the gravity map sheet had not been published at the time this cover was issued.

