

CONFIDENTIAL  
CONFIDENTIAL

Union Oil Company of California

UCA05.1207

In Reply Give No.

PROPERTY OF UNION  
GEOTHERMAL DIV.  
GEOLOGY DEPARTMENT

UNION 76

CONFIDENTIAL

GMT 916-1

January 20, 1984

TO: Alex Schriener  
FM: Brian P. Hausback *Brian Hausback*  
RE: SURFICIAL GEOLOGY OF THE MEDICINE LAKE HIGHLAND

#### INTRODUCTION

Medicine Lake Highland is a youthful volcanic complex lying in northeastern California, 35 miles northeast of Mount Shasta (figure 1). The Highland lies in the portion of the Tertiary to Recent Cascade volcanic arc that stretches from British Columbia to Mt. Lassen in California. This younger portion of the Cascade range is a north to south-trending narrow line of volcanoclastic-dominated strato-volcanoes including such peaks as Mt. Shasta, Mt. St. Helens, and Mt. Rainier. The Medicine Lake Highland is one of three strikingly similar Quaternary lava flow-dominated shield-like volcanic systems lying on the east side of the chain of major strato-volcanoes. The other two are Newberry Volcano in Oregon and the Simco Volcanics in Washington.

The goal of this study was to produce a geologic map (plate 1) and report outlining the surficial geology of the central Medicine Lake Highland in order to guide future geothermal exploration in the area.

#### PREVIOUS STUDIES

The first comprehensive geologic study in the Medicine Lake Highland was by Charles Anderson (1941) who mapped the area in reconnaissance and outlined the general stratigraphy upon which all successive work has been based. Recently, there has been abundant interest in the area and the attached bibliography summarizes these works. Of particular interest to the surficial mapping are studies by Mertzman (1977b), several bachelor theses (Burnell, 1974; Hackett, 1974; Huffman, 1981; Hughes, 1974; Thomas, 1981; Walter, 1975; Weaver, 1976) completed at Franklin and Marshall College, Pennsylvania under the guidance of Mertzman, as well as studies by Fink (1983a).

Presently, Julie Donnelly-Nolan of the U. S. Geological Survey is mapping the Highland in detail but this work will not be available for a couple of years.

### REGIONAL GEOLOGY

Basement rocks are nowhere exposed in the Medicine Lake Highland. Regional outcrop and structural trends would suggest one of two basement types: Sierran granitic rocks or Klamath terrane rocks (metavolcanics, metasediments, ultramatics, serpentinites, granitics). The type of basement underlying the Highland is suggested by xenoliths found in the extrusive rocks. These include granitic and gneissic rocks suggesting a Sierran basement; no xenoliths were found to represent any of the distinctive Klamath Terrain suite of rocks.

The oldest rocks surficially exposed adjacent to the Medicine Lake Highland is the Cedarville Series (Anderson, 1941), a set of lavas and pyroclastics of late Tertiary age. Directly overlying the Cedarville Series is the Warner Basalt, a voluminous set of high-alumina olivine basalts spread over much of the Modoc Plateau region (Alturas Sheet, CDMG, 1958). One K-Ar date on a Warner Basalt by Thompson (1981) yielded about 1 m.y.. The Medicine Lake volcanic edifice is apparently built on the plateau of Warner Basalt. Neither the Cedarville Series nor Warner Basalt were encountered within the mapped area (plate 1).

Overlying the Warner Basalt-dominated Modoc Plateau is a belt of Pleistocene to Recent volcanic rocks stretching from Mt. Shasta to Medicine Lake. The transition between the stratovolcanic rocks of Mt. Shasta and the shield volcanic series at Medicine Lake has not been studied, hence the relationship between these two closely-spaced volcanic edifices is still not understood.

### STRATIGRAPHY OF THE MEDICINE LAKE HIGHLAND

#### General Statement

At 600 km<sup>3</sup>, the Medicine Lake extrusive system is the largest of the Cascade volcanoes, the next largest being its neighbor, Mt. Shasta (Julie Donnelly-Nolan, personal communication). Stratovolcanoes such as Mt. Shasta typically form steep-sided cones composed of intercalated pyroclastics and lavas of strongly porphyritic andesites and dacites (including minor rhyolite and basalt) with typical phenocryst assemblages of plagioclase, hornblende, pyroxene, quartz and sanidine. In

comparison, the Medicine Lake Highland is a flattened shield-shaped mountain, with a central basin morphologically reminiscent of the collapsed caldera at Oregon's Newberry Volcano and is composed almost entirely of lava flows with a complete range from rhyolite to basalt (figure 2). As opposed to the strato-volcano rocks, Medicine Lake intermediate-composition lavas contain fewer and much finer-grained phenocrysts. None of the Medicine Lake lavas contain quartz or hydrous phenocrysts (e. g. biotite, hornblende). The likely explanation for the limited phenocryst assemblages Medicine Lake is that the magmas were erupted at relatively high temperatures. The temperatures of the silicic lavas must have been higher than the quartz liquidus and hot enough to preclude the crystallization of hydrous phenocrysts, above about 870°C (Maaloe and Wyllie, 1975; Naney, 1982).

Anderson (1941) pointed out a general lack of basaltic rocks in the central Medicine Lake Highland. This central paucity of basalts has been noted at several large volcanic systems and is termed a basaltic shadow zone. It is generally attributed to the existence of a silicic, at least partially molten, magma chamber. Basaltic liquids cannot rise through molten or plastic rhyolitic magmas because of the large density contrast.

#### Hornblende Rhyolite Obsidian

The oldest unit on the periphery of the Medicine Lake Highland is a set of hornblende-bearing rhyolites. These rhyolites occur only on the west side of the mapped area (plate 1). On the southeast flank of Red Cap Mtn. this unit appears as dense, black, poorly-outcropping glassy porphyritic rhyolite (plagioclase + sanidine? hornblende) and has been dated at the summit of Red Cap Mtn. at  $1.01 \pm 0.05$  m.y. (Mertzman, 1982). On the southeast side of Dock Well there are two exposures of similar, but entirely devitrified, rhyolite with an age of  $1.18 \pm 0.06$  m.y. (Mertzman, 1982). These exposures may represent either old lava flows, domes or extremely shallow intrusive stocks. These silicic bodies are the only rocks in the Medicine Lake Highland found to contain hydrous phenocrysts and thus their genetic relationship to the Recent Medicine Lake magmatic system is probably remote.

#### Old Rhyolite, Dacite; Rampart Andesite and Basalt

Situated along the periphery of the Highland are a series of rhyolite, dacite, andesite and basalt exposures that appear to be the oldest lavas of Medicine Lake affinity. The Old Rhyolite and Dacite members are typically black glassy

flow-banded lavas, either aphyric or with minor plagioclase and pyroxene phenocrysts. Most exposures appear to be flows overlain by Shield Basalts and Andesites, however, the Old Dacite exposed southwest of Schonkin Spring is an elongate body with consistently-oriented internal foliations suggesting that it is a large dike-like body intruding a Shield Basalt exposure. K-Ar dates on the old rhyolites range from 0.61 to 0.33 m.y. (Mertzman, 1982; refer to Explanation Sheet of Plate 1).

The Shield Andesites and Basalts are typically younger than the Old Rhyolites and are exposed almost exclusively along the foot of the Highland. These basic to intermediate lavas have flow fronts with subdued to rounded topographic expressions compared with younger lavas. This is probably due to the subsequent deposition of the widespread Andesite Tuff, which created a topographic smoothing effect, covering-up craggy surfaces and filling-in topographic depressions.

#### Andesite Tuff

The Andesite Tuff directly overlies the older sequence of rhyolite to basalt flows. This distinctive gray-colored (red to orange when weathered) scoriaceous ash flow tuff covers a large area on the northwest flank of the volcano and is the only marker bed in the older Medicine Lake stratigraphy. It is a poorly exposed unit since it deposited principally in topographic depressions. In outcrop it breaks into large tabular blocks displaying densely welded to completely unwelded textures.

Anderson (1941) reported that the Andesite Tuff underlies the Medicine Lake volcanic stratigraphy, however, one small, possibly resurgent, exposure within the Medicine Lake depression indicates that the tuff was erupted from the Medicine Lake magmatic system. Examination of the size distribution of the maximum lithic clast incorporated in the tuff (figure 3) shows a systematic decrease in size to the northwest, indicating a northwest transport direction from the vicinity of Medicine Lake, where the maximum size lithic was encountered. Donnelly-Nolan (1983) estimated a rough volume of about 2 km<sup>3</sup> based on rather tenuous thickness control (she used an average thickness of 20 feet). This small volume estimate would not allow for much of a caldera collapse. However, the distribution of Rampart Andesite eruptive centers along an elliptical arc (plate 1) surrounding Medicine Lake (active immediately following the Andesite Tuff eruption and covering up any ring-fractures) indicates that there was a ring-fracturing event associated with the Andesite Tuff eruption, whether or not any measurable caldera collapse actually occurred.

## Rampart Andesites

The most common unit encountered in the Medicine Lake region is the crystal-poor Rampart Andesites. These voluminous flows issued from an elliptical set of vents surrounding Medicine Lake as well as several peripheral vents on the northerly and southerly slopes of the Highland. These flows directly overlie the Andesite Tuff and retain distinct flow morphology in areas that have not been glaciated. The flows compose a section of at least 120 feet thick along the northwestern Highland rim and it is probable that the true thickness is several times greater. Individual flows have distinctive platy-jointed interiors and massive to vesicular tops. The platy-joints are generally subhorizontal but commonly fold and swirl, suggesting that they follow flow-generated shear surfaces. The strike of these joints (and the fold axes of intraflow folds) are roughly perpendicular to the direction of local flow. Mertzman (1981, 1977) reports two K-Ar dates on these andesitic lavas of  $0.21 \pm 0.05$  m.y. and  $0.07 \pm 0.04$  m.y. These K-Ar dates analysed by Mertzman are suspect for several reasons:

1. Potassium analyses were attained from XRF procedures, a highly inaccurate, low precision method.
2. Sample splits were taken for potassium analysis prior to treating the remaining sample for argon analysis. Treating the K-split with different preparatory acids than the corresponding Ar-split introduces inhomogeneity into the sample.
3. Whole rock samples were heated to 250°C prior to argon extraction. Temperatures in excess of 100°C are well known to change the original argon isotopic abundances in whole rock samples and give spurious dates, some too old, some too young (McDougall, et al., 1976; Robert Drake, personal communication).

## Post Rampart Andesite Eruptive Units

Following, and in part overlapping with, the voluminous Rampart Andesite eruptions, the Medicine Lake magmatic system substantially diversified, producing numerous flows of various lithologies and textures. These flows are primarily aligned along three major linear zones on the Highland (discussed in the Structure section).

This post Andesite Tuff eruptive sequence is primarily composed of basalt through basaltic andesite with several voluminous rhyolitic to dacitic lavas. There appears to be a compositional gap at the intermediate composition series-andesite (see Correlation Chart with explanation to plate 1). Several of these younger units will be discussed here; for more comprehensive descriptions see the explanation to plate 1.

#### Lake Basalt

Three notable exceptions to the "basaltic shadow zone" in the central Highland are the young Lake Basalt and the Modoc basalts at Little Mt. Hoffman and the southwest foot of Mt. Hoffman. These occurrences cast doubt on the existence of a large Holocene silicic molten chamber beneath Medicine Lake.


The Lake Basalt is a voluminous sequence of olivine basalt flows that erupted primarily from the area of Lyons Peak with two eruptive cones to the west within the Medicine Lake basin itself. The highest occurrence of the basalt is on the southwest flank of Lyons Peak but the original eruptive cones are covered by a younger sequence of basaltic andesites. The flows are draped over the eastern rim of the basin and flowed far down the flank of the Highland.

#### Older Modoc Lavas

Subsequent to the Lake Basalt a sequence of medium-grained diktytaxitic high-alumina basalts vented from numerous cones situated primarily on the flanks of the Highland and concentrated in the northeastern sector. Most of these eruptive vents retain a very sharp cone morphology since they largely postdate the Pleistocene glaciation. Texturally and chemically these basalts are indistinguishable from the older Warner Basalt series.

#### Fissure-Erupted Basaltic Andesite

An unusual eruptive event at Medicine Lake vented andesitic to basaltic tephra from a major fracture zone trending southwest from Lyons Peak. The resultant poorly sorted tephra unit mantles the topography for a mile around the fracture zone. Subsequent to the tephra eruption, and probably during the last stages of it, numerous violent explosions occurred along the fracture zone creating large (up to 200 feet deep) craters (figure 4). These explosions cut through the agglutinated tephra, locally ejecting blocks of underlying Lake Basalt up to 10 feet or more in average diameter (figure 5) out of the craters onto the surrounding surface.



R. 4 E. 121°30' 2140000 FEET R. 4 E. 428 430 431 25' 433 41°30' 625000m E

REVISIONS	union
	UNION OIL COMPANY OF CALIFORNIA - GEOTHERMAL DIVISION
	<b>GLASS MOUNTAIN PROSPECT</b>
	<b>GEOLOGIC MAP OF THE</b>
	<b>MEDICINE LAKE HIGHLAND, CALIFORNIA</b>
	BY BRIAN P. HAUSBACK
	MARCH 1984
	SISKIYOU COUNTY, CALIFORNIA
	Scale 1:62500 DRAWN: DANA WILLIS April, 1983

Original color map in  
File G-10-910-1

## Youngest Bimodal Volcanism

The youngest eruptive phase at Medicine Lake consists of a bimodal eruption sequence of obsidian flows and Young Modoc basalts and basaltic andesites along the Little Glass Mountain and Glass Mountain structural trends. The basic members of this sequence are confined to the periphery of the Highland.

The youngest of these units are high silica rhyolite obsidian flows at Glass Mountain and Little Glass Mountain (both with several smaller satellite vents). The pumice tephra deposits associated with the obsidian flows were dated at about 1000 years by <sup>14</sup>C methods (Chesterman, 1955; Luedke and Smith, 1981). Based on tephra chronologic studies by Heiken (1981), Glass Mountain is the youngest unit and immediately postdates the eruption of Little Glass Mountain which, in turn, postdates basaltic eruptions probably from Cinder Butte and/or High Hole Crater.

Along the Glass Mountain structural trend are two distinct parallel lines of obsidian domes and flows. The southwestern line of vents is slightly older than the northeastern line of rhyolite vents, (including Glass Mountain) and is composed entirely of dacites, including the Hoffman Dacite (Dy).

## STRUCTURE

### Medicine Lake Structural Setting

The Modoc Plateau is dominated by north-northwest trending Basin and Range normal faults. The Medicine Lake edifice covers most of these faults and thus, in large part, postdates their activity. However, these older structures continue to act as deep crustal zones of weakness allowing magma to vent along them. The relative ease of magma rise along these zones may be the reason for the higher temperatures of magma at Medicine Lake compared to Mt. Shasta to the west.

There are two main orientations of structures in the Medicine Lake Highland: northwest and northeast. The Glass Mountain structure trends about N20W and is defined by the alignment of vents from Black Mountain to the south through the parallel vents of Glass Mountain and the Hoffman Dacite flow, and extending northward through aligned Older Modoc cinder cones to the prominent Basin and Range Gillem Fault (figure 6) in the Lava Beds National Monument. The high relief of the abrupt eastern margin of the Medicine Lake Highland over the Modoc Plateau can be attributed to normal faulting along this structure occurring either before or during the Pleistocene eruptive period.



### Little Glass Mountain Structural Trend

The other major orientation of structures and alignment of volcanic vents is about N30E. There are two zones following this orientation: the Little Glass Mountain and Lyons Peak Structural Trends. These zones do not extend into the surrounding regional structural pattern, suggesting that their formation is peculiar to the stress field associated with magma rise at Medicine Lake.

The Little Glass Mountain Structural Trend is defined by alignment of obsidian domes, and other more basic eruptive units young ground fissures. Along this structural trend the sequence of formation is: basaltic vent eruptions, pumice tephra eruptions and formation of local explosion craters, ground fissuring, and finally, the extrusion of the obsidian domes (from the same vents that issued the pumice tephra). Much of this activity (tephra eruptions, ground fissuring and dome extrusions) was nearly synchronous and took place about 1000 years ago.

The ground fissures (figure 7) along this trend are believed by Fink (1983) to have formed from the near-surface approach of silicic dikes. The cracks have dilatational opening patterns up to 12 feet wide and greater than 30 feet deep and seem to be associated with local topographic bulges. The extrusion of obsidian domes directly from these fissures is strong evidence that these structures have formed from the upward-shouldering of silicic magma along this northeasterly trending zone.

### Lyons Peak Structural Trend

Parallel to the Little Glass Mountain structure is a set of ground fissures and prominent explosion craters southwest of Lyons Peak. This zone lacks any silicic extrusions but was the locus of a basalt flow along its southerly extension and the eruption of a tephra of andesitic basalt (detailed in stratigraphy section) on the southwest flank of Lyons Peak. This set of structures and associated eruptive units immediately predates the 1000 year old pumice tephra erupted from Glass Mountain.

### Other Structures

One other set of faults cuts east-west across Medicine Mountain on the south side of the Highland and to the west of Fourmile Hill on the north side of the Highland. These linear

topographic escarpments are too high to be flow fronts and do not have the arcuate morphology of glacial cirques. They tend to downdrop the outside of the central Highland possibly suggesting that they are resurgent structures associated with the central 'caldera'.

#### SURFICIAL ALTERATION

Surface exposures displaying hydrothermal alteration are rare in the Medicine Lake Highland. This is conceivably due to the masking-effect that the widespread pumice tephra eruptions had on the area 1000 years ago. Nevertheless, through this young mantle there are two exposure displaying intense argillic hydrothermal alteration. The first and most prominent is the Hot Spot located on the Glass Mountain Structural Trend just northwest of Glass Mountain (plate 1). This is the only active zone of mild fumarolic activity in the Highland. At this locality the 1000 year old pumicious soil is thoroughly altered to clays. In the largest fumarole at this site a temperature of 168°F was recorded on July 25, 1983 by extending a thermometer 12 feet down into the vent. The other area of surficially exposed argillic alteration (and local silicification) is the area between Schonkin Springs and Crystal Springs, immediately west of Medicine Lake. This area is not an active hydrothermal zone but there are numerous localized occurrences of argillically altered Andesite Tuff, Rampart Andesite, Shield Basalt and associated cinder deposits.

## REFERENCES

- Allen, C. C., Jercinovic, M. J., and Allen, J. S. B., 1982, Subglacial volcanism in north-central British Columbia and Iceland: *Journal of Geology*, V. 90, pp. 699-715.
- Anderson, Jr., A. T., 1974, Evidence for a picritic, volatile-rich magma beneath Mt. Shasta, California: *Journal of Petrology*, V. 15, part 2, pp. 243-67.
- Anderson, C. A., 1940, Hot Creek lava flow: *American Journal of Science*, V. 238, No. 7, pp. 447-92.
- \_\_\_\_\_, 1941, Volcanoes of the Medicine Lake highland, California: *University of California Publications Bulletin of the Department of Geological Sciences*; Louderback, G. D. et al, eds., V. 25, No. 7, pp. 347-422.
- Barsky, C. K., 1975, Geochemistry of basalts and andesites from the Medicine Lake Highland, California: Ph.D. dissertation (unpublished), Washington University.
- Broker, M. M., Christopherson, K., and Haller, R., 1983, E-field ratio telluric survey near Medicine Lake in the Medicine Lake Highlands Caldera, Siskiyou County, California: U. S. Geological Survey Open File Report 82-900, 10 p.
- Brown, L., and Mertzman, S. A., 1979, Negative inclination anomalies from the Medicine Lake Highland lavas, northern California: *Earth and Planetary Science Letters*, V. 42, pp. 121-126.
- Burnell Jr., J. R., 1974, Geology and Petrology of the southeast portion of the Medicine Lake Highland, California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 44 p.
- Calex, Model 70 Fluxgate Magnetometer, Instruction Manual.
- Carmichael, I. S. E., 1967, The mineralogy of Thingmuli, a Tertiary volcano in eastern Iceland: *The American Mineralogist*, V. 52, pp. 1815-1841.
- Champion, D. E., Dalrymple, G. B., and Kantz, M. A., 1981, Radiometric and paleomagnetic evidence for the Emperor reversed polarity event at  $0.46 \pm 0.05$  M.Y. in basalt lava flows from the eastern Snake River Plain, Idaho; *Geophysical Research Letters*, V. 8, No. 10, pp. 1055-1058.

Chesterman, C. W., 1955, Age of the obsidian flow at Glass Mountain, Siskiyou County, California: American Journal of Science, V. 253, pp. 418-24.

Christiansen, R. L., 1982, Volcanic hazard potential in the California Cascades: in status of volcanic prediction and emergency response capabilities in volcanic hazard zones of California, Martin, R. C. and Davis, J. F., eds.: California Division of Mines and Geology, Special Publication 63, pp. 41-59.

Christiansen, R. L. and Lipman, P. W., 1966, Emplacement and thermal history of a rhyolite lava flow near Fortymile Canyon, southern Nevada: Geological Society of America Bulletin, V. 77, pp. 671-84.

Christiansen, R. L., and Lipman, P. W., Carr, W. J., Beyers, Jr., F. M., Orkild, P. P., and Sargent, K. A., 1977, Timber Mountain-Oasis Valley caldera complex of southern Nevada: Geological Society of America Bulletin, V. 88, pp. 943-59.

Condie, K. C., and Hayslip, D. L., 1975, Young bimodal volcanism at Medicine Lake volcanic center, northern California: Geochimica et Cosmochimica Acta, V. 39, pp. 1165-78.

Couch, R., Gemperle, M., 1982, Aeromagnetic measurements in the Cascade Range and Modoc Plateau of northern California -- Report on work done from December 1, 1980 to May 31, 1981: U. S. Geological Survey Open File Report 82-933, 31 p.

\_\_\_\_\_, 1982, Aeromagnetic measurements in the Cascade Range and Modoc Plateau of northern California -- Report on work done from June 1, 1980, to November 30, 1980: U. S. Geological Survey Open File Report 82-932.

Crandell, D. R., 1972, Glaciation near Lassen Park, northern California: U. S. Geological Survey Professional Paper 800-C, pp. C179-C189.

Cummings, D., 1964, Eddies as indicators of local flow direction in rhyolite: U. S. Geological Survey Professional Paper 475-D, Article 134, pp. D70-D72.

Davies, K. D., Vassell, R. K., Miles, R. C., Foley, M. G., and Bonis, S. B., 1978 Fluvial transport and downstream sediment modifications in an active volcanic region: in Fluvial Sedimentology; Miall, A. D., ed., Canadian Society of Petroleum Geology Memoir 5, pp. 61-84.

- Day, A. L., and Allen, E. T., 1925, The hot springs of Lassen National Park: Carnegie Institution of Washington, No. 360, pp. 85-123 (?).
- Doell, R. R., and Cox, A., 1962, Determination of the magnetic polarity of rock samples in the field: U. S. Geological Survey Professional Paper 450-D, Article 151, pp. D105-D108.
- Donnelly-Nolan, J. M., 1983, Two ash flow tuffs and the stratigraphy of Medicine Lake volcano, northern California Cascades: Geological Society of America Bulletin, Abstracts with Programs, V. 15, No. 5, p. 330.
- Duffield, W. A., and Fournier, R. O., 1974, Reconnaissance study of the geothermal resources of Modoc County, California: U. S. Geological Survey, Open File Report.
- Eaton, J. P., 1966, Crustal structure in northern and central California from seismic evidence: in Geology of northern California Division of Mines and Geology, Bulletin 190, pp. 419-26.
- Eichelberger, J. C., 1974, Origin of andesite and dacite: the petrographic and chemical evidence for volcanic contamination: Ph.D. dissertation (unpublished), Stanford University.
- \_\_\_\_\_, 1975, Origin of andesite and dacite: Evidence of mixing at Glass Mountain in California and at other circum-Pacific volcanoes: Geological Society of America Bulletin, V. 86, pp. 1381-91.
- \_\_\_\_\_, 1981, Mechanism of magma mixing at Glass Mountain, Medicine Lake Highland volcano, California: in Guides to some Volcanic Terranes in Washington, Idaho, Oregon, and northern California, U. S. Geological Survey Circular 838, pp. 183-189.
- Eichelberger, J. C., and Gooley, R., 1977, Evolution of silicic magma chambers and their relationship to basaltic volcanism; American Geophysical Union, Monograph 20, The Earths Crust, Heathcock, J. G., ed., pp. 57-77.
- Elston, W. E., and Smith, E. I., 1970, Determination of flow direction of rhyolitic ash-flow tuffs from fludial textures: Geological Society of America Bulletin, V. 81, pp. 3393-3406.

Fink, J. H., 1980, Gravity instability in the Holocene Big and Little Glass Mountain rhyolitic obsidian flows, northern California: Tectonophysics, V. 66, pp. 147-166.

\_\_\_\_\_, 1980, Surface folding and viscosity of rhyolite flows: Geology, V. 8, pp. 250-254.

\_\_\_\_\_, 1981, Surface structures of Little Glass Mountain: in Guides to some Volcanic Terranes in Washington, Idaho, Oregon, and northern California, U. S. Geological Survey Circular 838, pp. 183-189.

\_\_\_\_\_, 1983a, Structure and emplacement of a rhyolitic obsidian flow: Little Glass Mountain, Medicine Lake Highland, northern California: Geological Society of America Bulletin, V. 94, pp. 362-380.

\_\_\_\_\_, 1983b, Structural evidence for dikes beneath silicic domes, Medicine Lake Highland Volcano, California: Geology, V. 11, pp. 458-461.

Fink, J. H., Pollard, D. D., and Donnelly-Noian, J., 1982, Structural relationships among feeder dikes, silicic domes and ground cracks in the Medicine Lake Highland and Mount St. Helens Volcanoes (abstract): EOS, V. 63, No. 45, p. 1143.

Finn, C., 1982, A magnetic study of Medicine Lake volcano, California (abstract): EOS, V. 63, No. 45, p. 1150.

Finn, C., and Williams, D. L., 1982, Gravity evidence for a shallow intrusion under Medicine Lake volcano, California: Geology, V. 10, pp. 503-507.

Froggatt, P. C., 1982, Review of methods of estimating rhyolitic tephra volumes; application to the Taupo Volcanic Zone, New Zealand: Journal of Volcanology and Geothermal Research, V. 14, pp. 301-318.

Gardner, M. C., 1964, Cenozoic volcanism in the High Cascade and Modoc Plateau provinces of northeast California: Ph.D. dissertation (unpublished), University of Arizona.

Gerlach, D. C., and Grove, T. L., 1982, Petrology of Medicine Lake Highland volcanics: Characterization of endmembers of magma mixing: Contributions to Mineralogy and Petrology, V. 80, pp. 147-159.

- Grove, T. L., Gerlach, D. C., and Sands, T. W., 1982, Origin of Calc-Alkaline series lavas at Medicine Lake volcano by fractionation, assimilation and mixing: Contributions to Mineralogy and Petrology, V. 80, pp. 160-182.
- Grove, T. L., Gerlach, D. C., Sands, T. W., and Baker, M. B., Origin of Calc-Alkaline series of lavas at Medicine Lake volcano by fractionation, assimilation and mixing: Connections and clarifications: Personal communication.
- Hackett, W. R., 1974, Geology and petrology of a portion of the Medicine Lake highland, northern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 34 p.
- Hannah, J. C., 1977, Tectonic setting of the Modoc region, northeastern California: California Division of Mines and Geology, Special Report 129, pp. 35-39.
- Hart, W. K., 1976, The geology and petrology of the northern one-half of the Bray Quadrangle, northeastern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 71 p.
- Heiken, G., 1978, Petrologic evidence for a small magma chamber below the Medicine Lake highland, California: Geological Society of America Abstracts with Programs, V. 10, No. 3, pp. 108-109.
- \_\_\_\_\_, 1981, Holocene Plinian tephra deposits of the Medicine Lake Highland, California: in Guides to some Volcanic Terranes in Washington, Idaho, Oregon and northern California, U. S. Geological Survey Circular 838, pp. 177-181.
- Higgins, M. W., 1973, Petrology of Newberry Volcano, central Oregon: Geological Society of America Bulletin, V. 84, pp. 455-488.
- Higgins, M. W., and Waters, A. C., 1970, A re-evaluation of basalt-obsidian relations at East Lake Fissure, Newberry Caldera, Oregon: Geological Society of American Bulletin, V. 81, pp. 2835-2842.
- Hotchkiss, W. R., 1968, A geologic and hydrologic reconnaissance of Lava Beds National Monument and vicinity, California: U. S. Geological Survey, Water Resources Division, Open File Report, 30 p.

- Huffman, A. R., 1981, The geology, petrology, and geochemistry of the northwestern Humboldt and southwestern Medicine Lake Quadrangle, northern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 72 p.
- Hughes, J. M., 1974, Geology and petrology of the western Medicine Lake highlands, and Garner Mountain area, northern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 81 p.
- James, D. E., 1966, Geology and rock magnetism of Cinder Cove Lava Flows, Lassen Volcanic National Park, California: Geological Survey of America Bulletin, V. 77, pp. 303-12.
- Kane, P., 1982, Pleistocene glaciation, Lassen Volcanic National Park: California Geology, V. 35, No. 5, pp. 95-105.
- Kilbourne, R. T., and Anderson, C. L., 1981, Volcanic history and "active" volcanism in California: California Geology, V. 34, No. 8, pp. 159-168.
- Laux, G. M., 1970, Geology of the Modoc National Forest: Unpublished manuscript, Modoc National Forest.
- Luedke, R. G. and Smith, R. L., 1981 Map showing distribution, composition, and age of late Cenozoic volcanic centers in California and Nevada: U.S. Geological Survey, Miscellaneous Investigation Series, Map I-1091-C, scale 1:1,000,000.
- \_\_\_\_\_, 1982, Map showing distribution, composition, and age of late Cenozoic volcanic centers in Oregon and Washington: U.S. Geological Survey, Miscellaneous Investigation Series, Map I-1091-D, scale 1:1,000,000.
- Lydon, P. A., Gay, T. E., and Jennings, C. W., compilers, 1960, Geological map of California, Westwood Sheet, scale 1:250,000: California Division of Mines and Geology.
- MacDonald, G. A., 1966, Geology of the Cascade Range and Modoc Plateau: in Geology of northern California, California Division of Mines and Geology, Bulletin 190, pp. 65-190.
- MacDonald, G. A., and Katsura, T., 1965, Eruption of Lassen Peak, Cascade Range, California, in 1915; Example of mixed magmas: Geological Society of America Bulletin, V. 76, pp. 475-482.



- McBirney, A. R., 1978, Volcanic evolution of the Cascade Range: Annual Review of Earth and Planetary Science, V. 6, pp. 437-456.
- McDougall, I., Watkins, N. D., and Kristiansson, L., 1976, Geochronology and paleomagnetism of a Miocene-Pliocene lava sequence at Bessastadaa, Eastern Iceland: Amer. J. Sci., v. 276, p. 1078-1095.
- Maaloe, S. and P. J. Wyllie, 1975, Water content of a granite magma deduced from the sequence of crystallization determined experimentally with water-undersaturated conditions: Contrib. Mineral Petrol., v. 52, p. 175-191.
- Mahood, G. A., 1980, Geological evolution of a Pleistocene rhyolitic center-Sierra la Primavera, Jalisco, Mexico: Journal of Volcanology and Geothermal Research, V. 8, pp. 199-230.
- Mertzman, Jr., S. A., 1977a, Recent volcanism at Schonchin and Cinder Buttes, northern California: Contributions to Mineralogy and Petrology, V. 61, pp. 231-243.
- \_\_\_\_\_, 1977b, The petrology and geochemistry of the Medicine Lake Volcano, California: Contributions to Mineralogy and Petrology, V. 62, pp. 221-247.
- \_\_\_\_\_, 1981, Pre-Holocene silicic volcanism on the northern and western margins of the Medicine Lake Highland, California: in Washington, Idaho, Oregon, and northern California; U. S. Geological Survey Circular 838, pp. 163-169.
- \_\_\_\_\_, 1982, K-Ar results for silicic volcanics from the Medicine Lake Highland, northeastern California - A summary: Isochron/West, No. 34, pp. 3-7.
- Naney, M. T., 1982, Phase equilibria of rock-forming ferromagnesian silicates in granitic systems: Am. J. Sci., in press.
- Noble, D. C., Drake, J. C., and Whallon, M. K., 1969, Some preliminary observations on compositional variations within the pumice- and scoria-flow deposits of Mount Mazama: in Proceedings of the Andesite Conference, McBirney, A. R. ed.; Oregon State Department of Geology and Mineral Industries, Bulletin 65, pp. 157-164.

- Pakiser, L. C., 1964, Gravity, volcanism, and crustal structure in the southern Cascade Range, California: Geological Society of America Bulletin, V. 75, pp. 611-620.
- Peterman, A. E., Carmichael, I. S. E., and Smith, A. C., 1970, Sr<sup>87</sup>/Sr<sup>86</sup> Ratios of Quaternary lavas of the Cascade Range northern California: Geological Society of America Bulletin, V. 81, pp. 311-318.
- Powers, H. A., 1932, The lavas of the Modoc Lava Bed Quadrangle, California: The American Mineralogist, V. 17, No. 7, pp. 253-294.
- Reed, M., 1978, Chemistry of thermal waters in selected geothermal areas of California: California Division of Oil and Gas, Report TR15, pp. 26-31.
- Sans, J. R., 1972, Glass inclusions in olivines from Holocene cinder cones, Medicine Lake, California (abstract): American Geophysical Union Transactions, V. 53, p. 547.
- Schmid, R., 1981, Descriptive nomenclature and classification of pyroclastic deposits and fragments: Neues Jahrbuch Fuer Mineralogie Monatshefte, V. 4, pp. 190-196.
- Schriener, A., 1982, Geological discussion, Glass Mountain Unit Area: Union Oil Company, Geothermal Division, Geology Department, pp. 10.
- Self, S., and Rampino, M. R., 1981, The 1883 eruption Krakatau: Nature, V. 294, pp. 699-704.
- Smith, R. L., and Bailey, R. A., 1968, Resurgent cauldrons: in Coats, R. R., Hayes, R. L., and Anderson, C. A., eds., Studies in Volcanology, A memoir in honor of Howell Williams; Geological Society of America, Memoir 116, pp. 613-662.
- Sparks, R. S. J., Self, S., and Walker, G. P. L., 1973, Products of ignimbrite eruptions: Geology, V. 1, pp. 115-118.
- Sparks, S. J., 1975, Stratigraphy and geology of the ignimbrites of Vulsini Volcano, central Italy: Geol. Rundsch, V. 64, pp. 497-523.
- Spera, F. J., and Crisp, J. P., 1981, Eruption volume, periodicity, and caldera area: Relationships and inferences on development of compositional zonation in silicic magma chambers: Journal of Volcanology and Geothermal Research, V. 11, pp. 169-187.

- Taylor, S. R., Copp, A. C., and Graham, A. C., 1969, Trace element abundances in andesites; II. Saipan, Bougainville and Fiji: Contributions to Mineralogy and Petrology, V. 23, pp. 1-26.
- Thompson, K., 1981, Geology and petrology of the Bartle Quadrangle, northern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 86 p.
- U.S.G.S., 1981, U.S.G.S. projects in the Cascade Range, January 1981.
- Vantine, J., 1982, Comments on the hydrology of Glass Mountain and Medicine Lake KGRA based on information inferred from U.S.G.S. topographic maps: Union Oil Company, Geothermal Division, Santa Rosa.
- Vessell, R. K., and Davies, D. K., 1981, Nonmarine sedimentation in an active fore arc basin: Society of Economic Paleontologists and Mineralogists Special Publication No. 31, pp. 31-45.
- Walker, G. P. L., 1971, Grain-size characteristics of pyroclastic deposits: Journal of Geology, V. 79, pp. 696-714.
- Walter, R. C., 1975, Geology and petrology of the northwest portion of the Medicine Lake highland, California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 54 p.
- Weaver, S. G., 1976, Geology and petrology of the southeast portion of the Bray Quadrangle, northern California: B. A. Thesis, unpublished, Franklin and Marshall College, Pennsylvania, 67 p.
- Williams, H., 1935, Newberry volcano of central Oregon; Geological Society of America Bulletin, V. 46, pp. 253-304.
- \_\_\_\_\_, 1941, Calderas and their origin: University of California Publications, Bulletin of the Department of Geological Sciences, Louderback, G. D., ed.; V. 25, No. 6, pp. 239-346.
- Withjack, M. O., and Scheiner, C., 1982, Fault patterns associated with domes - an experimental and analytical study: American Association of Petroleum Geologists Bulletin, V. 66, n. 3, pp. 302-316.

Wohletz, K. H., and Sheridan M. F., 1983, Hydrovolcanic explosions II. Evolution of basaltic tuff rings and tuff cones: American Journal of Science, V. 283, pp. 385-413.

Wolff, J. A., and Wright, J. V., 1981, Rheamorphism of welded tuffs: Journal of Volcanology and Geothermal Research, V. 10, pp. 13-34.

Wright, J. V., Smith, A. L., and Self, S., 1980, A working terminology of pyroclastic deposits: Journal of Volcanology and Geothermal Research, V. 8, pp. 315-336.

## FIGURES

Frontpiece - View of snow covered Little Glass Mountain,  
Shasta in background

1. Location Map  
(from Christiansen, 1982, p. 42)
2. Topography of Northeast California  
(from Christiansen, 1982, p. 44)
3. Isopleth Map of Ave. Maximum Lithic Clast (Andesite Tuff)
4. Explosion Crater north of Undertakers Camp
5. Large Lake Basalt block ejected from explosion crater,  
SW side of Lyons Peak
6. View to North, Gillem Bluff, Lava Beds National  
Monument
7. Ground Fissures, Northeast of Little Mt. Hoffman

## PLATE

Plate 1 Geologic Map with Explanation of the Medicine  
Lake Highland, California

