

# UURI

EARTH SCIENCE LABORATORY  
391 CHIPETA WAY, SUITE C  
SALT LAKE CITY, UTAH 84108-1295  
TELEPHONE 801-524-3422

April 9, 1990

Randolph C. Thompson  
Unocal Geothermal Division  
3576 Unocal Place  
P.O. Box 6854  
Santa Rosa, California 95406

Dear Randy:

The lithologies in well 13 are similar to those previously studied in the "6", "8", and "17" series groups. However, unlike the other wells, no granodiorite or microgranodiorite is present in these samples from well 13.

Three major rock types are present. 1) Porphyritic to aphyric basalts and porphyritic basaltic andesites are commonly amygdular. 2) Coarser-grained rocks with abundant clino- and ortho- pyroxene are termed microdiabase, implying an intrusive origin for this rock type. Some of the microdabasases also contain minor amounts of biotite or hornblende. Phenocrysts in the coarser-grained diabases are optically-textured. 3) Quartz latites or rhyodacites exhibit a variety of devitrification textures and probably were originally glassy.

In general, predominantly aphyric and amygdular basaltic andesite flows are present to 1200'. However, at 300' and 800', porphyritic dacite is also present. The dacites probably represent flows; they are flow-banded and contain altered phenocrysts of biotite and perthitically intergrown feldspars in glomeroporphyritic aggregates. Angular fragments of oxidized andesitic basalt in sample 1200' may represent cinders that are part of a volcanic cone at the base of the predominately basaltic flow section.

Samples from 1300' and 1400' are mostly porphyritic andesite and represent a flow or possibly, a shallow intrusive body. Sample 1460' is a mixture of many rock types and represents the contact

between the andesite and an underlying thick (1500'-1800') rhyodacite. The cuttings in 1460' include basalt and andesite flow rocks and rhyodacite flows or tuffs. Because such a variety of lithologies are present in 1460' and not in the overlying section, it is likely that this depth represents an erosional disconformity in the volcanic section.

The rhyodacite displays several different textures. Relict perlitic texture is evident at the top (1500') and near the bottom (at 1800'), coarser-grained rhyodacite is present. This crystallization pattern suggests that the rhyodacite was deposited as a flow or a flow-dome complex; it may have cooled quickly to a glass near the surface (now 1500') and more slowly near its center (now 1800'). Spherulitic devitrification is also common in the center of the rhyodacite body. At the base of the rhyodacite, sample 1890' contains many cuttings with an argillic matrix and rounded andesite clasts along with resorbed quartz phenocrysts. This rock is probably an intraformational mudflow with an ashy matrix. One chip in sample 1890' is of coarse-grained graphic intergrowths of feldspar and quartz which could have been derived from a granitic highland.

Below 2000', the rocks consist predominately of diabase that is gouged or mylonitic and highly altered.

Samples from 2000' to 2200' are mostly microdiabase which probably represents a brecciated intrusive body. Within the microdiabase are altered phenocrysts of a ferromagnesian mineral, probably hornblende but possibly biotite, which are completely altered (deuterically?) to smectite. Clinopyroxene phenocrysts subophitically enclose plagioclase laths in the coarse-grained diabase at 2100'.

Samples from 2300' to 2400' are mostly hornblende microdiabase. The presence of potassium feldspar at 2300' is somewhat problematical. There is certainly secondary adularia in this sample but it is also possible that some of the potassium feldspar is primary. In the latter case and considering the presence of hornblende, some of the lithologies at 2300' and 2400' may be termed porphyritic rhyodacite or quartz latite. Either way, samples 2300' and 2400' represent a different rock unit than from 2000' to 2200'.

Some of the microbreccia in microdiabase at 2300' and in andesite at 2370' may have been caused by hydrothermal brecciation; both samples contain rounded, hematitic fragments within a silicified matrix.

Sample 2450' is nearly identical to sample 2370', and sample 2400' is similar to sample 2300'. Samples 2370' and 2450' are vesicular basaltic andesites and both are intensely mineralized. This phenomenon may indicate that vesicular lithologies provide good permeability pathways for hydrothermal fluids.

Samples 2500' and 2560' contain two rock types; fine-grained, glassy andesite and coarser-grained diabase, both of which are probably just textural variations of the same rock. In sample 2500', both rock types are vesicular and the vesicles are filled with fluid inclusion-rich (hydrothermal) quartz and/or epidote. The glassy, aphyric andesite probably represents the cooled borders of the intrusive body. In chips of diabase at 2500', the clinopyroxene is coarse-grained and subophitically-encloses plagioclase.

Samples 2560' and 2600' are both intensely altered and mineralized, making the original rock type difficult to determine. However, these samples appear to be mostly microdiabase (flooded with secondary K-feldspar?). Sample 2560' also contains some porphyritic rhyodacite; large quartz and zoned plagioclase phenocrysts are present in a devitrified matrix of quartz and potassium feldspar. Some fragments of the rhyodacite appear sericitized. Relict shard and pumice texture is evident in some chips indicating that the rhyodacite is at least partially tuffaceous.

The rocks from well 13 are mineralogically zoned with depth. Argillic alteration predominates down to about 1000'. Smectite and zeolite-filled amygdules are common in the argillic zone, although some calcite-filled amygdules are present at 600' and 800'. X-ray diffraction analysis of the zeolite indicates that it is stilbite, a low-temperature, fibrous zeolite.

The first occurrence of hydrothermal veining (chlorite-smectite and quartz+pyrite) is at 1000'. The top of the propylitic zone is at 1100', where epidote first appears as a trace mineral in veins. Veins of intergrown quartz and calcite (+/- epidote and wairakite) in samples from 1100' to 1460' suggest that this may be a zone of extensive boiling. Dark green, highly birefringent smectite is common in this zone and is especially abundant in sample 1400'. All of the lithologies in sample 1460' are highly altered and it is likely that this interval is a lithologic contact that was permeable and promoted hydrothermal alteration.

In general, the rhyodacite flow from 1500' to 1890' is not as altered as the overlying and underlying rocks. Chips of andesitic or diabasic rocks within the rhyodacite interval commonly contain veins and much epidote but the rhyodacite itself is commonly just silicified. Some chips of the (originally glassier?) rhyodacite are also sericitized. X-ray diffraction analyses of the clay fractions from samples 1600', 1710' and 1890' indicate the presence of mixed-layer illite-smectite with about 10% smectite interlayers. Such illitic illite-smectites usually indicate temperatures up to 260°C (e.g. McDowell and Elders, 1980).

The microdiabases from 2000' to 2300' are brecciated and veined. In thin-section, sheared and microbrecciated or strained crystals in a dark chloritic matrix characterize mylonitic chips of diabase. Traces of talc in this interval may be associated with

the fault gouge. Actinolite is present in this interval both in veins with hydrothermal quartz and as a (deuteric?) alteration product of primary clinopyroxene and/or amphibole in the diabase.

Dark green smectite is also a common alteration mineral in the interval from 2000' to 2300'. At 2200', this smectite occurs in veins with actinolite and quartz. Therefore, it must be a high-temperature variety of smectite (e.g. Eberl, Whitney and Khoury, 1978). From the X-ray diffraction pattern, it is not possible to distinguish this clay as saponite (magnesium-rich smectite) or a smectite-rich chlorite-smectite. Nevertheless, it appears to be identical in XRD and in thin-section to the clay in the lower portions of well 8 (termed smectite) and well 17 (termed chlorite-smectite). In well 13, the smectite is mostly absent in the hottest part of the well (below 2300') and a very chloritic mixed-layer chlorite-smectite is the only clay mineral that persists to the bottom of the well.

From 2300' down to 2600', well-defined hydrothermal veins of prehnite+epidote and adularia+quartz (+/- epidote and actinolite) are common. Evidence for hydrothermal brecciation at 2370' and 2450' suggests that some boiling may have occurred locally. The presence of veins of anhydrite at 2370' and 2470' may indicate increased permeability due to hydrothermal brecciation and boiling.

Potassium feldspar flooding of the diabase below 2300' is extensive. Because potassic alteration is typical of ascending and cooling solutions, the formation of K-feldspar in this zone may indicate the positioning of these rocks in or close to a major upflow zone (e.g. Giggenbach, 1984).

The secondary mineral assemblage from 2300' to 2600' is certainly indicative of deposition from high-temperature hydrothermal fluids. Epidote commonly forms above 240°C; prehnite above 250°C; and actinolite at temperatures greater than 250°C (e.g. Browne, 1978, 1984; Hulen and Nielson, 1986; Bird et al., 1984). Traces of grossular garnet in a vein with prehnite and epidote at 2560' may indicate temperatures above 300°C in this zone.

To conclude, the rocks in the lower portion of well 13 contain an abundance of unambiguous high-temperature hydrothermal minerals (particularly prehnite) in well-defined veins. The secondary (and perhaps metamorphic) mineral assemblage of biotite, actinolite and clinopyroxene which complicated interpretation of the hydrothermal alteration in the granodiorite of well 17 is not present.

At this point in your investigation, you should have the basis of a stratigraphic and structural framework for this area. If you don't and are willing to provide me with the thin-sections from, the depths of samples from, and the relative position of, wells 6, 8, 13 and 17, I would be very interested in working on this aspect of your investigation. Please contact me if you are interested in this proposal or if you have any questions concerning the X-ray diffraction or petrographic analyses of the rocks from well 13. Thanks for the opportunity to continue working on these interesting rocks.

Sincerely,

*Susan Juch Lutz*

Susan Juch Lutz  
Manager,  
X-ray Diffraction Lab

References:

- Bird, D.K., Schiffman, P., Elders, W.A., Williams, A.E., and McDowell, S.D., 1984, Calc-silicate mineralization in active geothermal systems: *Econ. Geol.*, v. 79, p. 671-695.
- Browne, P.R.L., 1978, Hydrothermal alteration in active geothermal fields: *Ann. Rev. Earth Planet. Sci.*, v.6, p 229-250.
- Browne, P.R.L., 1984, Lectures on geothermal geology and petrology: United Nations Univ., Geoth. Training Prog., Rept. 1984-2, 92p.
- Eberl, D., Whitney, G. and Khoury, H., 1978, Hydrothermal reactivity of smectite: *American Mineralogist*, v. 63, p. 401-409.
- Giggenbach, W.F., 1984, Mass transfer in hydrothermal alteration systems- a conceptual approach: *Geochim. et Cosmochim. Acta*, v. 48, p. 2693-2711.
- Hulen, J.B. and Nielson, D.L., 1986, Hydrothermal alteration in the Baca geothermal system, Redondo dome, Valles caldera, New Mexico: *J. Geophys. Res.*, v. 91, p. 1867-1886.

UNOCAL GEOTHERMAL DIVISION - RANDOLPH THOMPSON

MINERALOGY, APPROX. WT.%  (or) RELATIVE ABUNDANCE

WELL 13  
MEDICINE LAKE  
BUCK + CLAY XRD  
SAMPLE NO.

SAMPLE NO.	MINERALOGY										CEMENT W/REPUTE										
	QUARTZ	PLAGIOCLASE	K-FELDSPAR	ANHYDRITE	MICA	HEMATITE	PURITE	SPH. KUTTE	LEUCOENE UNOV	CHALCOPHOS		AMPHIBOLE	EPIDOTE	PREHNITE	WAIKAKITE	TALC	SMECTITE	MIXED-LAYER CLAY	MIXED-LAYER CLAY-SMECTITE	LITE-SMECTITE	TRACED BY XRD
1800 BUCK	2	14	19	32	3	3	2	3	1	TR	TR	4			9	4				8	CEMENT W/REPUTE
1890 BUCK	2	20	14	30	4	3	4	7		TR						9	3				
CLAY															TR	46	54				
2000 BUCK	2	30	9	4	9	2	2	15	1	1					1	16					6
2100 BUCK	5	35	6	3	10	1	2	4	9	TR					1	10	4				10
CLAY															44	56					
2200 BUCK	2	45		3	15	2	3	10	TR	TR	TR				16	1					3
CLAY															2	44	54				
2300 BUCK	9	31	10	2	9	1	12	8	2	1	TR				1	9	5				
2370 BUCK	10	26	11		2	5	1	13		14	5						11				1
CLAY																	99	1			
2400 BUCK	6	45	11	2	10	1	4	7	1	1					8	1					3
2450 BUCK	10	23	6	3	3	1	TR	15		16	6						11				6
CLAY																	100				
2500 BUCK	5	40	11	1	4	3	10	6	2	8							10				
2560 BUCK	7	31	7	1	7	3	13	3	1	12	1						11	2			
CLAY																	91	4			
2600 (C) BUCK	10	26	5	1	4	1	17		5	15	4						9				8

MM = PREDOMINANT M = MAJOR m = MINOR Tr = TRACE ? = TENTATIVE IDENTIFICATION



**SUMMARY OF X-RAY DIFFRACTION ANALYSIS**  
UNIVERSITY OF UTAH RESEARCH INSTITUTE, EARTH SCIENCE LABORATORY

S. Lutz  
3-9-90



SAMPLE FOOTAGE	ROCK TYPES															
	BASALT ANDESITE	FORPHYRITIC ANDESITE	MICRODIORITE	RHYODIORITE + RTZ LATHES	FORPHYRITIC RHYODIORITE	MICRO GRANODIORITE	HORNBLANDITE MICROGRANODIORITE	DACITE	RHYOLITE ASH FLOW TUFF	VOUCANIC CONGLOMERATE	BOULDER MICROBRECCIA	VEIN FRAGS.				
160	a MM	m														
200	m	MM														FLOW-BANDED
300								MM								DEVITRIFIED, SOME PERLITIC TEX, ALTER. BIOTITE
400	a MM		TR													
600	a MM															
800								MM								MUCH ALTERED BIOTITE, SOME CPX + OPX, + FLOW-BANDED
1000	a M	m	m								m					
1100	a MM								TR	m						
1200	a MM									m						ANGULAR FRAGS. + HEM. = CINDERS? SPHERULITIC
1300		a MM		m						m						
1400	m	MM								m	m					ALTERED
1460	a m	M		M				m		m	m					DEVITRIFIED, ALSO RELECT SHARD TEX. PERLITIC TEX, SPHERULITIC DEVITR. EPIDOTE
1500			TR	MM							TR					
1600	a m		m	M						TR	TR					
1710			m	MM												SPHERULITIC DEVITR. CEMENT
1800	m		m	M							TR					COARSE GR. WICK + OPX CEMENT
1890	a M			M		TR?			M							ONE CLAST GRAPHIC FSPAR a - FILLED W/ QZ
2000			MM							M	m					CPX > BIO BIOTITE -> SMECTITE
2100	TR		MM	TR						M	m					CPX + OPX >> BIO; COARSE RELECT PERLITIC TEX.
2200			MM							m	m					CPX + OPX >> BIO COARSE GR. = DIABASE
2300			MM							M	m					X CPX + HB. HYDROTHERMAL CROSS-CUTTING VEINS
2370	a MM									m	M					a - FILLED WITH QZ + EP HYDROTHERMAL BRECCIA
2400			MM							M	m					sim TO 2300 Mylonite
2450	a MM										M					a - MINERALIZED sim TO 2370
2500	a M		m							M	m					a - FILLED WITH QZ, EP SAME ROCK-DIFE TEX.
2560	m		M		m				m?	m	m					LARGE PLAG IN DEVITRIFIED MATRIX
2600	m		M							m	m					

a = ANHYDULAR

GRAPHIC F<sub>2</sub>

MORE XLN  
THAN 300'

MIXTURE  
OF ROCKS

ROUND  
ANDESITE  
CLASTS IN  
CLAYEN  
MATRIX

BRECCIA?

WELL 13  
MEDICINE LAKE

ROCK TYPES OBSERVED DURING  
RECONNAISSANCE PETROGRAPHY



SAMPLE FOOTAGE	VEINLETS & VEINLET FRAGMENTS													
	QTZ	QTZ + PY	QTZ + CC	ANH ± CC	CHL ± CC	SMECTITE	WAIR ± CC	EP ± CC	EPIDOTE	PREHNITE	KF ± QTZ		ACT ± QTZ	TALC
160														MAGN → HEM, GLASSY
200														SOME SMECTITE BIO → SMECT
300														SMECT, ZEOLITES IN AMYGDULES. MAGN → HEM
400														CC-FILLED AMYGDULES, HEM
600														ZEOLITE, CC, SMECT-FILLED AMYGDULES
800														FIRST PY, CHL
1000														* FIRST EP
1100	✓	✓ <sub>leuc</sub>	✓		✓	✓				✓ <sup>+</sup> QTZ				* BOILING ASSEMBLAGE? MAGN → HEM QTZ + WAIR IN AMYGDULES
1200	✓	✓	✓		✓			✓ <sup>+</sup> CC						EP + CHL + PY, LEUCOKENE
1300	✓	✓	✓		✓	✓		✓	✓					MUCH OK. GREEN SMECTITE, TR. CHALC. AMY
1400		✓	✓					✓						PLAG → EP + CHL, LEUCOKENE,
1460	✓		✓					✓ <sup>+</sup> QTZ						SILICIFIED, BIG WAIR VEINS
1500	✓				✓			✓ <sup>+</sup> QTZ						EP IN MICRADIABASE SILICIFIED ANHYDRATE
1600	✓	✓	✓		✓									"
1710	✓													"
1800	✓	✓							✓ <sup>+</sup> CHL					" LEUCOKENE
1890	✓	✓			✓			✓ <sup>+</sup> EP PY						SOME SERICITE
2000		✓			✓				✓ <sup>+</sup> QTZ			✓ <sup>+</sup> EP	✓	FIRST ACTINOLITE, TALC MUCH SMECTITE
2100	✓							✓ <sup>+</sup> QTZ	✓					
2200					✓			✓					✓ <sup>+</sup> ACT	ACT + QTZ + SMECTITE FIRST PREHNITE
2300	✓ <sub>leuc</sub>	✓	✓ <sup>+</sup> EP						✓ <sup>+</sup> QTZ	✓	✓		✓	* BOILING ASSEMBLAGE? KF FLOODING?
2370				✓ <sup>+</sup> EP	✓			✓	✓ <sup>+</sup> EP	✓ <sup>+</sup> EP				INTENSELY MINERALIZED, HEM
2400	✓								✓ <sup>+</sup> QTZ	✓ <sup>+</sup> ACT	✓ <sup>+</sup> EP			KF + ACT + EP VEINS KF FLOODING
2450				✓ <sup>+</sup> EP	✓				✓ <sup>+</sup> QTZ	✓ <sup>+</sup> EP	✓			MINERALIZED, HEM LEUCOKENE
2500		✓			✓				✓ <sup>+</sup> QTZ	✓ <sup>+</sup> EP	✓	✓ <sup>+</sup> EP		KF FLOODING
2560					✓				✓ <sup>+</sup> QTZ	✓ <sup>+</sup> EP		✓		LOTS OF ACTINOLITE + CHL IN DIABASE, SOME SERICITE
2600	✓				✓				✓ <sup>+</sup> QTZ	✓ <sup>+</sup> EP	✓ <sup>+</sup> PREH			HEM + LEUCOKENE KF FLOODING, MINERALIZED

TR. GARNET + PREHN + EP

WELL 13  
MEDICINE LAKE

VEINLETS OBSERVED DURING  
RECONNAISSANCE PETROGRAPHY