

Geol. Soc. of Amer. Vol. 7, No. 5 1975

6207182

that slight environmental differences greatly influenced the morphology of the stromatolites. Near Rio Blanco, Colorado, large domed stromatolites can be traced into thin algal laminites over a distance of a few hundred meters. The large domed stromatolites are interpreted as growing on a promontory where wave activity was significant. In protected areas where wave activity was less, the domed structures were replaced by thin algal laminites. Many of the domed structures are elongated perpendicular to the shore line.

Modern stromatolites from Shark Bay are found only where high salinity has eliminated grazing of algae by other organisms. The absence of gastropods and pelecypods associated with the stromatolite units in the Green River Formation probably implies that high salinity was a necessary condition for stromatolite development in Eocene Lakes Gosute and Uinta also.

✓ GEOPHYSICAL STUDIES OF A GEOTHERMAL AREA IN THE SOUTHERN RAFT RIVER VALLEY, IDAHO

Mabey, Don R., Ackermann, Hans, Zohdy, Adel A.R.,
Hoover, Donald B., Jackson, Dallas B., and
O'Donnell, James E., U.S. Geological Survey,
Denver Federal Center, Denver, Colorado 80225

The U.S. Geological Survey in cooperation with the Atomic Energy Commission has made gravity, magnetic, refraction seismic, resistivity, audio-magnetotelluric, self-potential, and telluric current surveys in a study of the geothermal resources in the southern Raft River Valley. The geophysical data indicate that the maximum thickness of Cenozoic sedimentary and volcanic rock underlying the valley is about 2 km and that the valley is bounded by normal faults on the east and south and by a complex system of faults on the west. Large gravity, magnetic, and total field resistivity highs within the valley east of Jim Sage Mountains reflect a mass at a relatively shallow depth, which is probably igneous rock but too old to relate directly to a geothermal system. The seismic interpretation divides the valley into four areas where the Cenozoic rocks have distinctive seismic velocities. These areas appear to relate to known or inferred structures and to a suspected zone of shallow warm water. Resistivity anomalies reflect compositional variations in the Cenozoic rocks and variation in degree of induration and alteration. Although no large reservoir of hot water in the Cenozoic rocks has been identified yet, the resistivity soundings show a 2-5 ohmmeter resistivity unit with a thickness of 1 km underlying a large area of the valley, which may in part be indicative of hot water. Observed self-potential anomalies are believed to indicate zones where warm water is ascending to near the surface. The warm water occurring in springs and wells in the area northeast of The Narrows may relate to deep circulation control by the intersection of north-trending faults east of the Jim Sage Mountains with a northeast-trending structure passing through The Narrows.

REGIONAL GRAVITY AND MAGNETIC STUDIES OF THE SNAKE RIVER PLAIN

Mabey, Don R., Peterson, Donald L., and Wilson, Carol W.,
U.S. Geological Survey, Denver Federal Center, Denver,
Colorado 80225

The western and eastern parts of the Snake River Plain are distinctly different, as shown by the gravity and magnetic anomalies. Over the western part gravity and magnetic highs are approximately coextensive

with t
and by
depth
Group.
magnet
the an
lying
appear
Superi
of the
appear
which
The Sn
the re
faults
plain
the ea
Miocen
Basin-
that r
in the
subsid

INTERN
ALLUVI
Mac
U
D
F

A smal
source
form c
alluvi
logic
bution
strati

The
size a
develo
of the
sed re
and sh
these

Hea
deposi
fan-he
degree
of the

Ano
mum te
from t

Bot
distri
alluvi
econom